

1 Synoptic drivers of the August 2024 record-breaking rainfall in the Chadian
2 Sahara: Dynamics, thermodynamics, and socio-economic consequences

3
4
5 Claudin Wamba Tchinda¹· Frédéric Saha^{2,3}· Alain T. Tamoffo⁴
6
7
8

9 ¹Climate Laboratory for Environmental Modeling and Atmospheric Physics (LEMAP), Faculty
10 of Science, University of Yaoundé 1, Yaoundé, Cameroon

11 ²Faculty of Arts, Letters and Social Sciences, University of Yaoundé 1, Yaoundé, Cameroon

12 ³Community Resilience Association (RESCOM-Tchad), Moundou, Chad

13 ⁴Climate Service Center Germany (GERICS), Helmholtz-Zentrum Hereon, Hamburg, Germany
14

15 Corresponding author: Claudin Wamba Tchinda wambatchindaclaudin@yahoo.fr
16

17 Claudin Wamba Tchinda's orcid: <https://orcid.org/0009-0007-5715-5897>

18 E-mail: wambatchindaclaudin@yahoo.fr
19

20 Frédéric Saha's ORCID: 0000-0003-0327-9417

21 fredericsaha72@gmail.com
22

23 Alain T. Tamoffo's ORCID: 0000-0001-8482-8881

24 alaintamoffotchio@gmail.com
25
26
27
28
29
30
31
32
33
34
35
36

37 Abstract

38 This study examines the synoptic atmospheric mechanisms behind the extreme rainfall
39 event of August 2024 in northern Chad and their devastating socio-economic impacts. Based on
40 the CHIRPS, TAMSAT, and ERA5 datasets (1983–2024), Theil-Sen and Mann-Kendall analyses
41 reveal significant increases, over several decades, in precipitation and integrated water vapor
42 transport. Crucially, a transition phase in the early 2000s shifted the regional baseline from
43 persistent late-20th-century deficits into a moister "new normal," thereby culminating in the
44 record-breaking August 2024 anomaly.

45 During this month, unprecedented accumulations reached 100–400 mm over the Tibesti, Borkou,
46 and Ennedi massifs, representing parametric return periods of 124.5 years in CHIRPS and 451.4
47 years in TAMSAT, with standardized anomalies exceeding $+8\sigma$ (locally $>+12\sigma$). Our analysis
48 of lower-tropospheric convergence, specific humidity, vertical velocity (ω), and moist static
49 energy (MSE) reveals a major shift from the typical West African monsoon regime, validated via
50 a 1000-iteration grid-point Monte Carlo permutation test ($p \leq 0.05$). In August 2024, the
51 Intertropical Front (ITF) defined by the 15°C dew point isodrosotherm at 925 hPa shifted
52 abnormally northward, reaching 20°N – 22°N , which allowed moist air to penetrate deep into the
53 Saharan zone. This shift was driven by strengthened convergence at 850 hPa, anomalous
54 cyclonic vorticity, and a significant increase in low-level humidity, where specific humidity and
55 Total Column Water Vapor (TCWV) reached their absolute historical records (97.67th empirical
56 percentile). Furthermore, negative ω anomalies throughout the troposphere indicate a northward
57 extension of the monsoon's upward branch. Strong positive MSE anomalies and equivalent
58 potential temperature exceeding the 2σ statistical threshold over desert regions further highlight
59 a thermodynamic enrichment of the atmospheric column. Together, these signals point to a
60 highly effective dynamic-thermodynamic coupling that fueled intense convective systems. The
61 synchronization between these atmospheric conditions and the synoptic forcing of African
62 Easterly Waves (AEWs) between August 10 and 14 generated extreme local rainfall anomalies,
63 redefining the hydrological balance between aquifer recharge and increased risks of flash
64 flooding.

65 This hydroclimatic shift caused devastating impacts: nationwide, floods affected 2 million people
66 and caused 576 fatalities. Regionally, impacts were severe and heterogeneous, affecting 67,613
67 individuals in Borkou with 23,174 houses destroyed, and claiming 60 lives in Tibesti. Beyond
68 livestock, crop, and gold-mining infrastructure losses, the floods unearthed thousands of
69 conflict-era landmines, creating a grave secondary hazard. These findings underscore hyper-arid
70 zones' vulnerability to extreme hydrological events, highlighting the urgent need to transition
71 from reactive disaster relief to proactive climate adaptation.

72

73 Keywords: Extreme rainfall; Chadian Sahara; flooding; Intertropical Front; Moist Static Energy;
74 west African monsoon; Socio-economic impacts

75

1. Introduction

Extreme precipitation episodes constitute one of the most critical manifestations of contemporary climate variability, posing a growing challenge to the resilience of human societies (Seneviratne et al., 2021). Their impact is particularly devastating in arid and semi-arid regions, where structural water scarcity renders natural and socio-economic systems extremely vulnerable to hydrometeorological anomalies (Saha et al., 2020). Under global warming, the intensification of hydrological cycle has led to a documented increase in the frequency and magnitude of these events, often exceeding local adaptive capacities (Allan and Soden, 2008; Trenberth et al., 2015; Ehtasham et al., 2024; Adeyeri, 2025). Specifically, surpassing critical warming thresholds could increase intense precipitation in Central Africa by up to 25%, exacerbating flooding risks and population exposure (Zakariah et al., 2024; Tamoffo et al., 2024a). This vulnerability is especially concerning for the Lake Chad Basin, where projections indicate a resurgence of rainfall extremes alongside an extension of dry spells (Adeyeri et al., 2019).

The Sahara and its southern margins operate under a climatic regime traditionally governed by tropospheric subsidence and erratic rainfall. In this environment, intense rain events arise from exceptional atmospheric configurations linked to the dynamics of the Saharan Heat Low (SHL) and regional moisture convergence (Selami et al., 2021). This dynamics is part of the rainfall recovery observed since the end of the 20th century, driven by oceanic warming and a strengthened monsoon moisture budget (Sindikubwabo et al., 2018; Biasutti, 2019; Tamoffo et al., 2026). However, paleoclimatic records serve as a reminder that this system is prone to non-linear and abrupt responses; the history of "Lake Mega-Chad" demonstrates that transitions between aridity and "African Humid Periods" can be sudden (Simon et al., 2015; Pausata et al., 2017; Pausata et al., 2020).

Northern Chad, a confluence point between the central Sahara and the Sahel, represents a highly sensitive climatic transition zone where the balance between water resources and agro-pastoral activities is precarious (Raimond et al., 2014; Mahamat Nour et al., 2017). Rainfall distribution is strictly governed by the latitudinal position of the Intertropical Front (ITF), the surface expression of the Intertropical Convergence Zone (ITCZ) marking the boundary between the dry harmattan and the moist monsoon (Nicholson, 2013). An anomalous northward migration of the ITF allows for the injection of Atlantic moisture into the heart of the Sahara, triggering deep convection (Sultan and Janicot, 2003; Parker et al., 2005). This dynamics is further modulated by intra-seasonal oscillations (ISO), which influence water vapor transport and moist static energy (MSE) fluxes (Siwe et al., 2025; Wamba et al., 2023; Tchinda et al., 2026; Tamoffo et al., 2024b) and is often exacerbated by synoptic perturbations, such as African Easterly Waves (AEW) (Kiladis et al., 2006; Lafore et al., 2017).

Beyond dynamics, thermodynamic and radiative factors are decisive. MSE offers a diagnostic framework integrating temperature, humidity, and the vertical structure of the atmosphere (Neelin and Held, 1987; Holloway and Neelin, 2009). High MSE values indicate a warmer and moister boundary layer, which directly enhances the convective available potential energy while

simultaneously reducing convective inhibition, facilitating the organization of mesoscale convective systems responsible for extreme accumulations (Romps, 2015). Recent studies demonstrate that these events result from a synergy between thermodynamic instability, radiative flux anomalies, and dynamic forcings such as moist enthalpy advection and potential vorticity anomalies (Kenfack et al., 2024, 2025; Tchana et al., 2025). Simultaneously, vertical velocity characterizes the strength of updraft, where negative ω anomalies represent the classic signature of organized convection in West Africa (Taylor et al., 2017).

At the global scale, 2024 was characterized by unprecedented warming and a surge in hydrometeorological extremes (Zhang et al., 2025). In August 2024, exceptionally intense rainfall hit the Saharan regions of Chad and the Lake Chad Basin, leading to catastrophic socio-economic impacts, including unusual flooding, infrastructure collapse, agricultural losses, and widespread population displacement (OCHA, 2024). This crisis highlighted the acute vulnerability of local socio-ecosystems, which are historically adapted to a hyper-arid regime with low mean annual rainfall. In such environments, even a minor absolute increase in precipitation represents a severe hydrometeorological hazard, triggering disproportionate societal disruptions due to the lack of infrastructure designed for torrential rain. While the 2024 anomaly aligns with a broader Sahelian trend associated with an anomalously northward ITCZ (Zhang et al., 2025; Nicholson, 2013), the specific atmospheric mechanisms driving rainfall in the true Saharan desert remain insufficiently documented. Most existing literature focuses on the Sahelian and Sudanian belts, leaving a gap in our understanding of the joint dynamic and thermodynamic processes and their socio-economic consequences within the hyper-arid Sahara and the Lake Chad Basin.

This study aims to fill this gap by characterizing the forcings behind the August 2024 extreme event through an integrated analysis of moisture convergence, remote moisture advection, vertical velocity, and MSE. By linking these physical mechanisms to field observations of socio-economic impacts, this work provides critical insights for climate adaptation and risk management in the Sahara and Lake Chad Basin (Biasutti, 2019; Vizy and Cook, 2022). The article is structured as follows: Section 2 details the study area, data and methodology; Section 3.1 characterizes the extreme rainfall event and its spatial-temporal evolution, while the rest of Section 3 explores the underlying dynamic and thermodynamic physical mechanisms; Section 4 details the associated socio-economic impacts and local vulnerabilities; and Section 5 concludes the study.

148

149 2. Study Area, Data and Methods

150 2.1. Study Area

This study focuses on northern Chad, extending 16°-24°N, 13°-25°E, an area administratively comprising the regions of Tibesti, Borkou, Ennedi West, and Ennedi East (Figure 1). The landscape is dominated by vast sandy plains and longitudinal dune systems. The regional climate is hyper-arid, characterized by high diurnal and seasonal thermal ranges and a mean annual precipitation below 50 mm (UNESCO, 1977; Nicholson, 2011). Despite its harsh

environment, the Saharan sector of Chad accounts for approximately 13% of the national population (approx. 2.73 million inhabitants; INSEED-Chad). Economic subsistence is primarily linked to oasis-based agriculture, specifically date palm cultivation, and nomadic pastoralism centered on camel herding. The primary urban centers include Faya-Largeau (pop. 100,000), Fada (pop. 50,000), Bardaï (pop 30,000) and Amdjarass (pop. 2,000). These cities, like all inhabited localities, are established in oases to take advantage of the meager water resources available, given the hyper-arid environment.




Figure 1. Study area in Chad. Panel A shows that this predominantly desert area is bordered to the south by the 100 mm isohyet. Panel B presents the four administrative provinces whose socioeconomic losses from the August 2024 floods were analyzed. It is worth noting that despite the desert environment, numerous settlements are located there, generally in oases.

167

168

169 2.2. Data sources

170 2.2.1. Satellite Precipitation Estimates

171 To analyze the long-term interannual variability, monotonic trends, and the specific
 172 magnitude of the August 2024 rainfall event, monthly precipitation datasets spanning the
 173 1983-2024 period were gathered from two distinct high-resolution satellite-based products: the
 174 Tropical Applications of Meteorology using SATellite Data v.3.1 (TAMSAT; Maidment et al.,
 175 2017) and the Climate Hazards Group InfraRed Precipitation with Station data v.2.0 (CHIRPS;
 176 Funk et al., 2015).

177 A critical constraint in the hyper-arid Borkou-Ennedi-Tibesti (BET) region of northern Chad is
178 the severe scarcity or total absence of operational, continuous in-situ rain gauges. Consequently,
179 hydro-climatic monitoring relies exclusively on satellite remote sensing. CHIRPS blends global
180 satellite infrared imagery ($0.05^\circ \times 0.05^\circ$ resolution) with available station observations to
181 generate long-term gridded time series, which is highly effective for drought and trend
182 monitoring (Funk et al., 2015). The dataset was accessed from the Climate Hazards Group
183 (<https://www.chc.ucsb.edu/data>). Concurrently, TAMSAT estimations are derived from Meteosat
184 Thermal Infrared (TIR) imagery, using a Cold Cloud Duration (CCD) threshold approach
185 calibrated against regional ground-based gauges (Maidment et al., 2017). TAMSAT features a
186 high spatial resolution of 0.0375° (~ 4 km), which is optimized for capturing convective rainfall
187 distributions across complex Saharan topography (Wainwright et al., 2021). Relying on both
188 products is necessary to address structural uncertainties inherent to satellite retrieval algorithms
189 over hyper-arid terrains where direct gauge validation is unavailable.

190

191 2.2.2. Atmospheric Reanalysis (ERA5)

192 Atmospheric dynamics, vertical coupling, and moisture transport patterns were analyzed
193 using the ERA5 reanalysis, the fifth-generation global atmospheric climate dataset developed by
194 the European Centre for Medium-Range Weather Forecasts (ECMWF) (Hersbach et al., 2020).
195 ERA5 employs a multi-decadal advanced data assimilation scheme, ensuring a physically
196 consistent co-dependence between dynamic and thermodynamic variables across data-sparse
197 regions such as the Sahara desert (Tarek et al., 2020; Ssenyunzi et al., 2020). Previous climate
198 assessments have confirmed that ERA5 accurately reproduces monsoon circulation, moisture
199 influxes, and synoptic-scale waves over the Sahelian and Saharan belts (Johannsen et al., 2019;
200 Cook and Vizzy, 2021).

201 For this study, data were extracted at a horizontal resolution of $0.25^\circ \times 0.25^\circ$ on standard
202 pressure levels spanning from the lower troposphere to the upper troposphere (850 hPa to 200
203 hPa). The monthly variables retrieved include the horizontal components of wind (u , v), vertical
204 velocity (ω), horizontal wind divergence (d), geopotential height (z), temperature (T), and
205 specific humidity (q). The analysis focuses on the month of August from 1983 to 2024. While
206 the total 42-year period is utilized for interannual trend estimations, the 1983-2023 window
207 ($N=41$ years) is strictly designated as the standard climatological baseline mean from which the
208 exceptional structural anomalies of August 2024 are evaluated.

209

210 2.2.3. Data on flood losses

211 Due to Chad's fragile socioeconomic and security context, systematic disaster recording
212 is primarily facilitated by international humanitarian organizations and United Nations agencies
213 in collaboration with the Chadian government. For the 2024 flood event, impact data were
214 obtained from the United Nations Office for the Coordination of Humanitarian Affairs (OCHA),
215 which synthesizes inter-agency field reports. These data are validated by the Ministry of

216 Territorial Administration and Decentralization, alongside the Ministry of Social Action,
217 Solidarity, and Humanitarian Affairs.

218 The dataset includes standardized metrics on:

- 219 • Human casualties (mortality and injury rates);
- 220 • Displaced populations and affected households;
- 221 • Agricultural losses (hectares of devastated cropland);
- 222 • Infrastructure damage (destruction of homes and commercial facilities);
- 223 • Livestock mortality.

224 While national summaries typically emphasize the southern and central provinces due to their
225 higher absolute population densities (e.g., Mayo-Kebbi Est, Lac, and Tandjilé), this study
226 specifically isolates data for the northern Saharan provinces. Although the absolute number of
227 victims in these arid regions is lower compared to the south, the relative impact, defined as the
228 ratio of affected individuals to the total provincial population, is among the highest recorded in
229 2024. This relative approach is essential for assessing the exceptional nature of climate extremes
230 in sparsely populated hyper-arid zones. Data was accessed via the OCHA Chad humanitarian
231 portal (<https://www.unocha.org/chad>).

232 2.3. Methodological Framework and Statistical Metrics

233 To diagnose the physical mechanisms driving the exceptional rainfall of August 2024, a
234 suite of diagnostic variables was employed. The methodological framework distinguishes
235 between thermodynamic precursors (the energy and moisture required for convection) and
236 dynamic forcing (the atmospheric mechanisms triggering vertical motion).

237 2.3.1. Climatological Anomalies and Spatial Significance

238 To highlight the physical anomalies driving the extreme event of August 2024, monthly
239 climatological anomalies at each grid point were computed as the direct departure of the
240 observed August 2024 values from the long-term historical mean, which was established over a
241 strict 41-year reference baseline period spanning from 1983 to 2023.

242 To assess whether these spatial anomalies represent a statistically significant departure from the
243 background noise or are merely a product of random internal climate variability, a
244 non-parametric Monte Carlo permutation test was systematically applied to the anomaly fields.
245 This approach is highly suitable for atmospheric dynamics over hyper-arid zones because it does
246 not rely on parametric assumptions of normal probability distributions (Livezey and Chen,
247 1983). For each variable, the historical time series (1983 - 2024) was randomly shuffled 1000
248 times to empirically reconstruct a local null distribution of climatological anomalies at each grid
249 point. The true observed August 2024 anomaly was then positioned within this empirical
250 distribution. Anomalies falling outside the 2.5th and 97.5th percentiles of the permuted
251 distribution ($p \leq 0.05$, two-tailed test) were diagnosed as statistically significant at the 95%

confidence level. These structurally robust grid points are explicitly highlighted via stippling on the corresponding spatial maps.

2.3.2. Long-Term Trend Analysis

The long-term monotonic trajectories of regional precipitation and column-integrated moisture transport over the 42 years (1983-2024) were quantified using the non-parametric Theil-Sen estimator, while trend significance was evaluated using the Mann-Kendall test. The Sen's slope represents the median of all slopes calculated between all possible pairs of data points in the time series, ensuring structural immunity to extreme outliers such as the August 2024 anomaly itself (Sen, 1968). To contextualize the physical magnitude of these trends, the relative change (Δ_{rel} , expressed in %) was calculated by multiplying the Sen's slope by the total length of the time series ($\Delta_t = 42$ years), normalized by the long-term climatological mean ($\bar{X}$) obtained from the 1983 - 2023 baseline:

$$\Delta_{rel} = \left(\frac{\text{Sen's Slope} \times \Delta_t}{\bar{X}} \right) \times 100 \quad (1)$$

2.3.3. Extreme hydro-climatic quantified diagnostics

To quantify the exact recurrence interval and rarity of the August 2024 precipitation, a parametric framework was implemented. Because precipitation over Saharan environments is highly intermittent, strictly non-negative, and positively skewed, standard Gaussian frameworks are physically invalid. Instead, a continuous two-parameter Gamma probability density function ($f(x)$) was fitted to the multi-decadal historical August series:

$$f(x, \alpha, \beta) = \frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-\frac{x}{\beta}} \text{ for } x > 0 \quad (2)$$

Where $\alpha > 0$ is the shape parameter, $\beta > 0$ is the scale parameter, and $\Gamma(\alpha)$ represents the standard Gamma function. These parameters were optimized via maximum likelihood estimation (MLE), providing asymptotically unbiased and numerically robust estimators for meteorological series (Husak et al., 2007). Once the cumulative distribution function ($F(x)$) was solved, the return period (T_r , in years) for the August 2024 rainfall magnitude (x_{2024}) was derived as the inverse of its exceedance probability:

$$T_r = \frac{1}{1 - F(x_{2024})} \quad (3)$$

To evaluate the severity of the low-tropospheric moisture anomalies without assuming a parametric background distribution, the historical specific humidity and total column water vapor values were ranked against the 42-year record using the non-parametric Weibull plotting position formula:

$$P = \frac{m}{N_{total} + 1} \quad (4)$$

where “m” represents the ascending rank of the observed value (with m = 42 for the absolute historical maximum) and $N_{total} = 42$ is the total number of years in the 1983 - 2024 series. The corresponding percentile rank is expressed as $PR = P \times 100$, allowing an objective determination of the rarity of the thermodynamic preconditioning. Following Makkonen (2006), this specific formulation represents the only mathematically consistent plotting rule for extreme value analysis, preventing the overestimation of return periods often induced by alternative empirical plotting positions and providing a realistic assessment of weather-related risks in hyper-arid environments.

2.4. Atmospheric Circulation and Thermodynamic Indicators

2.4.1. Intertropical Front (ITF) Position

The ITF serves as a critical diagnostic for monitoring the northward penetration of the West African Monsoon (WAM) into the Saharan heat low. In this study, the latitudinal position of the ITF is identified using the 15°C isodrosotherm (dew point temperature $T_d = 15^\circ\text{C}$) at the 925 hPa level, following the criteria established by Nicholson (2013). This boundary marks the interface between the moist south-westerly monsoon flow and the dry north-easterly Harmattan winds.

2.4.2. Integrated Vapor Transport (IVT)

To capture the horizontal moisture streams fueling the convective anomalies over northern Chad, the time series of monthly mean interannual vertically Integrated Vapor Transport (IVT) magnitude was computed. Originally conceptualized to distinguish concentrated filamentary moisture fluxes from broader background fields (Zhu and Newell, 1998), the analysis of IVT magnitudes has become a robust standard for diagnosing synoptic-scale moisture transport and associated hydrometeorological extremes (Gimeno et al., 2014). The zonal (IVT_u) and meridional (IVT_v) mass-weighted components ($\text{Kg.m}^{-1}.\text{s}^{-1}$) were obtained by integrating the product of the specific humidity (q) and horizontal wind components (u, v) across the active tropospheric column (Lavers and Ramos, 2019):

$$IVT_u = \frac{1}{g} \int_{P_{top}}^{P_{surf}} u \cdot q \cdot dp \quad ; \quad IVT_v = \frac{1}{g} \int_{P_{top}}^{P_{surf}} v \cdot q \cdot dp \quad (5)$$

where g is the standard gravitational acceleration (9.81m.s^{-2}), q (Kg.Kg^{-1}), u and v (m.s^{-1}), and dp is the pressure increment (Pa). Vertical integration bounds were restricted from $p_{surf} = 1000$ hPa to $p_{top} = 200$ hPa. This configuration captures the full structural depth of the monsoon circulation,

including the coupled lower-tropospheric moisture surges preconditioned by the Saharan heat low, the horizontal shear zones along the African Easterly Jet (AEJ), and upper-level transitions near the Tropical Easterly Jet (TEJ) (Gimeno et al., 2016; Awolaye et al., 2025). The total IVT magnitude (IVT_{mag}) was then derived as:

$$IVT_{mag} = \sqrt{IVT_u^2 + IVT_v^2} \quad (6)$$

In the context of the African monsoon belt, evaluating IVT_{mag} is critical as vertically coupled moisture fluxes exceeding specific synoptic thresholds have been directly linked to enhanced organized convection, intense spatially focused precipitation events, and regional aerosol scavenging dynamics (Awolaye et al., 2025).

327

2.4.3. Low-Tropospheric Convergence and Vertical Velocity

The dynamic mechanisms forcing local vertical ascent were diagnosed using two structural fields: low-tropospheric horizontal wind convergence and tropospheric vertical velocity (ω). Wind convergence ($-\nabla \cdot V$) was extracted directly at the 850 hPa level from the native ECMWF spectral model divergence archive (d) to prevent numerical truncation errors arising from post-processing finite-difference schemes. Concurrently, the vertical velocity (ω , expressed in $Pa.s^{-1}$) was examined across multiple pressure levels to assess the deep vertical coupling of the column. Negative values ($\omega < 0$) represent strong updrafts and synoptic-scale convective ascent (Taylor et al., 2017; Tchana et al., 2025).

2.4.4. Thermodynamic Columns and Convective Instability

To evaluate the thermodynamic structure and convective potential governing the anomalous August 2024 event, we analyze two complementary conserved thermodynamic tracers: Moist Static Energy (MSE) and equivalent potential temperature (θ_e). Moist Static Energy (MSE) is used to evaluate the convective potential and stability of the atmosphere (Neelin and Held, 1987; Kenfack et al., 2025). It integrates sensible heat, potential energy, and latent heat. It also quantifies the total energy content of an air parcel by integrating sensible heat, potential energy, and latent heat, and is defined as:

$$MSE = C_p T + gz + L_v q \quad (7)$$

Where:

C_p is the specific heat of dry air at constant pressure ($1004 J Kg^{-1} K^{-1}$); T is the absolute temperature (K), L_v is the latent heat of vaporization, z is the geopotential height (m). By examining MSE anomalies, we can establish if the August 2024 extremes were driven by an unusual amount of moisture (latent heat) or intense localised surface heating (sensible heat).

351 In parallel, the equivalent potential temperature θ_e (K), which dictates the absolute temperature a
352 water-vapor-saturated air parcel would attain if all its moisture condensed pseudo-adiabatically,
353 was calculated following Bolton's (1980) formulation:

354
$$\theta_e = \theta \cdot \exp\left(\frac{L_v q}{C_p T_L}\right) \quad (8)$$

355 Where θ is the dry potential temperature and T_L is the temperature at the lifted condensation
356 level. In the tropical troposphere, MSE and $C_p \theta_e$ are tightly coupled and approximately linearly
357 related ($MSE \approx C_p \theta_e$), as both conserve total energy during moist moist-adiabatic processes
358 (Neelin and Held, 1987).

359 From a diagnostic standpoint, an increase in low-level MSE (and thus θ_e) operates via a critical
360 dual thermodynamic mechanism: it enhances the thermal buoyancy of a lifted air parcel, thereby
361 expanding the convective available potential energy aloft, while simultaneously eroding the
362 convective inhibition barrier at the base of the atmospheric profile, facilitating spontaneous and
363 deep convective triggering.

364

365 3. Results


366

367 3.1. Multi-Decadal Rainfall Trends, IVT Evolution, and the Hydro-Climatic Transition

368 The interannual variation in August rainfall over the northern Chadian Sahara (16°–24°N,
369 13°–25°E) characterizes a hyper-arid hydro-climatic regime defined by extreme scarcity and
370 inherent high interannual variability. Historically, monthly totals over this region hover around a
371 very low climatological baseline, with a long-term mean of 12.7 mm for TAMSAT and 9.6 mm
372 for CHIRPS. This hyper-arid dynamic is intrinsically governed by the Saharo-Sahelian transition
373 zone, where precipitation depends strictly on the erratic northward excursions of the Intertropical
374 Front (ITF) and the seasonal migration of the West African Monsoon (WAM) (Nicholson et al.,
375 2013; 2018). An analysis of the 42-year time series (1983 - 2024), comparing the TAMSAT (Fig.
376 2a) and CHIRPS (Fig. 2b) satellite products reveal a remarkable structural consistency in
377 capturing regional climate variability. Both independent datasets accurately identify historical
378 dry cycles such as the severe drought years of 1984, 1990, and 2000, as well as past wet episodes
379 like 1988, 1999, and the 2018-2020 corridor. Although a systematic magnitude offset exists
380 between the two products due to their distinct retrieval algorithms, their long-term climate
381 trajectories are perfectly synchronized.

382

383



384

385 Figure 2: August precipitation trends and patterns (solid blue line) averaged over Northern Chad
 386 (16° - 24°N, 13° - 25°E) from 1983 to 2024. Panels display (a) TAMSAT and (b) CHIRPS
 387 satellite series. The solid red line represents the monotonic linear Sen's slope; the solid black line
 388 marks the long-term climatological mean.


389

390 Both datasets exhibit a robust, continuous, and statistically significant monotonic upward trend
 391 over the entire 1983 - 2024 period, with exceptionally high confidence levels ($p = 1.35 \times 10^{-5}$ for
 392 TAMSAT and $p = 1.06 \times 10^{-4}$ for CHIRPS). Rather than an abrupt structural regime shift or a
 393 statistical break in the trend line itself, the year 2003 marks a climatological transition period
 394 corresponding to the chronological midpoint. This year represents the exact chronological
 395 midpoint where the long-term linear Sen's slope intersects the historical baseline mean. Before
 396 2003, the regional signal was heavily weighted by the extreme aridity and persistent rainfall
 397 deficits of the late 20th-century drought, with the vast majority of years falling well below the

398 long-term average. After 2003, due to a sustained humidification trend, a profound transition
 399 occurred. The frequency of extreme dry years drops sharply, replaced by a dense clustering of
 400 positive anomalies and closely spaced rainfall peaks (notably post-2017), establishing a moister
 401 "new normal" background state over the Chadian Sahara.

402 This multi-decadal evolution is quantified by a positive Sen's slope of +0.44 mm/year for
 403 TAMSAT and +0.17 mm/year for CHIRPS, corresponding to substantial relative magnitudes of
 404 +145.1% and +74.4% over the 42-year record, respectively. This long-term wetting trend over
 405 Northern Chad aligns with the wider "greening" and hydrometeorological intensification signals
 406 documented across the central Sahel (Brandt et al., 2015).

407 To establish a clear physical connection between this long-term precipitation trend and
 408 large-scale atmospheric drivers, a multi-decadal time series of the Integrated Vapor Transport
 409 (IVT) during August was analyzed over the same domain (Fig. 3). As an integrated indicator of
 410 lower-tropospheric moisture supply and wind convergence, the IVT exhibits an exceptional,
 411 statistically significant co-variability with the satellite rainfall series.



412
 413 Figure 3: Same as in Fig. 2, but for Integrated Vapor Transport (IVT, $\text{kg m}^{-1} \text{s}^{-1}$). The dashed red
 414 line represents the Sen's slope; the solid black line marks the historical mean ($91.2 \text{ kg m}^{-1} \text{s}^{-1}$).
 415

416 The IVT time series confirms a robust atmospheric moisture-loading trend, featuring a highly
 417 significant Sen's slope of +1.28 units/year ($p = 8.50 \times 10^{-8}$) and a relative magnitude increase of
 418 +59.2%. More specifically, the IVT reproduces exactly the same physical behavior as the
 419 precipitation data:

- 420 1. Late 20th-century suppressed state: Persistent negative IVT anomalies dominate the
 421 1980s and 1990s, indicating a severely restricted moisture supply to the region.
- 422 2. The early-2000s hydro-climatic transition: The IVT line permanently crosses its historical
 423 climatological mean ($91.2 \text{ kg m}^{-1} \text{s}^{-1}$) around 2003, transitioning the regional atmosphere
 424 into a consistently heightened moisture-loading baseline.

3. The 2024 Extreme Manifestation: The record-breaking event of August 2024 sits at the absolute tail end of this multi-decadal trend, where the IVT reaches an unprecedented historical peak exceeding $150 \text{ kg m}^{-1}\text{s}^{-1}$.

The record-breaking precipitation anomaly of August 2024, which reached an unprecedented regional average of 69.71 mm/month in TAMSAT and 22.35 mm/month in CHIRPS, represents the absolute extreme manifestation of this long-term hydro-climatic transition. Rather than occurring as an isolated, random synoptic anomaly, the exceptional magnitude of the 2024 event was fundamentally favored by this altered, post-2003 moister background state. The long-term intensification of the atmospheric moisture transport background state effectively maximized the regional susceptibility to extreme monsoonal rainfall. This setup allowed contemporary dynamic and thermodynamic forces to trigger an unprecedented northward penetration of the WAM, shifting the regional risk paradigm from perpetual hyper-aridity to acute flash flood hazards.

This multi-decadal shift toward a structurally moister baseline suggests that the Chadian Sahara has transitioned into a hydroclimatic regime that is fundamentally more susceptible to intense convective forcing. Within this altered background state, the exceptional event of August 2024 does not emerge as an isolated, uncoupled anomaly but rather as the absolute tail-end expression of a long-term intensification in regional moisture loading. To fully understand how this large-scale climatological transition manifests on a local scale, it is necessary to move beyond domain-averaged time series and dissect the explicit spatial architecture, structural magnitude, and statistical return periods of the August 2024 anomalies across the hyper-arid core.

3.2. Magnitude and Statistical Significance of the August 2024 Rainfall Anomalies

In response to the long-term hydroclimatic transition and the regional-scale intensification of Integrated Vapor Transport (IVT), the August 2024 event manifested as a highly intense and spatially heterogeneous deployment, which had been masked until now by domain-averaged metrics. Examining fine-scale spatial structures thus becomes essential to map how this historical anomaly is distributed across the core hyper-arid northern sector of Chad. To evaluate this spatial architecture and the exceptional severity of the August 2024 rainfall within this hyper-arid zone, we examine the distribution of monthly accumulated precipitation alongside probabilistic Z-scores. Figure 4 illustrates the monthly accumulated rainfall and corresponding standardized anomalies over northern Chad, compared against the 41-year historical baseline climatology (1983–2023).

Although the preceding regional temporal analysis (Section 3.1) demonstrated a remarkable synchronization between the TAMSAT and CHIRPS datasets, the spatial characterization presented here (Figure 4) relies exclusively on the TAMSAT product. This methodological choice is motivated by TAMSAT's algorithmic design optimized for tracking African convective systems in arid zones. Furthermore, since the analysis critically depends on the calculation of standardized anomalies (Z-scores), the systematic magnitude offset observed between the two satellite products is mathematically normalized. This ensures that the spatial structure and

statistical significance of the 2024 rainfall plume remain robust and independent of the selected product.

The baseline climatology (Fig. 4a) highlights the extreme and uniform aridity that typically characterizes the region. Within the central Saharan sectors encompassing the massifs and plateaux of the Tibesti, Borkou, and Ennedi, mean August accumulations are nearly non-existent and remain strictly below 5 mm.




Figure 4: Spatial distribution, magnitude, and statistical significance of August precipitation over Northern Chad derived from the TAMSAT product. Panels display (a) the multi-decadal baseline

climatology (1983-2023), (b) absolute monthly rainfall accumulations for August 2024, and (c) the corresponding standardized anomalies (Z-score). Stippling in panel (c) denotes statistically significant anomalies ($p \leq 0.05$) derived from a grid-point-wise Monte Carlo simulation.

In contrast to this baseline state, the spatial structure of the August 2024 rainfall (Fig. 4b) exhibits a remarkable northward thrust of precipitation well beyond the 20°N parallel. This rainfall activity was organized around two primary, well-defined core nodes:

- The first developed over the rugged topography of the Ennedi and extended westward into the Borkou region, generating unprecedented absolute accumulations ranging from 150 to 400 mm.
- The second manifested further to the northwest, directly impinging on the volcanic massifs of the Tibesti region, with totals ranging from 100 to 250 mm.

The standardized anomaly field (Fig. 4c) provides a mathematical context for the extraordinary magnitude of this event. Localized departures from the historical mean are exceptional, with Z-scores systematically exceeding $+8\sigma$, and localized peaks transcending $+12\sigma$ within the Borkou and Ennedi sectors. Such values underscore the profound statistical deviation of this event from regional historical observations.

Finally, to ensure that this signal is reliable and does not result from random variability, a grid-point-wise statistical Monte Carlo significance test was applied. The resulting empirically derived significance mask (represented by the stippling in Fig. 4c) completely envelopes the Tibesti, Borkou, and Ennedi regions. The spatial continuity of this mask confirms that the northward advance of the monsoon and the associated hyper-arid precipitation constitute a statistically significant anomaly ($p \leq 0.05$) that is unprecedented since the beginning of the satellite era, indicative of an exceptional displacement of the West African Monsoon (WAM) boundaries during this period.

Table 1: Parametric Return Periods and Non-Parametric Thermodynamic Percentile Ranks for the August 2024 Extreme Event based on spatial averages over the Northern Chad study domain (16°N–24°N, 13°E –25°E). Rainfall metrics are based on a Gamma distribution fitted via Maximum Likelihood Estimation (MLE) over the 1983–2024 baseline. Specific humidity (q) and Total Column Water Vapor (TCWV) metrics represent empirical ranks and percentile ranks calculated using the Weibull formula over the same climatological period (N = 42 years).

Diagnostic Category	Parameter / Dataset	Observed Value (August 2024)	Empirical Rank (Out of 42)	Calculated Percentile Rank (%)	Estimated Return Period (T, years)
Pluviometric Metrics	CHIRPS Rainfall	22.35 mm	–	–	124.5

	TAMSAT Rainfall	69.71 mm	–	–	451.4
Thermodyna mics Metrics	ERA5 Specific Humidity q at 850 hPa	10.67g.kg ⁻¹	42	97.67%	–
	ERA5 Specific Humidity q at 925 hPa	11.90g.kg ⁻¹	42	97.67%	–
	ERA5 Column Water Vapor TCWV (Total Column)	14.83kg.m ⁻²	42	97.67%	–

While the spatial architecture of Figure 4 illustrates the geographic extent of this event along the Tibesti-Borkou-Ennedi corridor, it raises, above all, the question of its statistical return period and the underlying atmospheric conditions that allowed such anomalies to persist. To place the extreme nature of this plume within a historical context spanning several decades, we employ a combined diagnostic framework that links parametric return periods of precipitation to nonparametric empirical ranks of tropospheric moisture content. To quantify the historical rarity of this event, a rigorous dual-probabilistic diagnostics framework was applied to the core convective zone (16°–24°N, 13°–25°E).

The results, summarized in Table 1, contrast the parametric return periods (T) derived from the optimized two-parameter Gamma distribution against the empirical thermodynamic percentile ranks (PR) computed via the mathematically consistent Weibull plotting position (Makkonen, 2006). The regionalized August 2024 rainfall anomaly exhibits a return period of 124.5 years according to CHIRPS (mean: 22.35mm), and reaches up to 451.4 years according to the TAMSAT satellite product (mean: 69.71mm). The structural discrepancy between these two satellite estimates reflects well-known algorithmic sensitivities in hyper-arid, gauge-sparse regions. TAMSAT relies exclusively on Thermal Infrared (TIR) Cold Cloud Duration (CCD) thresholds calibrated specifically for African convective regimes (Maidment et al., 2014, 2017). Consequently, it effectively captured the persistent, deep cloud tops and prolonged convective activity that sustained this specific event, likely yielding a physically representative depiction of the local convective intensity. Conversely, while CHIRPS integrates TIR data, its methodology heavily relies on the blending of in-situ rain gauge observations for structural bias correction (Funk et al., 2015). In the hyper-arid northern territory of Chad, the severe scarcity of operational, real-time reporting gauges forces the CHIRPS algorithm to heavily weight its

background climatology, typically inducing a conservative underestimation during unprecedented, high-magnitude anomalies. Despite this systematic magnitude offset, both independent datasets structurally agree on the centennial to multi-centennial scale of the anomaly. This out-of-boundary pluviometric response was systematically fueled by an extraordinary, vertically coherent thermodynamic state throughout the tropospheric column. The monthly mean specific humidity (q) reached absolute historical records at both the lower-tropospheric base (11.90 g kg^{-1} at 925 hPa) and the core of the monsoonal layer (10.67 g kg^{-1} at 850 hPa).

This profound moistening is further corroborated by the Total Column Water Vapor (TCWV), which attained an unprecedented historical peak of 14.83 kg m^{-2} . For all three hydro-thermodynamic parameters, August 2024 ranked 42nd out of the 42-year climatological record, placing the month at the 97.67th empirical percentile of the historical distribution (Table 1). This joint analysis confirms that the historic rainfall observed in the Saharan regions of Chad resulted from a perfect synergy between, on the one hand, a local moisture reservoir reaching its absolute maximum, thereby preventing the inflow of dry air and lowering the condensation level due to lifting, and, on the other hand, a dynamically optimized triggering mechanism at the synoptic scale. However, while these extraordinary thermodynamic conditions created a highly conducive and moisture-saturated background state, they constitute a necessary but not sufficient condition for such extreme convective development. To fully understand how this localized precipitable water was dynamically lifted and organized into plumes of intense precipitation across the hyper-arid heartland, it is essential to shift our focus from hydrostatic column measurements to the synoptic-scale transient triggers that governed regional atmospheric circulation during August 2024.

3.3. Synoptic-Scale Triggering: African Easterly Waves (AEWs) Dynamics

To study the synoptic-scale atmospheric mechanisms underlying the exceptional precipitation anomalies observed in August 2024, Figure 5 presents a longitude-time (Hovmöller) diagram of the meridional wind and relative vorticity at 700 hPa , averaged over the Saharan belt (16°N – 24°N). During the first half of the month (from August 1 to 16), the regional atmosphere was featured by robust and well-defined activity of the AEWs. This dynamic is visually identifiable by distinct diagonal corridors propagating westward, composed of alternating meridional components of north (blue) and south (red) winds. It is important to note that between August 10 and 14, a high-amplitude wave trough penetrated the specific study area bounded by the meridians 13°E and 25°E (vertical dotted lines). This synoptic event was characterized by a powerful acceleration of the southerly wind exceeding $+4 \text{ m s}^{-1}$, which acted as a dynamic conveyor belt, ensuring significant moisture advection from the monsoon reservoirs in the low latitudes toward the hyper-arid interior. This hydrodynamic driving mechanism was closely linked to a significant core of cyclonic relative vorticity (green contour $\geq 1 \times 10^{-5} \text{ s}^{-1}$), ensuring the large-scale upwelling and vertical organization of the necessary cells. This intense dynamic forcing on a synoptic scale explains the onset, development, and

concentration of the extreme hydrometeorological events observed over the Tibesti, Borkou, and Ennedi mountain ranges during this specific period spanning several days. In contrast, the second half of August 2024 (after August 16) was marked by a notable change in weather conditions. The organized, in-phase, and propagating AEW structures have weakened considerably, as evidenced by the disruption of the diagonal wind bands. This state of suppression gave way to more stationary atmospheric features and localized, isolated vorticity maxima, centered mainly near the eastern boundary (25°E) during the last few days of the month. This evolution indicates that, while the first half of the month was dominated by transient wave-induced convective bursts, the second half shifted toward a more localized convective regime of thermodynamic origin, within the context of stationary synoptic patterns.




Figure 5: Longitude-time (Hovmöller) diagram of 700 hPa meridional wind (shaded, m.s^{-1}) and relative vorticity (contours, 10^{-5} s^{-1}) averaged over the Saharan band (16°N – 24°N) during August 2024. Dashed vertical lines delimit the core study domain (13°E – 25°E).

3.4. Climatological mean and anomalous state of lower-tropospheric (850 hPa) circulation, wind convergence, and Intertropical Front (ITF) positions.

The progressive AEW packets analyzed in the previous section are not merely isolated daily weather events; on the contrary, their recurring occurrence and intensity are likely to modulate the large-scale monthly atmospheric state. To quantify the net, integrated impact of

these synoptic drivers on the lower troposphere, this section examines the monthly mean dynamic configuration at 850 hPa. More specifically, Figure 6 displays the spatial distribution of 850 hPa wind convergence, horizontal wind vectors, and the latitudinal position of the Intertropical Front (ITF), defined by the 15°C dew point isodrosotherm. Three states are compared: the 1983–2023 August climatology (Fig. 6a), the August 2024 observed state (Fig. 6b), and the resultant absolute anomalies (Fig. 6c).


In the climatological mean (Fig. 6a), the solid black contour marking the ITF exhibits a zonal orientation localized between 15°N and 18°N, while the color shading reveals a well-defined band of low-level convergence (blue shading; -1 to $-3 \times 10^{-5} \text{ s}^{-1}$) strictly bounded to the Sahelian belt. South of this boundary, the horizontal wind field is characterized by persistent, large-scale southwesterly vectors exceeding 5 m s^{-1} . Conversely, inside the red box defining the northern Chadian study domain (16°N–24°N, 13°E–25°E), the background shading is dominated by positive divergence values (light yellow to white; $+0.5$ to $+2 \times 10^{-5} \text{ s}^{-1}$), followed by strong northeasterly wind arrows. Dynamically, this indicates that significant low-level convergence is normally restricted to the humid monsoonal flow over the Sahel. Northern Chad, including the Tibesti, Borkou, and Ennedi massifs, is climatologically dominated by a strongly divergent, subsiding regime associated with the descending branch of the Hadley cell and dry Harmattan winds. These synoptic conditions maintain absolute atmospheric stability and structurally inhibit deep convection, a configuration perfectly aligned with the established canonical structure of the West African Monsoon (WAM) (Sultan & Janicot, 2003; Nicholson, 2013).

In August 2024 (Fig. 6b), the thick black ITF line displays a pronounced northward deformation, looping up to the 20°N–22°N latitudinal band, while the core of negative divergence values (blue patches; -2 to $-4 \times 10^{-5} \text{ s}^{-1}$) undergoes a substantial northward migration into the Saharan interior. Simultaneously, the horizontal wind vectors show a total reorientation within the study domain, where the traditional northeasterlies are replaced by an uncharacteristic influx of southwesterly and southerly arrows penetrating past 20°N. This interior flow was embedded within a broader synoptic configuration, characterized by enhanced cyclonic circulation gyres visible both to the west and east of the study domain. Visually and physically, this confirms a radical breakdown of the typical desert circulation. This deep latitudinal displacement allowed intense southwesterly monsoonal surges to bypass their usual boundaries and directly establish a powerful dynamic convergence engine over the complex terrain of the Tibesti massif, the Borkou depressions, and the Ennedi plateau. This localized convergence provided the forced mechanical ascent necessary to overcome the hyper-arid background stability, shifting a normally stable region into a highly active convective state. This structural configuration closely mirrors the diagnostic frameworks of extreme multi-day rainfall events documented in arid northern Africa, where topographically locked moisture flux convergence drives unprecedented precipitation events (Vondou et al., 2025).

The absolute anomaly field (Fig. 6c) reveals a vast, cohesive cell of enhanced convergence anomalies (blue shading; reaching -0.6 to $-1.6 \times 10^{-5} \text{ s}^{-1}$) spanning the 15°N–22°N domain, overlaid by a prominent cyclonic (counter-clockwise) wind vector circulation centered

627 over the Chad-Niger border. The solid green contour, tracking the +1 K dew point anomaly
628 threshold, extends well north of 15°N and runs spatially in-phase with the maximum
629 convergence anomalies. Physically, this feature indicates that the dynamic lifting engine was
630 perfectly co-located with a highly anomalous tropospheric moisture loading, which effectively
631 suppressed dry entrainment and favored widespread condensation over the Saharan massifs.
632 Mathematically, this cyclonic wind anomaly explicitly proves that horizontal wind convergence
633 was complemented by a substantial increase in low-tropospheric cyclonic vorticity. Such an
634 environment is vital for providing the mesoscale organization and hydrodynamic longevity
635 required to sustain organized and long-lived convective systems in extreme environments
636 (Lavaysse et al., 2009; Flamant et al., 2018). Crucially, the extensive stippling mask confirms
637 that these enhanced convergence anomalies are statistically significant ($p \leq 0.05$) based on the
638 1000-iteration Monte Carlo permutation test. Because this significant signal is spatially in-phase
639 with the anomalous southwesterly moisture advection, it confirms a total, non-random disruption
640 of the Saharan divergent regime, representing a profound, transient northward expansion of the
641 Sahelian convective climate boundary into the hyper-arid core of Northern Chad (Taylor et al.,
642 2017; Biasutti, 2019). In summary, the out-of-boundary rainfall event of August 2024 resulted
643 from a perfect synergy between the exceptional northward migration of the ITF, reinforced
644 low-level convergence at 850 hPa, and the establishment of a regional cyclonic circulation that
645 optimized the vertical lifting of warm, moist air masses over Northern Chad. These findings
646 strongly align with recent literature linking increased WAM variability to a higher frequency of
647 hydroclimatic extremes along the Sahara-Sahel transition zone (Janicot et al., 2008; Taylor et al.,
648 2017; Biasutti, 2019; Tamoffo et al., 2025,2026c).

649



650

651 Figure 6: Spatial distribution of lower tropospheric dynamics at 850 hPa in August, showing
 652 wind convergence (shading; $\times 10^{-5} \text{ s}^{-1}$), horizontal wind vectors (arrows; m s^{-1}), and tropospheric
 653 moisture boundaries. The panels show: (a) the climatological mean from 1983 to 2023, (b)
 654 conditions observed in August 2024, and (c) absolute anomalies for August 2024 relative to the
 655 climatological reference. In panels (a) and (b), the solid black contour line defines the absolute
 656 position of the intertropical front (ITF) via the dew point isodrosotherm at 15°C (288.15 K). In
 657 panel (c), the solid green line follows the dew point anomaly threshold of $+1 \text{ K}$ (ΔT_d) associated
 658 with the monsoon intrusion, and the stippling indicate regions where convergence anomalies are
 659 statistically significant at a 95% confidence level ($p \leq 0.05$), based on a Monte Carlo permutation
 660 test with 1,000 iterations. The reference vector located at the bottom right of each figure
 661 corresponds to a wind speed of 5 m s^{-1} . The red frame in panel (a) delineates the specific study
 662 area in northern Chad (16°N – 24°N , 13°E – 25°E).

663

664 3.5 Spatial Distribution of Low-Level Humidity and Moisture Flux

While the anomalous 850 hPa wind convergence and northward ITF displacement analyzed in Section 3.4 establish the necessary structural lifting engine over Northern Chad, dynamic forcing alone cannot trigger extreme precipitation without a sufficient and continuous supply of atmospheric water vapor. To assess the thermodynamic environment supporting this localized convergence, it is essential to examine the availability and transport of low-level moisture. Consequently, this section evaluates the spatial distribution of specific humidity and horizontal moisture flux fields at 850 hPa. Figure 7 displays the spatial distribution of 850 hPa specific humidity along with horizontal moisture flux vectors over West and Central Africa. Three configurations are compared: the 1983-2023 August historical climatology (Fig. 7a), the observed conditions in August 2024 (Fig. 7b), and the resulting absolute anomalies (Fig. 7c). The climatological mean (Fig. 7a) highlights a sharp meridional moisture gradient across the region. High specific humidity values (purple shading; exceeding 10 to 14 g kg⁻¹) remain strictly confined south of the 15°N–18°N latitudinal belt, marking the canonical northernmost boundary of the West African Monsoon (WAM) moisture pool. South of this transition zone, the horizontal moisture flux vectors exhibit a powerful, organized southwesterly routing, driving a continuous supply of monsoonal water vapor from the Atlantic toward the Sahelian corridor. Conversely, inside the red box defining the Northern Chadian study domain (16°N–24°N, 13°E–25°E), the lower troposphere is climatologically arid, characterized by low specific humidity values (light blue to white shading; dropping below 6 to 8 g kg⁻¹) and a weak, predominantly zonal moisture transport. This asymmetric structural configuration shows that, under typical summer conditions, the Saharan desert interior is structurally isolated from significant low-level tropical moisture influxes (Nicholson, 2013; Biasutti, 2019). In August 2024 (Fig. 7b), this mean baseline was heavily disrupted. The low-level moisture pool underwent an exceptional northward migration, pushing the 10 to 12 g kg⁻¹ specific humidity thresholds deep into the hyper-arid core of Northern Chad, well past 20°N. Concurrently, the horizontal moisture flux vectors reveal a massive, uncharacteristic reorientation. The climatological continental desert regime was entirely replaced by an intensified, cross-equatorial southwesterly moisture conveyor that bypassed its traditional Sahelian limits to directly feed the complex topography of the Tibesti, Borkou, and Ennedi massifs. Physically, this structural change confirms that the 850 hPa layer was subjected to a continuous, large-scale tropical moisture loading, satisfying a critical thermodynamic prerequisite for widespread and persistent high-latitude rainfall anomalies. The absolute anomaly field (Fig. 7c) provides a quantitative diagnostic of this hydroclimatic shift. A vast, spatially cohesive core of positive specific humidity anomalies (pink shading; reaching +2 to +4 g kg⁻¹) covers the entire 15°N–23°N Saharan interior. Crucially, the extensive stippling mask generated by the 1000-iteration Monte Carlo permutation test confirms that these moist anomalies are statistically significant ($p \leq 0.05$) over virtually the entire northern Chadian domain. This significant moisture surplus runs perfectly in-phase with robust anomalous southerly and southwesterly moisture flux arrows. This alignment explicitly proves that the local humidification of the desert boundary layer was structurally driven by enhanced active advection

from the tropical Atlantic and the monsoonal belt rather than localized evaporation. In summary, the out-of-boundary convective activation of August 2024 resulted from an optimal dynamical-thermodynamical coupling. The exceptional low-level moisture advection documented here (Fig. 7) acted in perfect synergy with the strong 850 hPa wind convergence anomalies previously analyzed. By shifting the normally dry and stable Saharan boundary layer into an uncharacteristically humid and moist state, this combined synoptic-scale mechanism effectively sustained a highly active and long-lived convective regional climate.




Figure 7: Spatial distribution of 850 hPa specific humidity and horizontal moisture flux fields in August over West and Central Africa. The red box outlines the specific study domain over Northern Chad (16°N–24°N, 13°E–25°E). Panels display: (a) the 1983–2023 climatological mean, (b) the observed conditions in August 2024, and (c) the absolute anomalies for August 2024 relative to the historical baseline. Shading denotes specific humidity (g kg^{-1}), and vectors represent horizontal moisture flux ($\text{g kg}^{-1} \text{ m s}^{-1}$). In panel (c), stippling highlights regions where specific humidity anomalies are statistically significant at the 95% confidence level ($p \leq 0.05$).


based on a 1000-iteration Monte Carlo permutation test. Reference vectors corresponding to 150 $\text{g kg}^{-1} \text{ m s}^{-1}$ (panels a, b) and 50 $\text{g kg}^{-1} \text{ m s}^{-1}$ (panel c) are displayed in the bottom-right corners.

3.6. Atmospheric Vertical Structure and Meridional Circulation Cross-Sections

The horizontal accumulation of low-level moisture and enhanced moisture fluxes documented in Section 3.5 over the Saharan interior cannot trigger extreme rainfall anomalies without a corresponding vertical mechanism capable of lifting this moist air mass throughout the troposphere. To understand the dynamic framework supporting these surface thermodynamic shifts, it is essential to examine the meridional-vertical structure of the atmospheric column. Consequently, this section investigates the vertical overturning regional circulation. Figure 8 presents the latitude-pressure cross-sections of vertical velocity (ω , shaded) and superimposed meridional-vertical circulation vectors ($\mathbf{v}$, ω) averaged over the 13°E–25°E longitudinal band for the August historical climatology (Fig. 8a), August 2024 (Fig. 8b), and their absolute anomalies (Fig. 8c). Following the standard meteorological convention, negative ω values (blue shading) denote upward atmospheric motion, whereas positive values (red shading) correspond to subsidence.

In the historical climatology (Fig. 8a), the regional overturning circulation exhibits a classic asymmetric Hadley-type cell configuration. Deep and intense ascending motions ($\omega < 0$; blue shading) are structurally confined between 10°N and 15°N, extending from the lower boundary layer up to the upper troposphere (~300 hPa). This deep vertical convective core represents the mean position of the summer monsoonal ascending branch. North of 18°N, including the northern Chadian study domain (16°N–24°N, highlighted by the vertical dashed lines), the troposphere is heavily dominated by widespread subsidence ($\omega > 0$; red shading), particularly in the middle and upper layers (600–200 hPa). This pervasive mid-tropospheric capping downwelling structurally stabilizes the Saharan atmosphere, explaining the hyper-arid baseline of the desert interior under typical summer conditions.

During August 2024 (Fig. 8b), this mean vertical circulation experienced a profound latitudinal reorganization. The core of deep monsoonal ascent underwent an exceptional northward migration and intensification, with robust negative ω fields (blue shading) spreading continuously into the Saharan interior up to 21°N–22°N. The circulation vectors reveal a powerful, uninterrupted vertical conveyor bridging the lower layers (1000–850 hPa) directly with the upper tropospheric outflow up to 200 hPa. This structural shift implies that the climatological Saharan subsidence capping was completely broken down and replaced by a deep, regional-scale ascending chimney, creating an environment highly favorable for sustained regional moist convection.



755

756 Figure 8: Latitude-pressure cross-sections of meridional-vertical atmospheric circulation and
 757 thermodynamic structure for August averaged over the 13°E - 25°E longitudinal band. Shading
 758 displays the vertical velocity (ω , $\times 10^{-2}$ Pa s $^{-1}$), where negative values denote upward motion and
 759 positive values correspond to subsidence. Vectors represent the combined meridional and vertical
 760 wind components (v and ω , with vertical velocity scaled for visualization). Panels show (a) the
 761 1983-2023 climatological mean, (b) August 2024, and (c) the absolute anomalies (August 2024
 762 minus climatology). In panel (c), stippling isolates regions where anomalies are statistically
 763 significant at the 95% confidence level ($p \leq 0.05$) based on a 1000-iteration Monte Carlo
 764 permutation test. Dashed vertical lines delimit the latitudinal domain of the study area (16°N -
 765 24°N).

766 The absolute anomaly field (Fig. 8c) confirms the statistical robustness of this dynamic
 767 displacement, with the 1000-iteration Monte Carlo permutation test isolating large areas of
 768 significant changes ($p \leq 0.05$; stippling mask) within the Saharan interior. A prominent, deeply

769 rooted negative ω anomaly (blue shading) occupies the 16°N–24°N latitudinal belt, indicating an
770 unprecedented intensification of upward vertical velocity relative to the climatology. Directly
771 south of this feature, between 15°N and 18°N and confined to the lower troposphere (1000–700
772 hPa), a localized positive ω anomaly (red shading) is visible, accompanied by
773 downward-pointing anomalous circulation vectors. When cross-examined with the absolute
774 fields (Figs. 8a, 8b), where blue shading (upward motion) persists during both periods, this
775 positive anomaly does not represent physical synoptic-scale subsidence or an invasion of dry
776 desert air. Instead, it indicates a localized, non-statistically significant reduction in the magnitude
777 of the monsoonal updrafts. Because the main convective core of the West African Monsoon
778 underwent a pronounced northward migration beyond 18°N in August 2024, the vertical ascent
779 within the traditional 15°N–18°N latitudinal belt was weaker in absolute terms than its strong
780 historical climatological counterpart, mechanically generating a positive anomaly value.

781 In summary, Figure 8 demonstrates that the extreme Saharan rainfall of August 2024 was driven
782 by an optimal dynamical-thermodynamical synergy. The exceptional low-level moisture
783 advection previously analyzed acted in perfect harmony with a major vertical reorganization of
784 the West African Monsoon circulation, providing the large-scale lifting engine necessary to
785 sustain deep and widespread atmospheric destabilization over Northern Chad.

786


787 3.7 Spatial patterns of moist static energy anomalies

788 While the anomalous meridional-vertical overturning analyzed in Section 3.6 provides the deep
789 dynamical lifting engine necessary to breach the Saharan subsidence capping, the longevity and
790 intensity of such regional updrafts depend heavily on the continuous availability of thermal and
791 latent energy at the surface. To map the horizontal footprint and intensity of the energy reservoir
792 feeding this enhanced vertical circulation, it is essential to examine the spatial distribution of key
793 atmospheric thermodynamic tracers. Consequently, this section investigates the regional patterns
794 of low-level energy accumulation.

795 Figure 9 illustrates the spatial distribution of low-level (at the 850 hPa pressure level) moist
796 static energy (MSE; shaded, kJ kg^{-1}) anomalies and the corresponding contours of equivalent
797 potential temperature (θ_e ; black contours, K) anomalies over the Sahelian and Saharan sectors.
798 As an integrative thermodynamic state variable combining sensible heat, geopotential, and latent
799 enthalpy contributions, low-level MSE serves as a robust diagnostic tool to link sub-synoptic
800 dynamic forcing (such as wind convergence) to the thermodynamic conditions required to fuel
801 deep convection (Neelin and Held, 1987; Romps, 2015). To guarantee the statistical robustness
802 of the featured structures, only highly significant anomalies exceeding twice the climatological
803 standard deviation ($> 2\sigma$) are represented in this figure. Robust positive MSE anomalies
804 dominate the central and eastern Saharan latitudes, displaying a well-defined zonal structure. The
805 core of these statistically significant thermodynamic anomalies is centered between 18°N and
806 25°N, directly encompassing the Northern Chad study area designated by the red bounding box.
807 This spatial configuration reflects an exceptional energetic enrichment of the lower troposphere
808 over an arid region typically characterized by a low historical energy baseline. Conversely, the

809 weaker or absent anomalies observed further south imply a significant northward shift of the
 810 moist monsoon energy reservoir into the Saharan transition zone. In accordance with the
 811 theoretical framework of Romps (2015), the accumulation of high boundary-layer MSE shifts the
 812 ascending air parcel's moist adiabat, directly maximizing its thermal buoyancy integral aloft and
 813 unlocking the potential for deep convective initiation. The contours of θ_e anomalies exhibit a
 814 spatial geometry that perfectly mirrors the MSE fields, with isolines strictly bounding the core of
 815 the energy maxima. The tight spatial co-location between the maximum MSE shading and the
 816 elevated θ_e contours highlights their direct thermodynamic covariation arising from shared
 817 conservation properties during moist adiabatic displacements within the planetary boundary
 818 layer. The northward extension of the 6 K and 10 K anomaly contours deep into the Sahara
 819 Desert confirms that the observed regional warming and moisture enrichment were
 820 co-dependent, fueled by the intense horizontal advection of moist monsoonal air coupled with
 821 anomalous surface diabatic heating. This closely coupled MSE- θ_e configuration marks a
 822 profound reorganization of the regional desert thermodynamic environment relative to its
 823 baseline. Such persistent, large-scale positive energy anomalies exceeding the 2σ threshold are
 824 recognized precursors to extreme rainfall over West Africa when spatially aligned with
 825 lower-tropospheric convergent dynamic structures (Taylor et al., 2017). In a broader context, the
 826 heavy lower-tropospheric energy loading illustrated in Figure 9 is consistent with recent
 827 mechanistic views linking the intensification of the West African monsoon system to a higher
 828 frequency of intense rainfall over North Africa (Biasutti, 2019). This thermodynamic
 829 configuration highlights the role of atmospheric moisture preconditioning and its interaction with
 830 the Sahelian dryline boundary, which are critical precursors for driving high-impact convective
 831 extremes across the region (Vizy and Cook, 2022). The accumulation of high surface MSE over
 832 Northern Chad acted as a powerful thermodynamic primer; once matched with the persistent 850
 833 hPa wind convergence and moisture pump documented in the preceding sections, it sustained the
 834 development and longevity of the severe convective systems observed in August 2024. This
 835 synergy aligns perfectly with the regional-scale mechanisms described by Akinsanola et al.
 836 (2019), whose moisture budget analyses demonstrate that rainfall intensification across the
 837 central and eastern Sahel is intrinsically driven by a co-forcing between dynamic convergence in
 838 the low-to-mid troposphere and the thermodynamic enrichment of surface moisture.

839



840

841 Figure 9. Spatial anomalies of moist static energy (MSE; shaded, kJ kg^{-1}) and equivalent
 842 potential temperature (θ_e ; black contours, K) at the 850 hPa pressure level. The map illustrates
 843 the coupling between regional thermodynamic variability and the spatial organization of MSE
 844 over the Sahel-Saharan region. The superposition of these fields highlights the northward
 845 expansion of high-energy air masses and the intensification of the monsoon-related energy
 846 reservoir.

847

848 3.8. Socio-economic impacts of the August 2024 extreme event

849 In 2024, Chad experienced the most catastrophic flooding in its recorded history. Nationwide, the
 850 impact was profound: nearly 2 million people were affected, resulting in 576 confirmed fatalities.
 851 The floods submerged approximately 432,800 hectares of agricultural land, destroyed 217,700
 852 homes, and led to the loss of 72,100 head of livestock. Furthermore, infrastructure damage was
 853 extensive, with 3,058 schools severely impacted. Geographically, the crisis was universal,
 854 affecting all 23 provinces. Notably, the Saharan provinces, typically spared due to their
 855 hyper-arid climate, suffered disproportionately heavy losses.
 856 In the desert zone, initial warnings were issued in July as flooding began in the south. However,
 857 torrential rains struck the northern desert region with unprecedented intensity between August
 858 9th and 14th, 2024. National meteorological records documented 126.5 mm of rainfall over eight
 859 days in areas where total annual precipitation rarely reaches 100 mm (Tchadinfos). These levels
 860 represent the highest rainfall totals in 60 years. Visual evidence (Fig. 10) corroborates these

861 records, showing water levels reaching knee-depth at the Faya-Largeau market, where
 862 commercial and material losses were severe.

863



864

865 Figure 10: Flooded streets and commercial stalls at the Faya-Largeau market (Source: Tchad Info
 866 Société).

867

868 Within the specific study area, the flooding impacts are categorized by human displacement,
 869 mortality, agricultural devastation, and infrastructural loss. While national reports often
 870 aggregate these figures, detailed provincial data is typically only prioritized where significant
 871 losses of life occur. Consequently, the situations in Borkou (4 deaths) and Tibesti (60 deaths) are
 872 better documented than in the Ennedi provinces, which reported no fatalities. However, OCHA
 873 situational reporting as of October 1, 2024, allows for a more granular assessment of the affected
 874 populations across all study provinces. This disparity highlights the critical need for developing
 875 nations to enhance civil protection services and systematic data collection during
 876 hydrometeorological disasters (Saha et al., 2018; Mazhin et al., 2021). Table 2 presents the
 877 damage breakdown for each province covered by this study.

878

879 Table 2: Losses due to flooding in the study provinces

880

Region	Number of families affected	Number of people affected	Number of dead recorded	Number of houses/shops destroyed	number of animals killed
Borkou	10,402	67,613	04	23,174 houses	861
Ennedi East	35,980	233, 872			
Ennedi West	1, 842	11,977			

Tibesti	23	150	60	Thousands shops	

Beyond the quantified losses in Table 2, the regional economy, heavily reliant on trade and mining, suffered significant shocks. Thousands of shops were destroyed, and merchandise was lost. In the Tibesti region, gold mining operations suffered immense equipment losses. A particularly grave secondary hazard emerged as the 2024 floods unearthed thousands of landmines remaining from previous conflicts, creating a long-term security and humanitarian risk for the local population.

Conclusion

This study elucidates the atmospheric mechanisms and long-term hydro-climatic shifts driving the unprecedented, record-breaking rainfall event observed in August 2024 across the hyper-arid Chadian Sahara (16°N–24°N, 13°E–25°E), while documenting its catastrophic socio-economic impacts. Based on a 42-year climatological reference period (1983–2024), the results demonstrate that, far from being an isolated synoptic fluctuation or one unrelated to other phenomena, the historical anomaly observed in August 2024 represents the ultimate manifestation of a persistent and statistically significant trend toward increased humidity over several decades. This long-term transition, evident since the early 2000s, is characterized by a gradual trend toward increased atmospheric moisture content, with a +59.2% rise in integrated water vapor transport (IVT). This altered baseline, which has grown wetter over recent decades, has maximized regional vulnerability to extreme monsoon precipitation, structurally shifting the regional hazard paradigm from perpetual hyper-aridity to severe hydrological risks.

On a regional scale, the exceptional magnitude of the August 2024 event is evidenced by accumulated monthly rainfall totals that reached up to 250–400 mm over the rugged topographies of the Tibesti, Borkou, and Ennedi massifs, representing the highest rainfall totals in 60 years. These accumulations correspond to extraordinary standardized anomalies systematically exceeding $+8\sigma$ (and locally transcending $+12\sigma$), which are shown to be statistically significant ($p \leq 0.05$) via grid-point Monte Carlo testing. Parametric probability frameworks reveal that this rainfall plume represents a centennial to multi-centennial phenomenon, with return periods estimated between 124.5 years (CHIRPS) and 451.4 years (TAMSAT). This extreme pluviometric response was sustained by an unprecedented, vertically coherent thermodynamic loading, wherein monthly specific humidity at 925 hPa and 850 hPa, alongside Total Column Water Vapor (TCWV), reached their absolute historical records (97.67th empirical percentile), establishing a highly conducive, moisture-saturated column over the Saharan interior. The synthesis of atmospheric diagnostics reveals that this record-breaking event was governed by an optimal, highly synchronized dynamic-thermodynamic coupling. Synoptically, the first half of the month was intensely triggered by high-amplitude African Easterly Waves (AEWs) between August 10 and 14, acting as a powerful dynamic conveyor belt for low-latitude moisture advection. On a monthly scale, this transient wave activity drove a radical breakdown of the typical desert circulation, forcing a profound northward migration of the Intertropical Front

919 (ITF), which reached latitudes of 20°-22°N, supported by a synchronized intensification of the
920 West African Monsoon (WAM).

921 Consequently, the climatological low-level divergence and mid-tropospheric subsidence capping
922 that historically stabilize the desert interior were completely replaced by robust 850 hPa wind
923 convergence anomalies, a sharp increase in lower-tropospheric specific humidity, and an
924 uninterrupted, deep vertical ascending chimney reaching the upper troposphere (~200 hPa). This
925 is evidenced by pronounced negative ω anomalies extending from the lower to the upper
926 troposphere, which perfectly coupled in space with robust lower-tropospheric moist static energy
927 (MSE) and equivalent potential temperature (θ_e , derived via Bolton's formulation) anomalies
928 exceeding the 2σ statistical threshold over Northern Chad. This substantial energetic loading
929 indicates a profound thermodynamic enrichment of the atmospheric column in regions typically
930 dominated by dry air and moisture deficits.

931 The accumulation of high boundary-layer energy acted as a powerful thermodynamic primer;
932 once matched with the persistent low-level convergence pump and horizontal moisture
933 advection, it provided the necessary instability to sustain the vertical lifting and longevity of the
934 convective systems, resulting in the exceptional rainfall totals observed.

935 The socio-economic consequences of this hydroclimatic shift were historically devastating. On a
936 national scale, the floods affected nearly 2 million people, claiming 576 lives, submerging
937 432,800 hectares of agricultural land, and destroying 217,700 homes and 3,058 schools. Within
938 the hyper-arid study domain, the impacts were severe and spatially heterogeneous: the Borkou
939 province recorded over 67,000 affected individuals and the destruction of 23,174 structures,
940 while the Tibesti region suffered a heavy human toll with 60 confirmed fatalities. Beyond
941 structural and livestock losses, the severe rainfall caused major economic shocks to local trade
942 and gold mining operations due to massive equipment destruction. Crucially, the floods
943 generated a grave secondary humanitarian hazard by unearthing thousands of conflict-era
944 landmines, introducing long-term security risks for local populations.

945 Finally, the scale of this disaster underscores the profound vulnerability of hyper-arid regions to
946 sudden hydrological hazards. Given that the observed intensification of the regional hydrological
947 cycle suggests an increasing recurrence of such high-impact convective extremes under global
948 climate change, there is an urgent need for regional authorities to transition from reactive disaster
949 management to proactive climate adaptation strategies. Furthermore, these findings provide a
950 robust empirical benchmark for improving the representation of monsoon-desert interfaces in
951 climate models, highlighting the necessity for future research utilizing high-resolution
952 convection-permitting simulations to track the long-term evolution of these saharan
953 hydroclimatic shifts.

954

955 CODE AVAILABILITY

956 Figures shown in this study are plotted using the NCAR Command Language (NCL;
957 <https://doi.org/10.5065/D6WD3XH5>, NCAR Command Language, 2017) and QGIS. Codes can
958 be obtained from the corresponding author.

959

960 DATA AVAILABILITY STATEMENT

961 The ERA5 data were obtained from the Copernicus Climate Change Service (C3S) Climate Data
962 Store (<https://cds.climate.copernicus.eu>). Climate Hazards Group InfraRed Precipitation with
963 Station data (CHIRPS) is available through the link <https://www.chc.ucsb.edu/data/chirps3>

964 The TAMSAT rainfall estimates and derived products are based on Meteosat thermal infra-red
965 (TIR) imagery provided by EUMETSAT (<https://data.tamsat.org.uk/data-download/rainfall/>)
966

967 AUTHOR CONTRIBUTIONS

968 CWT: conceptualization; data analysis; formal analysis; investigation; methodology; writing
969 original draft; project administration; review and editing. FS: supervision; conceptualization;
970 data analysis; investigation; project administration; writing original draft; review and editing.
971 ATT: methodology; resources; validation; writing; review; project administration and editing.
972

973 COMPETING INTERESTS

974 The authors declared that they have no conflict of interests.
975

976 ACKNOWLEDGEMENTS

977 The authors thank all the observational and reanalysis data providers used in this study and the
978 research of the International Joint Laboratory “Dynamics of Terrestrial Ecosystems in Central
979 Africa: A Context of Global Changes” (IJL DYCOCA/LMI DYCOFAC).

980 The authors acknowledge the use of Gemini AI (Google) for linguistic refinement and stylistic
981 editing of the manuscript
982

983 References

984

985 Adeyeri, O. E.: Hydrology and climate change in Africa: Contemporary challenges, and future
986 resilience pathways. *Water*, 17, 2247, <https://doi.org/10.3390/w17152247>, 2025

987 Adeyeri, O. E., Lawin, A. E., Laux, P., Ishola, K. A., & Ige, S. O.: Analysis of climate extreme
988 indices over the Komadugu-Yobe basin, Lake Chad region: Past and future occurrences.
989 *Weather and Climate Extremes*, 23, 100194, <https://doi.org/10.1016/j.wace.2019.100194>,
990 2019.

991 Akinsanola, A. A., & Zhou, W.: Dynamic and thermodynamic factors controlling increasing
992 summer monsoon rainfall over the West African Sahel. *Climate Dynamics*, 52,
993 4501–4514, <https://doi.org/10.1007/s00382-018-4394-x>, 2018.

994 Armitage, S. J., Bristow, C. S., & Drake, N. A.: West African monsoon dynamics inferred from
995 abrupt fluctuations of Lake Mega-Chad. *Proceedings of the National Academy of*
996 *Sciences*, 112, 8543–8548, <https://doi.org/10.1073/pnas.1417655112>, 2015.

997 Awolaye, P. O., Adebisi, A. A., Abiodun, B. J., Balogun, I. A., Akinbobola, A., Oluleye, A., &
998 Ajayi, V. O.: Moisture and dust in motion: The dual role of integrated vapour transport
999 over West Africa. *Theoretical and Applied Climatology*, 156, 666,
1000 <https://doi.org/10.1007/s00704-025-05913-1>, 2025.

1001

- 1002 Biasutti, M.: Rainfall trends in the African Sahel: Characteristics, processes, and causes. *WIREs*
1003 *Climate Change*, 10, e591, <https://doi.org/10.1002/wcc.591>, 2019a.
- 1004 Brandt, M., Mbow, C., Diouf, A. A., Verger, A., Samimi, C., & Fensholt, R.: Ground- and
1005 satellite-based evidence of the biophysical mechanisms behind the greening Sahel.
1006 *Global Change Biology*, 21, 1610–1620, <https://doi.org/10.1111/gcb.12807>, 2015.
- 1007 Bolton, D.: The computation of equivalent potential temperature. *Monthly Weather Review*, 108,
1008 1046–1053, [https://doi.org/10.1175/1520-0493\(1980\)108](https://doi.org/10.1175/1520-0493(1980)108), 1980.
- 1009 Cook, K. H., & Vizy, E. K.: *Hydrodynamics of regional and seasonal variations in Congo Basin*
1010 *precipitation*. Springer Science and Business Media LLC, 59, 1775-1797,
1011 <https://doi.org/10.21203/rs.3.rs-879134/v1>, 2021.
- 1012 Ehtasham, L., Sherani, S. H., & Nawaz, F.: Acceleration of the hydrological cycle and its impact
1013 on water availability over land: An adverse effect of climate change. *Meteorology*
1014 *Hydrology and Water Management*, 12, <https://doi.org/10.26491/mhwm/188920>, 2024.
- 1015 Flamant, C., Knippertz, P., Fink, A. H., Akpo, A., Brooks, B., Chiu, C. J., Coe, H., Danuor, S.,
1016 Evans, M., Jegede, O., Kalthoff, N., Konaré, A., Lioussé, C., Lohou, F., Mari, C.,
1017 Schlager, H., Schwarzenboeck, A., Adler, B., Amekudzi, L., ... Yoboué, V.: The
1018 dynamics–aerosol–chemistry–cloud interactions in west africa field campaign: Overview
1019 and research highlights. *Bulletin of the American Meteorological Society*, 99, 83–104,
1020 <https://doi.org/10.1175/bams-d-16-0256.1>, 2018.
- 1021 Gaetani, M., Janicot, S., Vrac, M., Famien, A. M., & Sultan, B.: Robust assessment of the time of
1022 emergence of precipitation change in West Africa. *Scientific Reports*, 10, 7670
1023 <https://doi.org/10.1038/s41598-020-63782-2>, 2020.
- 1024 Gimeno, L., Nieto, R., Vázquez, M., & Lavers, D. A.: Atmospheric rivers: A mini-review,
1025 *Frontiers in Earth Science*, 2, 2, <https://doi.org/10.3389/feart.2014.00002>, 2014.
1026
- 1027 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J.,
1028 Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X.,
1029 Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren,
1030 P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer,
1031 A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S.,
1032 Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F.,
1033 Villaume, S., and Thépaut, J.: The ERA5 global reanalysis, *Q. J. Roy. Meteor. Soc.*, 146,
1034 1999–2049, <https://doi.org/10.1002/qj.3803> , 2020.
- 1035 Holloway, C. E., & Neelin, J. D. (2009). Moisture vertical structure, column water vapor, and
1036 tropical deep convection. *Journal of the Atmospheric Sciences*, 66(6), 1665–1683.

- 1037 <https://doi.org/10.1175/2008jas2806.1>
- 1038 Husak, G. J., Michaelsen, J., & Funk, C.: Use of the gamma distribution to represent monthly
1039 rainfall in Africa for drought monitoring applications. *International Journal of*
1040 *Climatology*, 27, 935–944, <https://doi.org/10.1002/joc.1441>, 2006.
- 1041 Janicot, S., Thorncroft, C. D., Ali, A., Asencio, N., Berry, G., Bock, O., Bourles, B., Caniaux, G.,
1042 Chauvin, F., Deme, A., Kergoat, L., Lafore, J.-P., Lavaysse, C., Lebel, T., Marticorena,
1043 B., Mounier, F., Nedelec, P., Redelsperger, J.-L., Ravegnani, F., ... Ulanovsky, A.:
1044 Large-scale overview of the summer monsoon over West Africa during the AMMA field
1045 experiment in 2006. *Annales Geophysicae*, 26, 2569–2595,
1046 <https://doi.org/10.5194/angeo-26-2569-2008>, 2008.
- 1047 Johannsen, F., Ermida, S., Martins, J., Trigo, I. F., Nogueira, M., & Dutra, E.: *Cold bias of ERA5*
1048 *summertime daily maximum land surface temperature over iberian peninsula*, *Remote*
1049 *Sensing*, 11, 2570, <https://doi.org/10.20944/preprints201909.0268.v1>, 2019.
- 1050 Kenfack, K., Djiotang Tchotchou, L. A., Marra, F., Bellomo, K., Tamoffo, A. T., Tchana, B. C.,
1051 Donfack, F. C., Fotso-Nguemo, T. C., Tanessong, R. S., Djomou, Z. Y., & Vondou, D. A.:
1052 Radiative anomalies associated with extreme precipitation of November 2023 in
1053 equatorial Central Africa. *Atmospheric Research*, 321, 108090.
1054 <https://doi.org/10.1016/j.atmosres.2025.108090>, 2025.
- 1055 Kenfack, K., Marra, F., Yepdo Djomou, Z., Angennes Djiotang Tchotchou, L., Tchio Tamoffo,
1056 A., & Appolinaire Vondou, D.: *Dynamic and thermodynamic contribution to the October*
1057 *2019 exceptional rainfall in West Central Africa*, *Weather and Climate Dynamics*, 5,
1058 1457–72, <https://doi.org/10.5194/egusphere-egu25-347>, 2024.
- 1059 Lafore, J. P., Biasutti, M., Knippertz, P., & Taylor, C.: The west african monsoon. In *World*
1060 *Scientific Series on Asia-Pacific Weather and Climate*, World Scientific, 35–49,
1061 https://doi.org/10.1142/9789813200913_0004, 2017.
- 1062 Lavaysse, C., Flamant, C., Janicot, S., Parker, D. J., Lafore, J.-P., Sultan, B., & Pelon, J. (2009).
1063 Seasonal evolution of the West African heat low: A climatological perspective. *Climate*
1064 *Dynamics*, 33(2–3), 313–330. <https://doi.org/10.1007/s00382-009-0553-4>
- 1065 Livezey, R. E., & Chen, W. Y.: Statistical Field Significance and its Determination by Monte
1066 Carlo Techniques. *Monthly Weather Review*, 46–59.
1067 [https://doi.org/10.1175/1520-0493\(1983\)111](https://doi.org/10.1175/1520-0493(1983)111), 1983.
- 1068 Maidment, R. I., Grimes, D., Allan, R. P., Tarnavsky, E., Stringer, M., Hewison, T., Roebeling,
1069 R., & Black, E.: The 30 year TAMSAT African Rainfall Climatology And Time series
1070 (TARCAT) data set. *Journal of Geophysical Research: Atmospheres*, 119, 10-619,
1071 <https://doi.org/10.1002/2014jd021927>, 2014.

1072 Maidment, R. I., Grimes, D., Black, E., Tarnavsky, E., Young, M., Greatrex, H., Allan, R. P.,
 1073 Stein, T., Nkonde, E., Senkunda, S., & Alcántara, E. M. U.: A new, long-term daily
 1074 satellite-based rainfall dataset for operational monitoring in Africa, *Scientific Data*, 4,
 1075 170063, <https://doi.org/10.1038/sdata.2017.63>, 2017.

1076 Makkonen, L.: Plotting positions in extreme value analysis. *Journal of Applied Meteorology and*
 1077 *Climatology*, 45, 334–340, <https://doi.org/10.1175/jam2349.1>, 2006.

1078 Mazhin, S. A., Farrokhi, M., Noroozi, M., Roudini, J., Hosseini, S. A., Motlagh, M. E.,
 1079 Kolivand, P., & Khankeh, H.: Worldwide disaster loss and damage databases: A
 1080 systematic review. *Journal of education and health promotion*, 10, 329,
 1081 https://doi.org/10.4103/jehp.jehp_1525_20, 2021.

1082 (N.d.). Retrieved January 17, 2026, from
 1083 https://www.iom.int/sites/g/files/tmzbd1486/files/1028/228493/avis-td-proc-24-02_wash-activitie
 1084 s.pdf

1085 Neelin, J. D., & Held, I. M.: Modeling tropical convergence based on the moist static energy
 1086 budget. *Monthly Weather Review*, 115, 3–12,
 1087 [https://doi.org/10.1175/1520-0493\(1987\)115](https://doi.org/10.1175/1520-0493(1987)115), 1987.

1088 Nicholson, S. E.: *Dryland climatology*. Cambridge University Press, 2011.

1089 Nicholson, S. E.: The west african sahel: A review of recent studies on the rainfall regime and its
 1090 interannual variability. *ISRN Meteorology*, 2013, 1–32,
 1091 <https://doi.org/10.1155/2013/453521>, 2013.

1092 Nilsson, E., Becker, P., & Uvo, C. B.: Drivers of abrupt and gradual changes in agricultural
 1093 systems in Chad, *Regional Environmental Change*, 20, 75,
 1094 <https://doi.org/10.1007/s10113-020-01668-9>, 2020.

1095 OCHA “United Nations Office for the Coordination of Humanitarian Affairs”: TCHAD :
 1096 Situation des inondations Rapport de situation no03 (01/10/2024), 2024.

1097 Parker, D. J., Burton, R. R., Diongue-Niang, A., Ellis, R. J., Felton, M., Taylor, C. M.,
 1098 Thorncroft, C. D., Bessemoulin, P., & Tompkins, A. M.: The diurnal cycle of the
 1099 West African monsoon circulation. *Quarterly Journal of the Royal Meteorological*
 1100 *Society*, 131, 2839–2860, <https://doi.org/10.1256/qj.04.52>, 2005.

1101 Pausata, F. S. R., Emanuel, K. A., Chiacchio, M., Diro, G. T., Zhang, Q., Sushama, L., Stager, J.
 1102 C., & Donnelly, J. P.: Tropical cyclone activity enhanced by Sahara greening and reduced
 1103 dust emissions during the African Humid Period. *Proceedings of the National Academy*
 1104 *of Sciences*, 114, 6221–6226, <https://doi.org/10.1073/pnas.1619111114>, 2017.

1105 Pausata, F. S. R., Gaetani, M., Messori, G., Berg, A., Maia de Souza, D., Sage, R. F., &

- 1106 deMenocal, P. B.: The greening of the Sahara: Past changes and future implications. *One*
1107 *Earth*, 2, 235–250, <https://doi.org/10.1016/j.oneear.2020.03.002>, 2020.
- 1108 Raymond, D. J., & Jiang, H.: A theory for long-lived mesoscale convective systems. *Journal of*
1109 *the Atmospheric Sciences*, 47, 3067–3077, [https://doi.org/10.1175/1520-0469\(1990\)047](https://doi.org/10.1175/1520-0469(1990)047),
1110 1990.
- 1111 Romps, D. M.: MSE Minus CAPE is the True Conserved Variable for an Adiabatically Lifted
1112 Parcel. *Journal of the Atmospheric Sciences*, 72, 3639–3646,
1113 <https://doi.org/10.1175/jas-d-15-0054.1>, 2015.
- 1114 Saha, F., Tchindjang, M., Dzana, J. G., & Nguemadjita, D.: Natural hazards in the Far-North
1115 Region of Cameroon and the hydrological extremes dynamics of the Chari-Logone
1116 system. *Physio-Géo. Physical geography and environment*, 15, 69–88,
1117 <https://doi.org/10.4000/physio-geo.10719>, 2020.
- 1118 Saha F., Tchio N. D., Tchindjang M., Voundi É. & Mbevo F. P.: Production of so-called
1119 “natural” risks in major urban centers of Cameroon, *Nat. Sci. Soc.*, 26, 418–433, DOI:
1120 <https://doi.org/10.1051/nss/2019003>, 2018.
- 1121 Selami, N., Sèze, G., Gaetani, M., Grandpeix, J.-Y., Flamant, C., Cuesta, J., & Benabadi, N.
1122 (2021). Cloud cover over the Sahara during the summer and associated circulation
1123 features. *Atmosphere*, 12(4), 428. <https://doi.org/10.3390/atmos12040428>
- 1124 Seneviratne, S. I., Zhang, X., Adnan, M., Badi, W., Dereczynski, C., Luca, A. D., ... & Allan, R.:
1125 Weather and climate extreme events in a changing climate, *Working group I contribution to*
1126 *the sixth assessment report of the IPCC*, 1513–1766
1127 <https://doi.org/10.1017/9781009157896.013>, 2021.
- 1128 Sen, P. K.: Estimates of the regression coefficient based on kendall’s tau. *Journal of the*
1129 *American Statistical Association*, 63, 1379–1389,
1130 <https://doi.org/10.1080/01621459.1968.10480934>, 1968.
- 1131 Sindikubwabo, C., Li, R., & Wang, C.: Abrupt change in Sahara precipitation and the associated
1132 circulation patterns. *Atmospheric and Climate Sciences*, 08, 262–273,
1133 <https://doi.org/10.4236/acs.2018.82017>, 2018.
- 1134 Siwe, A. N., Sandjon, A. T., Tchotchou, L. A. D., Tchinda, C. W., Vondou, D. A., & Nzeukou,
1135 A.: Study the impact of intraseasonal oscillations on water vapour transport in central
1136 africa: The case of 25–70 day oscillations. *International Journal of Climatology*, 46,
1137 e70150, <https://doi.org/10.1002/joc.70150>, 2025.
- 1138 Ssenyunzi, R. C., Oruru, B., D’ujanga, F. M., Realini, E., Barindelli, S., Tagliaferro, G., von
1139 Engeln, A., & van de Giesen, N.: Performance of ERA5 data in retrieving Precipitable
1140 Water Vapour over East African tropical region. *Advances in Space Research*, 65,
1141 1877–1893, <https://doi.org/10.1016/j.asr.2020.02.003>, 2020.

- 1142 Sultan, B., Janicot, S., & Diedhiou, A.: The west african monsoon dynamics. Part I:
1143 Documentation of intraseasonal variability. *Journal of Climate*, 16, 3389–3406,
1144 [https://doi.org/10.1175/1520-0442\(2003\)016, 2003](https://doi.org/10.1175/1520-0442(2003)016, 2003).
- 1145 Tamoffo, A. T., Weber, T., & Saleem, F.: Population exposed to compound extreme hot & humid
1146 and wet & windy events in Africa. *Environmental Research Letters*, 20, 014038,
1147 <https://doi.org/10.1088/1748-9326/ad9d5f>, 2024a.
- 1148 Tamoffo, A. T., Weber, T., Cabos, W., Monerie, P.-A., Cook, K. H., Sein, D. V., ... Jacob, D.:
1149 West african monsoon system's responses to global ocean–regional atmosphere coupling.
1150 *Journal of Climate*, 37, 4291–4312, <https://doi.org/10.1175/jcli-d-23-0749.1>, 2024b.
- 1151 Tamoffo, A. T., Mouassom, F. L., Weber, T., Ghomsi, F. E. K., & Nana, H. N.: Pattern
1152 recognition of West African monsoon extreme rainfall events using convolutional neural
1153 networks, *Geophysical Research Letters*, 53, e2026GL124425,
1154 <https://doi.org/10.1029/2026gl124425>, 2026.
- 1155 Tamoffo, A. T., Mouassom, F. L., Weber, T., & Jacob, D.: Constraining high-emission future
1156 West African monsoon with physics-weighted deep learning ensembles. *Communications*
1157 *Earth & Environment*, 7, <https://doi.org/10.1038/s43247-026-03839-8>, 2026c.
- 1158 Tarek, M., Brissette, F. P., & Arsenault, R.: Evaluation of the ERA5 reanalysis as a potential
1159 reference dataset for hydrological modelling over North America. *Hydrology and Earth*
1160 *System Sciences*, 24, 2527–2544, <https://doi.org/10.5194/hess-24-2527-2020>, 2020.
- 1161 Tarnavsky, E., Grimes, D., Maidment, R., Black, E., Allan, R. P., Stringer, M., Chadwick, R., &
1162 Kayitakire, F. Extension of the TAMSAT satellite-based rainfall monitoring over Africa
1163 and from 1983 to present. *Journal of Applied Meteorology and Climatology*, 53,
1164 2805–2822, <https://doi.org/10.1175/jamc-d-14-0016.1>, 2014.
- 1165 Taylor, C. M., Belušić, D., Guichard, F., Parker, D. J., Vischel, T., Bock, O., Harris, P. P., Janicot,
1166 S., Klein, C., & Panthou, G.: Frequency of extreme Sahelian storms tripled since 1982 in
1167 satellite observations. *Nature*, 544, 475–478, <https://doi.org/10.1038/nature22069>, 2017b.
- 1168 Tchana, B. C., Djomou, Z. Y., Kenfack, K., Tamoffo, A. T., Mouassom, F. L., Tchotchou, L. A.
1169 D., Vondou, D. A., & Mbane, C. B.: Analysis of air temperature anomalies during
1170 abnormal rainfall events in Western Central Africa from September to November 2023,
1171 *Theoretical and Applied Climatology*, 156, 288,
1172 <https://doi.org/10.1007/s00704-025-05528-6>, 2025.
- 1173 Tchinda, C. W., Sandjon, A. T., Tchotchou, A. L. D., Vondou, D. A., & Nzeukou, A.: Impact of
1174 the 25-70 day intraseasonal oscillation on extreme rainfall distribution over Central
1175 Africa, *Journal of the European Meteorological Society*, 4, 100035,
1176 <https://doi.org/10.1016/j.jemets.2026.100035>, 2026.
- 1177 Thorncroft, C. D., Bessemoulin, P., & Tompkins, A. M.: The diurnal cycle of the West African
1178 monsoon circulation. *Quarterly Journal of the Royal Meteorological Society*, 131,

1179 2839–2860, <https://doi.org/10.1256/qj.04.52>, 2005.

1180 Unesco.: Map of the world distribution of arid regions: explanatory note, 1977.

1181 Vizzy, E. K., & Cook, K. H. Distribution of extreme rainfall events and their environmental
1182 controls in the West African Sahel and Soudan. *Climate Dynamics*, 59, 997–1026,
1183 <https://doi.org/10.1007/s00382-022-06171-x>, 2022.

1184 Vondou, D. A., Maranan, M., Fink, A., & Knippertz, P.: Meteorological conditions
1185 leading to a catastrophic, rain-induced landslide in Cameroon in October 2019,
1186 *Quarterly Journal of the Royal Meteorological Society*, 152, 775,
1187 <https://doi.org/10.5194/egusphere-egu25-4516>, 2025.

1188 Wainwright, C. M., Finney, D. L., Kilavi, M., Black, E., & Marsham, J. H.: Extreme rainfall in
1189 East Africa, October 2019–January 2020 and context under future climate change.
1190 *Weather*, 76, 26–31, <https://doi.org/10.1002/wea.3824>, 2020.

1191 Wamba Tchinda, C., Tchakoutio Sandjon, A., Djiotang Tchotchou, A. L., Nzeudeu Siwe, A.,
1192 Vondou, D. A., & Nzeukou, A.: The influence of intraseasonal oscillations on rainfall
1193 variability over Central Africa: Case of the 25–70 days variability. *Scientific Reports*, 13,
1194 19842, <https://doi.org/10.1038/s41598-023-46346-y>, 2023.

1195 Zhang, W., Zhou, T., Ye, W., Zhang, T., Zhang, L., Wolski, P., Risbey, J., Wang, Z., Min, S.-K.,
1196 Ramsay, H., Brody, M., Grimm, A., Clark, R., Ren, K., Jiang, J., Chen, X., Fu, S., Li, L.,
1197 Tang, S., & Hu, S.: A year marked by extreme precipitation and floods: Weather and
1198 climate extremes in 2024. *Advances in Atmospheric Sciences*, 42, 1045–1063,
1199 <https://doi.org/10.1007/s00376-025-4540-4>, 2025.

1200 Zhu, Y., & Newell, R. E.: A proposed algorithm for moisture fluxes from atmospheric rivers.
1201 *Monthly Weather Review*, 126, 725–735, [https://doi.org/10.1175/1520-0493\(1998\)126](https://doi.org/10.1175/1520-0493(1998)126),
1202 1998.
1203