

North Atlantic sea level budget revisited

Zhe Song^{1,2}, Anny Cazenave¹, William Llovel², Andrea Storto³ and Marie Bouih⁴

¹Université de Toulouse, LEGOS (CNES/CNRS/IRD/UT3), 31401 Toulouse, CEDEX 9, France

²Université de Bretagne Occidentale, CNRS, Ifremer, IRD, Laboratoire d'Océanographie Physique et Spatiale (LOPS), IUEM, 29280, Plouzané, France

³Institute of Marine Science, National Research Council of Italy, Rome, Italy

⁴Magellium, 31520 Ramonville St Agne, France

Correspondence to: Zhe Song (zhe.song@univ-brest.fr) and Anny Cazenave (anny.cazenave@univ-tlse3.fr, anny.cazenave@gmail.com)

Ocean Science

2nd Revisioned ~~version~~

22 ~~6~~ July 2026

33 Abstract

34 Based on satellite altimetry, GRACE space gravimetry and Argo-based steric data down to
 35 2000m, recent studies have shown that the North Atlantic sea level budget (i.e., altimetry-based
 36 sea level minus sum of components) of the past two decades is not closed, with strong regional
 37 residuals in the North Atlantic. This was ~~s observed but not suggested to result from salinity~~
 38 ~~errors reported since 2015 in some Argo float measurements.~~ In this study, we revisit the
 39 North Atlantic sea level budget, using satellite altimetry, GRACE and GRACE-FO data,
 40 different Argo products and ~~an two~~ ocean reanalyses (CIGAR ~~and ORAS5~~) over the 2004-
 41 2022 time span. The ocean reanalyses ~~is are~~ used to estimate the manometric contribution, an
 42 alternative to using GRACE data, as well as the deep ocean contribution to the sea level budget,
 43 not yet fully sampled by Argo. Analyzing different data sets allows us to ~~assessevaluate~~ their
 44 impact on the previously reported non-closure of the North Atlantic sea level budget. We first
 45 find that using the CIGAR ocean reanalysis-based manometric component significantly
 46 reduces the residuals of the North Atlantic sea level budget compared to GRACE. We also find
 47 that accounting for the deep ocean (below 2000m) thermal expansion (using the CIGAR
 48 reanalysis) allows for ~~reducing~~ 30% ~~reduction of~~ the North Atlantic budget residuals when
 49 using ~~both~~ GRACE ~~and CIGAR~~ for the manometric component, ~~while the mean residual trend~~
 50 ~~is reduced by a factor of 2 when using CIGAR for the manometric sea level.~~ ~~In the latter~~
 51 ~~case~~ ~~With manometric~~, the budget is closed within data uncertainties. ~~The North Atlantic~~
 52 ~~halosteric component based on Argo and CIGAR in the upper 2000m displays a small decrease~~
 53 ~~since the early 2010s. However, this negative trend becomes stronger after 2016. The 2010-~~
 54 ~~2016 halosteric decrease may reflect a real salinity increase in the region, although salinity~~
 55 ~~measurement errors may have impacted the halosteric component after that date.~~

56

57 1 Introduction

58 While many studies have been devoted to assessing the global mean sea level budget over the
 59 satellite altimetry era (1993 to present) (e.g., Barnoud et al., 2021; Bouih et al., 2025; Chen et
 60 al., 2020; Dieng et al., 2017; Horwath et al., 2022; Llovel et al., 2023; Nerem et al., 2018;
 61 WCRP, 2018) , only a few have focused on the regional sea level budget, with mixed results
 62 (e.g., Bouih et al., 2025; Camargo et al., 2023; Frederikse et al., 2016; Hamlington et al., 2020;
 63 Mu et al., 2024; Royston et al., 2020). [At the global scale, the two main contributors to the](#)
 64 [global mean sea level rise are ocean mass \(known as barystatic sea level, Gregory et al., 2019\)](#)
 65 [and thermosteric change. At the global scale, the main terms of the sea level budget are the](#)
 66 [global mean sea level rise, ocean mass \(known as barystatic sea level, Gregory et al., 2019\),](#)
 67 [and thermosteric sea level changes.](#) At regional scale, however, other factors play a non-
 68 negligible role, such as the halosteric (i.e., salinity-related) component, which makes the
 69 regional sea level budget more complex to assess. [In effect, a spurious salinity drift \(giving rise](#)
 70 [to an important halosteric component decrease\) has been reported recently in some Argo-based](#)
 71 [measurements \(Liu et al., 2020; Ponte et al., 2021; Wong et al., 2023\), that is supposed to have](#)
 72 [a strong impact on the closure assessment of the regional sea level budget.](#) Considering all
 73 ocean basins, a recent study (Bouih et al., 2025) investigated the regional sea level trend budget
 74 closure over the 2004-2022 time span, using altimetry-based sea level data, GRACE space
 75 gravimetry for the regional ocean mass variations (also called manometric component, Gregory
 76 et al., 2019) and Argo data for the steric (sum of thermosteric and halosteric terms) component
 77 down to 2000m (global mean trends removed from all data sets). Bouih et al. (2025) also
 78 considered several ocean reanalyses to compute the manometric component (an alternative to
 79 using GRACE data), following the approach developed by Camargo et al. (2023), i.e., using
 80 the sterodynamic sea level (Gregory et al., 2019) computed by the reanalysis and correcting it
 81 for the local steric effect. The Bouih et al. (2025)’s study [used the Argo-based product from](#)

SIO (SCRIPPS Institution of Oceanography) that corrected for the spurious salinity drift reported recently in some Argo-based salinity measurements (Liu et al., 2020; Ponte et al., 2021; Wong et al., 2023). They showed that in the Pacific, Indian and South Atlantic oceans, the sea level budget trend residuals (altimetry-based sea level minus sum of components) were non-significant, considering the data uncertainties. On the other hand, strong positive residuals were found in the North Atlantic, whatever the manometric component considered (i.e., either from GRACE or from the ocean reanalyses). ~~These authors suspected the Argo-based spurious halosteric component (impacted by the Argo-based salinity drift) as the cause of the non-closure of the sea level budget in the North Atlantic.~~

In the present study we revisit the question of the North Atlantic sea level budget over the 2004-2022 time span, using a variety of different data sets for the components of the sea level budget: different GRACE mascon solutions and different Argo-based gridded products down to 2000m, as well as ~~antwo~~ ocean reanalyses that provide an estimate of the steric signal from the deep ocean. Our objective is to evaluate the impact of each product on the currently reported non-closure of the North Atlantic sea level budget. An important addition, compared to the Bouih et al. (2025)'s study, consists of accounting for the deep ocean warming below 2000m (not sampled by Argo) using estimates from the ocean reanalyses. This paper is organized as follows. Section 22 presents the data and analysis method. Section 33 displays the results for the North Atlantic sea level budget, both in terms of trend maps and time series. ~~The sea level budget excluding the North Atlantic (average over all other oceans) is also briefly discussed in that section.~~ A synthesis of the results is presented in the Discussion section (section 4) with some highlights on the few main messages arising from this study.

104 **2 Data and methods**

105 **2.1 Data**

106 *2.1.1 Altimetry-based total sea level*

107 Sea level variations have been continuously measured by satellite altimetry since 1993. In this
 108 study, we use the daily $0.25^\circ \times 0.25^\circ$ gridded sea level anomaly data, version DT2021 available
 109 from the Copernicus Climate Change Service (C3S) (<https://cds.climate.copernicus.eu/>). The
 110 data set is further corrected for the TOPEX-A instrumental drift that affected the first 6 years
 111 of the time series, but the correction has no impact on our assessment that starts in 2004 (Ablain
 112 et al., 2017). The Jason-3 radiometer drift that impacts the wet troposphere correction (Brown
 113 et al., 2023) is corrected for. The altimetry data set is also corrected for the absolute Glacial
 114 Isostatic adjustment (GIA) effect, using the ICE6G-D model from Peltier et al. (2018).

115 *2.1.2 GRACE-based ocean mass*

116 The Gravity Recovery and Climate Experiment (GRACE) and GRACE Follow-On (GRACE-
 117 FO) are joint missions by the National Aeronautics and Space Administration (NASA) and the
 118 German Aerospace Center (DLR) (Tapley et al., 2019). These satellites have been measuring
 119 temporal variations in Earth's gravity field since 2002. This data set is essential for estimating
 120 mass redistribution in the oceans, terrestrial water storage, ice sheets, and glaciers.
 121 GRACE/GRACE-FO data are generally available in two forms: Spherical Harmonic (SH)
 122 coefficients and Mass Concentration (mascon) solutions. In this study, we utilize the latest
 123 Release 6 (RL06) mascon solutions provided by three different institutions: the Center for
 124 Space Research (CSR, (Save et al., 2016)), the Jet Propulsion Laboratory (JPL, (Watkins et al.,
 125 2015)), and the Goddard Space Flight Center (GSFC, (Loomis et al., 2019)). These mascon
 126 solutions, at monthly temporal resolution, are corrected for the geocentric motion (degree-1)
 127 using the Sun et al. (2016) solution. The C_{20} and C_{30} coefficients are derived from Satellite

128 Laser Ranging (SLR) data. The GIA correction for GRACE is based on the ICE6G-D model
 129 from (Peltier et al. (2018)). Additionally, the GAD product derived from AOD1B models
 130 (Dobslaw et al., 2017; Flechtner et al., 2014), which represents non-tidal atmospheric and
 131 oceanic mass redistribution, is added back over the ocean areas to restore the ocean bottom
 132 pressure, which combines the effects of ocean mass and atmospheric loading. For alignment
 133 with altimetry-based sea level data which are corrected for the inverse barometer, we remove
 134 the global mean atmospheric pressure at every grid mesh in GRACE/GRACE-FO ocean mass
 135 data using the spatial mean of the GAD product at each month (Chen et al., 2019). In this study,
 136 we use the ensemble mean of the three mascon solutions.

137 *2.1.3 Argo-based gridded data*

138 The Argo program is an international observational network that deploys a global array of
 139 autonomous profiling floats to measure temperature and salinity in the upper 2000m of the
 140 ocean. Several institutions process the raw data from these floats and publish gridded
 141 temperature (T) and salinity (S) data sets. In this study, we utilize three distinct gridded
 142 products: the Scripps Institution of Oceanography (SIO, Roemmich and Gilson, 2009), the
 143 Japan Agency for Marine-Earth Science and Technology (JAMSTEC, Hosoda, 2007), and the
 144 Met Office Hadley Centre (EN4 product, version 2.2, Cheng et al., 2014). Note that the SIO
 145 product rejects salinity profiles exceeding a difference of 0.1 psu when comparing to historical
 146 estimate based on the WOCE Global Hydrographic Climatology (Roemmich and Gilson, 2009).
 147 This correction tends to withdraw the salinity drift reported in Argo floats since 2016 (Liu et
 148 al., 2020; Ponte et al., 2021; Wong et al., 2023). Note that the other two products do not apply
 149 such a correction. All three data products are available at [https://argo.ucsd.edu/data/data-](https://argo.ucsd.edu/data/data-access/)
 150 [access/](https://argo.ucsd.edu/data/data-access/) (downloaded in October 2025). The thermosteric, halosteric and total steric sea level
 151 time series are computed from these gridded temperature and salinity data using the Gibbs
 152 SeaWater (GSW) Oceanographic Toolbox (McDougall et al., 2011) which implements the

153 2010 Thermodynamic Equation Of Seawater (TEOS-10) standard. The gridded Argo-based
 154 time series have a spatial resolution of $1^{\circ} \times 1^{\circ}$ at monthly interval over January 2004 to
 155 December 2022.

156 2.1.4 Ocean reanalyses

157 We also used ~~two~~the ocean reanalysis data sets: ~~t(1) the~~ CNR-ISMAR Global Historical
 158 Reanalysis (CIGAR, Storto and Yang, 2024), ~~and (2) the Ocean Reanalysis System 5 (ORAS5,~~
 159 ~~Zuo et al., 2019) available from the Copernicus Climate Change Service~~
 160 ~~(<https://www.copernicus.eu/en/copernicus-services>).~~

161 CIGAR is a reanalysis system developed by Storto and Yang (2024). Its resolution is $1^{\circ} \times 1^{\circ}$. It
 162 is based on the NEMO ocean model version 4.0.7 (<https://www.nemo-ocean.eu/>) and is forced
 163 by the ERA5 atmospheric reanalysis from the European Center for Medium-range Weather
 164 Forecast (ECMWF) (<https://www.ecmwf.int/en/forecasts/dataset/ecmwf-reanalysis-v5>). The
 165 system uses a three-dimensional variational (3D-Var) scheme to assimilate in situ profiles from
 166 the EN4 data set. Notably, CIGAR does not assimilate satellite altimetry data, which allows
 167 for independent comparisons. The system consists of 32 ensemble members generated through
 168 varying configurations (e.g., perturbed initial conditions and atmospheric forcing). In this study,
 169 we use the ensemble mean of the 32 members to derive the thermosteric, halosteric and steric
 170 sea level.

171 ~~(Bouih et al. (2025) considered the ensemble mean of five ocean reanalyses over 2004-2019,~~
 172 ~~including CIGAR, to compute the manometric component and showed that this ensemble mean~~
 173 ~~and CIGAR manometric components are highly similar. For that reason, here we only consider~~
 174 ~~CIGAR, but extend it until 2022.~~

175 ~~The ORAS5 ocean reanalysis is based on the NEMO model (v3.4.1) and uses the NEMOVAR~~
 176 ~~data assimilation system. It assimilates a wide range of observations, including satellite~~
 177 ~~altimetry and sea surface temperature, at an eddy-permitting resolution of $0.25^{\circ} \times 0.25^{\circ}$.~~

178 2.3.2.2 *Methods*

179 2.3.12.2.1 *Computation of the manometric component*

180 Manometric sea level change generally refers to the sea-level component associated with ocean
 181 mass variations. It is driven by water mass exchange with the continents, such as land ice melt
 182 and terrestrial water storage changes, as well as by mass redistribution within the ocean driven
 183 by ocean circulation. In this study, because the global mean ocean mass trend is removed, the
 184 manometric sea level refers to the regional redistribution of ocean mass around the global
 185 mean, with the GRD effect added as described below. In this study, we utilize two methods to
 186 obtain an estimate of the manometric component. One method is to use the ensemble mean of
 187 the GRACE mascon solutions from CSR, JPL, and GSFC to obtain the GRACE-based
 188 manometric estimate. The other method, an alternative to using GRACE solutions, relies on
 189 ocean reanalysis data and follows the approach developed by Camargo et al. (2023). Here, we
 190 use the sterodynamic sea level provided by the CIGAR ocean reanalysis, which combines steric
 191 and dynamic sea level changes. To isolate the manometric component, we subtract the local
 192 steric effect from the sterodynamic sea level and further add the contemporary Gravitational,
 193 Rotational, and Deformation (GRD) fingerprints (representing solid Earth deformations and
 194 gravitational effects due to present-day land ice melt and terrestrial water storage changes;
 195 Gregory et al., 2019), to obtain a manometric component comparable to GRACE. The sea level
 196 fingerprint data used in this study are based on monthly GRD fingerprint grids ($0.5^\circ \times 0.5^\circ$
 197 resolution) estimated by Adhikari et al. (2019). Since this original dataset ends in 2016, we
 198 extended the time series up to 2022 using linear extrapolation, assuming that the observed trend
 199 remains constant after 2016 (see Bouih et al., 2025 for details). This reanalysis-based estimate
 200 is hereafter referred to as CIGAR manometric sea level. ~~Manometric sea level change generally~~
 201 ~~refers to the sea-level component associated with ocean mass variations. It is driven by water~~
 202 ~~mass exchange with the continents (such as land ice melt and terrestrial water storage changes)~~

~~as well as mass redistribution within the ocean, driven by ocean circulation. However, in the context of this study, we focus exclusively on the internal mass redistribution component (with the global mean ocean mass trend removed). Therefore, the 'manometric sea level' here refers solely to the redistribution by the ocean circulation, of water mass already present in the oceans, with the GRDs effect added as explained above.~~

2.3.22.2.2 *Post processing of the data*

To ensure spatial consistency across all observing systems, all gridded data sets were spatially interpolated onto a $1^{\circ} \times 1^{\circ}$ grid and averaged at monthly interval. A three-month moving average filter was applied to the time series. For the spatial analysis, a common mask was applied to all gridded components to exclude regions with high uncertainty. This mask covers latitudes from 66°S to 66°N, excludes inland seas, and omits coastal regions where the distance from land is less than 300 km (see Bouih et al., 2025 for details). Seasonal signals (annual and semi-annual) are removed from the time series through a least-squares adjustment of 6 and 12-month sinusoids. For the spatial maps, this [linear](#) least-squares fit was used to calculate the trends at each grid point. Finally, the globally averaged trend of each data set computed over the study period was subtracted from each grid point before constructing the spatial trend maps to focus on regional spatial patterns. The study period spans from January 2004 to December 2022.

2.3.32.2.3 *Data uncertainties*

For the altimetry [datagrids](#), we use the uncertainties estimated by Prandi et al. (2021). For the GRACE-based manometric component, uncertainties [can beare](#) derived from the dispersion of the data sets with respect to the ensemble mean. For the CIGAR-based manometric, thermosteric, halosteric and steric data, the dispersion of the 32 realizations around the

ensemble mean was used to estimate the uncertainties. For the residual time series, the uncertainty was estimated by quadratically summing the error of each term of the budget equation using the law of error propagation, considering all terms of the sea level budget. Regarding the trend uncertainty, it is expressed as the standard error of the least-squares fit. To obtain the uncertainty at the 95% confidence level, we scaled the standard error by a factor of 2 (representing the 2-sigma interval).

3 Results

3.1 Comparison of the different manometric components over the North Atlantic

Figure 1a, b, shows the spatial manometric trend in the North Atlantic using the GRACE and CIGAR manometric data. Comparing Figure 1(a) and (b), we distinctly observe contrasting patterns between the GRACE and CIGAR manometric components. For the GRACE-based component, significant negative trends dominate most of the basin, with magnitudes gradually weakening from east to west. However, positive trends are observed in the western tropical Atlantic. In contrast, the CIGAR manometric component displays positive trends in the east and negative trends in the west, separated by a distinct dividing line, which likely corresponds to the position of the Mid Atlantic Ridge.

Figure 1c also shows the ‘Altimetry minus SIO-based steric 0-2000m’ trends. This map is supposed to represent the manometric component, plus any deep ocean contribution. Comparing Figure 1c with Figure 1a-b, we note a better agreement between ‘Altimetry minus SIO’ and CIGAR manometric in the Eastern Atlantic, both maps showing positive trends, while the GRACE manometric trends are negative in this region. On the other hand, it is worth noting that the positive trend area seen in the GRACE manometric map in the western Atlantic around 30°N is also seen in the ‘Altimetry minus SIO’ trend map, but not in the CIGAR manometric map.

Over-all, such comparisons point to problems in GRACE data in the eastern North Atlantic region.




Figure 1. Manometric sea level trends in the North Atlantic over 2004.01–2022.12 derived from (a) the mean of three GRACE mascon solutions, (b) the CIGAR ocean reanalysis and (c) ‘Altimetry minus SIO-based steric sea level 0–2000m’.

3.1 Comparison of the different steric products over the North Atlantic


3.1.1 Upper ocean (0–2000m) thermosteric, halosteric and steric products


Figure 2 Figure 1 Figure 1a–de shows the thermosteric sea level (TSL) trends over 2004–2022 over the North Atlantic for the three Argo products SIO, JAMSTEC and EN4, and CIGAR (0–2000m) reanalysis data. Similarly, Figure 3cd–hf shows the halosteric sea level (HSL) trends for the same data sets and period. As expected, due to thermohaline compensation particularly strong in the North Atlantic (e.g., Pardaens et al., 2011; Wang et al., 2010; Wunsch et al., 2007), the two components display opposite trends. Regarding the thermosteric component (Figure 3 Figure 1 Figure 1aa–d–e), while all four three products display similar spatial patterns, their magnitudes vary. especially The JAMSTEC and EN4 products show stronger positive trends around 20°N compared to the SIO product. However,

271 ~~all three products~~ indicate significant positive signals in the ~~w~~eastern North Atlantic. ~~The~~
 272 ~~three~~ Regarding the halosteric ~~component, the four~~ products share similar spatial patterns;
 273 however, the magnitude of the SIO product is ~~notably~~ smaller than ~~the other three~~
 274 ~~JAMSTEC and EN4~~, the ~~others~~ ~~latter two~~ being very similar. This difference likely results from
 275 the SIO processing that corrects for the instrumental salinity drift reported in Argo floats since
 276 2016, whereas the other products do not account for this. ~~The mapping methods used by the~~
 277 ~~different processing groups, in particular the assumed different correlation radii, may also~~
 278 ~~contribute to the observed differences.~~
 279 The steric sea level (SSL) trends are shown in ~~Figure 1~~ ~~Figure 1~~ ~~Figure 3g-i-1~~. The magnitude
 280 of the total steric change is smaller than that of the thermosteric component alone. This implies
 281 that the halosteric contribution plays a significant compensatory role in the North Atlantic
 282 (consistent with previous published studies; e.g., Bouih et al., 2025; Llovel and Hochet, 2025).
 283 Unlike in the case of the global mean, the influence of salinity changes on sea level cannot be
 284 ignored in this region.

285

286





288

289 Figure 1+2. Trends of thermosteric (a–d), halosteric (e–h), and steric (i–l) sea level
 290 change in the North Atlantic (0–2000m depth) over 2004–2022 derived from the SIO (a, e, i),
 291 JAMSTEC (b, f, j), and EN4 (c, g, k) Argo products and CIGAR (d, h, l) ocean
 292 reanalysis.


293

294 3.9.13.1.2 Ocean reanalyses-based Full depth thermosteric, halosteric and steric products

295 Here we In addition to Argo products, we used the CIGAR and ORAS5 ocean reanalysis data
 296 to compute thermosteric, halosteric, and steric sea level changes for both the the deep ocean
 297 (upper 2000m and the deep ocean below (2000 mm–6000m)). We consider two cases: (1) SIO
 298 upper ocean (0–2000) combined with CIGAR deep ocean (below 2000 m). Hereafter it is called
 299 SIO hybrid. (2) CIGAR full depth. Trends of thermosteric, halosteric, and steric full depth for
 300 SIO hybrid and CIGAR full depth over 2004–2022 are shown in Figure 2Figure 2Figure 3.
 301 In the upper 2000m, CIGAR and ORAS5 exhibits similar spatial patterns across all three

302 ~~components. Notably, the thermosteric sea level change displays a significant signal in the~~
303 ~~western North Atlantic, consistent with the Argo based results. However, marked~~
304 ~~Differences/repancies are seen between the two cases reanalyses in~~ around 20N° for the
305 ~~thermosteric and halosteric evident below 2000m.components. Both The CIGAR products~~
306 displays strong positive thermosteric and negative halosteric signals in the western North
307 Atlantic. ~~compared to ORAS5. In contrast, the deep ocean signals in ORAS5 are weak. The~~
308 ~~contrasting results of the two reanalyses for the deep ocean are puzzling and need to be~~
309 ~~investigated further.~~

310






Figure 223. Trends of thermosteric (a, b–d), halosteric (c, de–h), and steric (e, ffi–l) for SIO hybrid (a, c, e) and CIGAR full depth (b, d, f) sea level change in the North Atlantic over 2004–2022. The panels compare contributions from the upper 2000 m (a, b, e, f, i, j) and full depth below 2000 m (c, d, g, h, k, l) derived from the CIGAR (a, c, e, g, i, k) and ORAS5 (b, d, f, h, j, l) reanalyses. **TO BE REDRAWN**


3.9.33.2 Upper ocean Argo-based and CIGAR-based ocean reanalysis thermosteric, halosteric and steric time series over 2004–2022 (global mean trend removed)

Next, we examine the thermosteric, halosteric and steric time series (weighted averages of the gridded data in the North Atlantic) for the different Argo and ocean reanalysis product (Figure


326 ~~3Figure 3Figure 4~~). ~~Figure 3Figure 3Figure 4a,d~~ indicates that the thermosteric component ~~(0-~~
 327 ~~2000m)~~ displays ~~exhibits~~ a positive trend in all products ~~as of about 2010, although the rates~~
 328 ~~of rise vary~~. Regarding the halosteric component ~~-(0-2000m)~~ (~~Figure 3Figure 3Figure 4b,e~~),
 329 we observe that the ~~JAMSTEC curve exhibits spurious positive anomaly between 2008 and~~
 330 ~~2012, while the EN4 curve (and to a little extent also SIO and JAMSTEC) shows a strong~~
 331 ~~decrease as of 2016, possibly linked to the salinity drift in EN4 and JAMSTEC data sets, to~~
 332 ~~together with a potential physical salinity increase. The latter issue needs further investigation.~~
 333 ~~SIO and CIGAR products show a small negative trend that begins earlier than 2016, which~~
 334 ~~may represent a true physical salinity increase during this early period. Among the three Argo~~
 335 ~~products (Figure 3Figure 4a-c), SIO and EN4 show strong agreement in thermosteric,~~
 336 ~~halosteric, and steric sea level changes. In contrast, JAMSTEC deviates from the other two~~
 337 ~~products, particularly in the halosteric component, which displays an unrealistic increase~~
 338 ~~during the 2010s. Consequently, this discrepancy causes the total steric sea level change in~~
 339 ~~JAMSTEC to mismatch with the other two products. In the following, we will only consider~~
 340 ~~the SIO product (as mentioned before, corrected for salinity drift). exclude the JAMSTEC~~
 341 ~~product from the subsequent analysis.~~

342 ~~Figure 4d-i~~ presents the changes for CIGAR and ORAS5 in the upper 2000m ocean depth and
 343 ~~full depth from 2000m to 6000m (deep ocean). In the upper 2000m, a~~Although there is a little
 344 ~~difference in trend before 2010, they are consistent across components after 2010, although~~
 345 ~~their amplitudes slightly differ. However, in the deep ocean (2000–6000m), substantial~~
 346 ~~discrepancies exist between the two products. As mentioned above, ORAS5 exhibits a weak~~
 347 ~~signal in the deep ocean; its thermosteric trend is nearly zero until 2019. In contrast, CIGAR~~
 348 ~~shows a consistent and distinct upward trend. Regarding the halosteric change, ORAS5 shows~~
 349 ~~a small rising trend until 2019, whereas CIGAR shows a consistent and obvious decrease.~~
 350 ~~Finally, while both products exhibit rising trends in steric sea level change in the deep ocean,~~
 351 ~~their magnitudes differ significantly. Given the unrealistically stationary thermosteric signal~~

352 observed in ORAS5 below 2000m, we suspect that this product is unreliable in its deep-ocean
353 variability.
354



355



356

Figure 334. Upper ocean time series of thermosteric (a, d, g), halosteric (b, e, h), and steric (c, f, i) sea level time series change in the North Atlantic. Panels (a–c) show the upper 2000 m derived from Argo products (SIO, JAMSTEC/JAM, EN4), while panels (d–i) are derived from CIGAR and ORAS5 reanalyses for the upper 2000 m (d–f) and the deep ocean 2000–6000 m (g–i). All time series have applied three-month moving average filtering. Global mean trend is included.

Comparison between the SIO steric trends and altimetry sea level trends corrected for the manometric component

3.3 Comparison of the different manometric components over the North Atlantic

Figure 4a, b, shows the spatial manometric trends in the North Atlantic using the GRACE and CIGAR manometric data. Comparing Figure 4(a) and (b), we distinctly observe contrasting patterns between the GRACE and CIGAR manometric components. For the GRACE-based component, significant negative trends dominate most of the basin, with magnitudes gradually weakening from east to west. However, positive trends are observed in the western tropical Atlantic. In contrast, the CIGAR manometric component displays positive trends in the east and negative trends in the west, separated by a distinct dividing line, which likely corresponds to the position of the Mid Atlantic Ridge.

Figure 4c, d also shows the ‘Altimetry minus steric full depth’ trends for both the SIO hybrid case and CIGAR. These maps are supposed to represent the manometric component, plus the deep ocean contribution. Comparing Figure 4c, d with Figure 4a, b, we note a better agreement between ‘Altimetry minus steric full depth’ and CIGAR manometric in the Eastern Atlantic, both maps showing positive trends, while the GRACE manometric trends are negative in this region. On the other hand, it is worth noting that the positive trend area seen in the GRACE manometric map in the western Atlantic around 30°N is also seen in the ‘Altimetry minus steric full depth’ trend map, although with smaller extension, but not in the CIGAR manometric map.

Overall, such comparisons seem point to problems in GRACE data in the eastern North Atlantic region.

385 A recent study by Hightower et al. (2026) revisited the GIA correction applied to GRACE
 386 data. They developed a 3D GIA model, considering 3D viscosity fields derived from
 387 tomography and two commonly used ice loading histories. While traditional GIA models
 388 assume—A recent study by Hightower et al. (2026) revisited the GIA correction applied to
 389 GRACE data. They developed a 3D GIA model, considering 3D viscosity fields derived from
 390 tomography and two commonly used ice-loading histories. While traditional GIA models
 391 assume laterally homogeneous mantle viscosity, these authors show that because the Earth's
 392 rheology varies significantly -from place to place, this strongly affects how the Earth's mantle
 393 rebounds after ice melts, with important impact on the GIA correction applied to GRACE-
 394 based ocean mass data in many regions, in particular in the whole Northeast Atlantic. -Since
 395 here we use GRACE data corrected for the traditional GIA correction, the discrepancy
 396 between the GRACE and CIGAR manometric trend maps may result from an inadequate
 397 GIA correction.




Figure 4. Manometric sea level trends in the North Atlantic over 2004.01-2022.12 derived from (a) the mean of three GRACE mascon solutions, (b) the CIGAR ocean reanalysis, (c) 'Altimetry minus SIO hybrid' and (d) 'Altimetry minus CIGAR full depth'.

Figure 4. Manometric sea level trends in the North Atlantic over 2004.01-2022.12 derived from (a) the mean of three GRACE mascon solutions, (b) the CIGAR ocean reanalysis and (c) 'Altimetry minus SIO-based steric sea level 0-2000m'. TO BE REDRAWN

In addition to directly obtaining the steric sea level trends, we can also derive it indirectly using the sea level budget equation (i.e., altimetry sea level minus manometric component computed at each grid mesh of the gridded products). In Figure 5, we compare the direct steric estimate from SIO with indirect estimates derived by subtracting different manometric products (GRACE or CIGAR) from altimetry data. It is important to note that the indirect method inherently includes deep ocean contribution and relies on independent observations. From Figure 5, we clearly observe that the spatial trend pattern of SIO is more consistent with the "Altimetry minus CIGAR manometric" estimate, as both exhibit positive signals in the western North Atlantic. In contrast, the "Altimetry minus GRACE manometric" estimate displays a prominent positive signal in the eastern North Atlantic which is absent in the SIO data. This suggests that the GRACE manometric data contains substantial inaccuracies, particularly in the eastern North Atlantic, while "Altimetry minus CIGAR manometric" agrees rather well with the SIO steric trends. Such a result confirms previous finding based on the comparison between GRACE manometric trends and 'Altimetry minus SIO-based steric' trends that already highlighted potential errors in GRACE data in the eastern North Atlantic (section 3.1).




Figure 5. Steric sea level trends derived directly from SIO and indirectly from Altimetry minus manometric estimates (GRACE or CIGAR).

To highlight this further, Figure 6 Figure 5a, b shows the scatter plots between ‘Altimetry minus steric full depth’ for SIO hybrid and CIGAR full depth SIO gridpoint steric trends and “Altimetry minus GRACE manometric” and “Altimetry minus CIGAR manometric” trends using the gridded datasets in all cases. grid point trends. The data have been normalized.

ADD A FEW SENTENCES ABOUT THE CORRELATIONS

The scatter plots reveal distinct differences in the correlation between xxx SIO observations and xxx Altimetry minus the two manometric estimates. In the negative SIO region (corresponding to the lower half of Figure 6 and the negative value regions in Figure 5), where most points align with the diagonal in both cases, the “Altimetry CIGAR manometric” estimates points are clustered notably more tightly than those in the “Altimetry GRACE manometric” comparison. This indicates that in regions with negative steric sea level change, where both estimates correlate with SIO, the “Altimetry CIGAR manometric” estimate maintains a higher spatial correlation compared to “Altimetry GRACE manometric”. The abundance of data points in the quadrant characterized by negative SIO and positive ‘Altimetry GRACE manometric’ values reflects the situation eastern North Atlantic (Figure 5), where negative SIO observations are paired with positive “Altimetry GRACE manometric” estimates. In the positive SIO areas

(corresponding to the upper half of the plots), a portion of the data points in both estimates fall below the 1:1 correspondence line (indicated by points falling above the diagonal). However, this deviation is much more pronounced for “Altimetry-GRACE manometric”, leading to a larger spread away from the diagonal compared to “Altimetry-CIGAR manometric”. This suggests that while the linear consistency with SIO decreases for both products in positive steric sea level trend regions (such as the western North Atlantic), the correlation of “Altimetry-GRACE manometric” is notably weaker than that of “Altimetry-CIGAR manometric”.




Figure 546. Scatter plots of SIO steric sea level change versus Altimetry minus manometric estimates (GRACE/left or CIGAR/right). The data have been normalized and outliers exceeding 3σ were removed. TO BE REDRAWN

461 ~~3.11.3.4~~ *Residual trend maps in the North Atlantic* ~~(global mean trends removed)~~

462 In this section, we utilize data from the observing systems described above to compute spatial
463 maps of the sea level budget residuals.

464 ~~3.11.13.4.1~~ *North Atlantic budget residuals for* GRACE and CIGAR ~~different~~ *manometric and* 465 *different* *steric (0-2000m* and full depth *) components*

466 We computed the residuals by subtracting various combinations of manometric (GRACE,
467 CIGAR) and steric (SIO, ~~EN4,~~ and CIGAR, ~~and ORAS5~~ upper 2000 m and full depth)
468 components from the altimetry data (Figure 5 ~~Figure 7~~ Figure 6).

469

470 ~~As shown in Figure 5a,b,e,f (upper ocean), tFigure 6Figure 7a,e, the resulting~~ residuals
471 implicitly include the a-deep ocean steric contribution, ~~which is not accounted for since the~~
472 ~~steric data are limited here to the upper 2000 m. This is consistent with~~As previously s
473 ~~observed results by from Bo~~(Bouih et al., (2025) ~~uuh et al. (2025), who observed strong~~positive
474 residuals are seen in the Northeast Atlantic with GRACE, and in the Northwest Atlantic with
475 CIGAR. for both manometric cases.When the deep ocean accounted for (Figure 5c,d,g,h),
476 strong residuals are still seen in the Northeast Atlantic with GRACE, while the residual signal
477 is much attenuated with CIGAR.Similarly positive residuals are found in the eastern North
478 Atlantic for the GRACE manometric component and in the western North Atlantic for the
479 CIGAR manometric component.

480 Indeed, when using the CIGAR manometric component, the residuals decrease significantly
481 in both spatial variability and magnitude.

482

483

484 ~~From Figure 7Figure 6, we also observe that the spatial distribution of the residuals remains~~
485 ~~similar when the manometric component is held fixed while varying the steric components.~~

486 However, the residual magnitude is different. Specifically, when using the GRACE
487 manometric component, the positive signal derived from SIO is stronger in the eastern North
488 Atlantic compared to xxxxother steric products. When using the CIGAR manometric
489 component, the positive signals from SIO and CIGAR are stronger in the western North
490 Atlantic compared to EN4 and ORAS5.
491 To quantify the residuals, we computed the latitude-weighted spatial mean trend of each map.
492 Results are shown in Table 1 (remind that results in Table 1 do not account for the global mean
493 trends). From Table 1, we observe that regardless of the steric data used (whether from Argo
494 or ocean reanalyses), the residuals derived using the CIGAR manometric component are
495 always smaller than those derived using GRACE. Furthermore, for a given manometric
496 component, the residual with SIO is larger than that with EN4, even though SIO has corrected

for the salinity drift since 2016. Besides, the residual with CIGAR steric is larger than that with ORASS.




Figure 5. Maps of sea level residuals (mm/yr) in the North Atlantic. The residuals are calculated by subtracting manometric and steric components from satellite altimetry. The manometric component is derived from GRACE (a–d) and CIGAR (e–h). The steric component from SIO (a, e), SIO hybrid CIGAR DO (b, f), CIGAR UP (upper ocean, c, g) and CIGAR full depth (d, h).

~~Figure 76. Maps of sea level residuals (mm/yr) in the North Atlantic. The residuals are calculated by subtracting manometric and steric components from satellite altimetry. The manometric component is derived from GRACE (a–d) and CIGAR (e–h). These are combined with upper 2000 m steric estimates from SIO (a, e), EN4 (b, f), CIGAR (c, g), and ORAS5 (d, h). TO BE REDRAWN~~

To quantify the residual trends, we computed the latitude-weighted spatial mean trend of each map. Results are shown in Table 1. Regardless of the steric data used (whether from SIO or CIGAR), the residuals derived using the CIGAR manometric component are always smaller than those derived using GRACE, for both upper ocean and full depth.

Table 1. Trend values (mm/yr) of sea level residuals in the North Atlantic corresponding to different steric product SIO and CIGARs (0–2000m) for upper ocean and deep ocean, and the two manometric components. Global mean trends were removed. Uncertainties are 2-sigma errors of the least-squares fit.

Trend (mm/yr)	SIO	CIGAR SIO Hybrid EN4	SIO hybrid CIGAR	CIGAR full depth ORAS5
GRACE		1.52 ± 0.21	1.53 ± 0.21	0.90 ± 0.23
manometric	2.15 ± 0.19	± 0.22	0.21	0.20
CIGAR		0.86 ± 0.25	0.88 ± 0.25	0.24 ± 0.22
manometric	1.50 ± 0.23	± 0.23		0.16

3.11.1 North Atlantic budget residuals and deep ocean steric contribution

We further In this section, we added the deep ocean steric sea level change to the upper 2000 m steric estimates, considering both the SIO hybrid and CIGAR full depth cases, and ORAS5 deep steric data, and using either GRACE or CIGAR for the manometric components.

Resulting residual trends are also shown in Figure 8b. For all cases except the ORAS5-deep ocean case, the residuals derived using GRACE manometric data (Figure 8a-c) show a dipole pattern (positive in the east, negative in the west) compared to the results without the deep ocean component (Figure 7Figure 6). The Table 2 summarizes the residual trend values for all cases are also gathered in (as for Table 1., remind that results in Table 2 do not account for the global mean trends). Comparing results from Table 1 and Table 2, We note that the mean residual magnitude is reduced when the deep-ocean steric change is included: by about 30% when using the GRACE manometric component with SIO upper-2000 m steric sea level, and by about 70% when using the CIGAR manometric component together with the CIGAR full-depth steric component. We note that the mean residual magnitude is significantly reduced when the deep-ocean steric change is included: by ~30% when using GRACE for the manometric component and SIO for the upper 0-2000m steric sea level, and by ~50% when using CIGAR for both the steric (full depth and manometric component). Based on CIGAR, the deep steric contribution to the North Atlantic sea level budget (in addition to the global mean deep-ocean contribution) amounts to 0.62 ± 0.04 mm/yr. In contrast, when using the CIGAR manometric component (Figure 8e-h), the residuals decrease significantly in both spatial variability and magnitude. Notably, the sea level budget is effectively nearly closed within error bars when combining CIGAR manometric and CIGAR steric data. Finally, it is worth noting that for SIO, positive residuals are observed in the eastern North Atlantic regardless of whether GRACE or CIGAR manometric data is used.




Figure 8. Maps of sea level residuals (mm/yr) in the North Atlantic, calculated as in Figure 7 but with the inclusion of the deep ocean (below 2000 m) steric component. The deep ocean contribution is derived from CIGAR for panels (a–c, e–g) and from ORAS5 for panels (d, h). In panels a–c, the manometric component is from GRACE while it is from CIGAR in panels e–h.

Table 2. Trend values (mm/yr) over 2004–2022 of sea level residuals in the North Atlantic for four steric products and two manometric components, accounting for the deep ocean. For SIO and EN4, the deep ocean contribution from CIGAR is added. For CIGAR and ORAS5, their own deep steric contributions are considered. Global mean trends removed. Uncertainties are 2-sigma errors of the least squares fit.

Trend (mm/yr)	SIO + CIGAR	EN4 + CIGAR	CIGAR	ORAS5
	Deep-ocean	Deep-ocean	Full depth	Full depth
GRACE				
manometric	1.53 ± 0.21	0.81 ± 0.24	0.90 ± 0.23	0.87 ± 0.20
CIGAR				
manometric	0.88 ± 0.20	0.15 ± 0.21	0.24 ± 0.22	0.22 ± 0.15

3.5 Time series of the North Atlantic mean sea level budget (*global mean trends removed*)

Figure 6 displays the North Atlantic budget time series using CIGAR for the manometric component and two steric full depth configurations. Figure 6 shows that the CIGAR manometric component display a changing trend as of 2016, which origin remains to be explained. Because of this behavior, the residual curves are not purely linear. However, Figure 6 clearly shows a larger positive trend with SIO hybrid than with CIGAR full depth. Corresponding residual linear trends amount to 0.88 ± 0.20 mm/yr for the hybrid steric case and 0.24 ± 0.22 mm/yr for the CIGAR full-depth case. In the latter case, we may conclude that the North Atlantic sea level budget is closed within data uncertainties.

~~Based on the analysis above, the deep ocean contribution to the North Atlantic sea level budget cannot be ignored. To assess the influence of the deep ocean on the sea level budget, we computed time series for each component, including total sea level, manometric sea level, thermosteric sea level, and halosteric sea level. Compared to results discussed above, here we account for the global mean trend of each component.~~

~~North Atlantic mean sea level budget time series without deep ocean (global mean trends included)~~

~~Here, we have reintroduced the global mean trend of each component. Since the CIGAR manometric component does not fully account for the external sea water mass addition to the ocean due to land ice melt and terrestrial waters, we adopt GRACE data to represent the total ocean mass change (i.e., global mean trend accounted for). Regarding the steric sea level component, we selected the SIO product for Argo estimates, as it corrects for the salinity drift. For ocean reanalyses, we only consider the CIGAR reanalysis.~~

~~Figure 9 presents the sea level budget time series (global mean trends accounted for) in the North Atlantic without the deep ocean steric sea level change. Residual time series are also shown. The left/right panels use SIO/CIGAR steric sea level change in upper 2000m.~~

588 Comparing Figure 9 left and right panels, we observe that the halosteric components from SIO
589 and CIGAR slightly differ. For SIO, a slight halosteric decrease is observed over 2012–2015,
590 with a linear trend of -1.10 ± 0.94 mm/yr during the such period, then it increases to -3.20 ± 0.9
591 mm/yr over 2016–2022. Consequently, the SIO halosteric trend over 2012–2022 amounts to $-$
592 2.87 ± 0.21 mm/yr. In terms of global mean trend, the SIO halosteric time series does not show
593 any trend (as expected since the salinity drift is supposed to be corrected). Thus, the SIO
594 halosteric decrease observed in the North Atlantic may possibly reflect a true physical salinity
595 increase. As of 2016, the CIGAR halosteric decrease is slightly larger than for SIO, possibly a
596 consequence of the salinity drift error not corrected for in the assimilated EN4 salinity data.
597 The mean residual trends amount to 2.72 ± 0.19 mm/yr and 2.34 ± 0.21 mm/yr for the SIO and
598 CIGAR steric cases respectively, i.e., not significantly different. Results are gathered in Table
599 3.

600




Figure 9. Time series of North Atlantic sea level components (top) and residuals (bottom) for the upper 2000m. The analysis uses Altimetry and GRACE data, combined with steric estimates from SIO (left) and CIGAR (right). All time series have applied three-month moving average filtering. The global mean trend of each component is included.

Table 3. Components of the North Atlantic sea level budget over 2004-2022
(upper ocean 0-2000m, Global mean trends included)

Components of the North Atlantic	Linear trends (mm/yr)
Altimetry	4.45 ± 0.16
GRACE ocean mass	1.03 ± 0.13
Thermosteric SIO (0-2000m)/CIGAR (0-2000m)	$1.76 \pm 0.27 / 1.92 \pm 0.29$
Halosteric SIO (0-2000m)/CIGAR (0-2000m)	$-1.01 \pm 0.17 / -0.8 \pm 0.19$
Sum of steric and mass components SIO (0-2000m)/CIGAR (0-2000m)	$1.73 \pm 0.19 / 2.11 \pm 0.22$
Residuals SIO (0-2000m)/CIGAR (0-2000m)	$2.72 \pm 0.19 / 2.34 \pm 0.21$

~~North Atlantic sea level budget time series with the deep ocean contribution (global mean trends included)~~

~~In this section, we add the deep ocean contribution from the CIGAR reanalysis. As in section 3.5.1, the GRACE-based ocean mass component is considered, and for the upper 2000m steric component, both SIO and CIGAR data are used. The corresponding sea level budget time series for the North Atlantic are shown in Figure 10.~~

~~As illustrated in Figure 10, with the inclusion of the deep ocean contribution, the halosteric decline in both panels become more evident compared to the case without the deep ocean. Specifically, the CIGAR component also shows a slight decrease starting in 2012. However, unlike the SIO trend, which maintains a relatively stable decline throughout the study time span, the CIGAR decrease still exhibits a sudden intensification after 2016. Similarly, the thermosteric sea level also displays a stronger positive trend compared to the case without the deep ocean.~~

Consequently, the total steric sea level trend is larger than in the case without the deep-ocean contribution. Finally, the inclusion of the deep-ocean contribution leads to a reduction in the North Atlantic residual trends for both cases: for the estimate based on SIO (upper 2000 m), the trend decreases by about 30%, from 2.72 ± 0.19 mm/yr to 1.89 ± 0.21 mm/yr, whereas for the estimate based on CIGAR, it drops from 2.34 ± 0.21 mm/yr to 1.51 ± 0.23 mm/yr. Corresponding results are presented in Table 4.




Figure 10. Time series of North Atlantic sea level change components (top) and budget residuals (bottom). The left panels use a hybrid steric estimate (SIO for 0–2000m combined with CIGAR for 2000–6000m), while the right panels use the full-depth CIGAR steric estimate (0–6000m). In both cases, Altimetry and GRACE data are used. All time series have applied three-month moving average filtering. The global mean trend of each component is included.

Table 4. Components of the North Atlantic sea level budget over 2004–2022
(full depth 0–6000m, Global mean trends included)

Components of the North Atlantic	Linear trends (mm/yr)
Altimetry	4.45 ± 0.16
GRACE ocean mass	1.03 ± 0.13
Thermosteric SIO (0–2000m) + CIGAR (2000–6000m)/CIGAR Full depth	$4.47 \pm 0.24 / 4.64 \pm 0.27$
Halosteric SIO (0–2000m) + CIGAR (2000–6000m)/CIGAR Full depth	$-2.9 \pm 0.16 / -2.68 \pm 0.19$
Sum of steric and mass components (with SIO 0–2000m + CIGAR 2000–6000m)/CIGAR Full depth	$2.56 \pm 0.19 / 2.94 \pm 0.22$
Residuals (with SIO 0–2000m + CIGAR 2000–6000m)/CIGAR Full depth	$1.89 \pm 0.21 / 1.51 \pm 0.23$

~~3.35.2 Sea level budget of all oceans but without the North Atlantic Ocean, with the deep ocean contribution added (global mean trends included).~~

~~In Bouih et al. (2025)’s study, the sea level budget for the global ocean excluding the North Atlantic (without the deep ocean steric contribution) was not closed. In this section, we consider the deep ocean steric sea level change contribution to the sea level budget. We compute the sea level budget for the global ocean excluding the North Atlantic using the same data products as in Section 3.5.2. The corresponding results are shown in Figure 11.~~

~~In the left panel (using SIO data for the upper 2000 m), we observe that the halosteric sea level change is nearly zero. The residual is very small, estimated at 0.16 ± 0.08 mm/yr. Thus,~~

considering the whole oceanic domain without the North Atlantic, the sea level budget is closed when SIO data are used for the upper 2000 m ocean layer. In contrast, the right panel (using CIGAR data for all depths) shows a distinct decrease in halosteric sea level starting in 2016, due to the assimilation of EN4 data in CIGAR (not corrected for the salinity drift). Nevertheless, the residual trend of the right panel is small (0.51 ± 0.11 mm/yr), suggesting quasi closure of the sea level budget.




Figure 11. Time series of sea level components (top) and residuals (bottom) for the global ocean excluding the North Atlantic. The left panels use a hybrid steric estimate (SIO for 0–2000 m combined with CIGAR for 2000–6000m), while the right panels use the full depth CIGAR estimate (0–6000m). All time series have applied three-month moving average filtering. The global mean trend of each component is included.

664 *3.35.3 North Atlantic sea level budget time series with the deep ocean contribution (global mean*
 665 *trends removed)*

666 Here, we compute the North Atlantic sea level budget (including the deep ocean contribution)
 667 with the global mean sea level trends of all components removed. The objective is to determine
 668 how errors in the regional components impact which part of the non-closure of the North
 669 Atlantic sea level budget, may result from errors in the regional components (i.e.,
 670 independently from errors in the global mean trends). Figure 12 presents the budget time series
 671 using CIGAR GRACE for the manometric component. We consider two steric cases: the
 672 hybrid case (SIO 0–2000 m plus CIGAR deep ocean) and the CIGAR full-depth case.

673 Table 2 summarizes the trend values for the components and residuals.

674 The mean residual trends are estimated at 1.53 ± 0.21 mm/yr for the hybrid steric case and
 675 0.88 ± 0.20 mm/yr for the CIGAR full-depth case. Comparing these results (where global mean
 676 trends are removed) with those in Figure 10 (where global mean trends are included), we
 677 observe very little difference in the mean residual trends. This suggests that errors in the global
 678 mean trends are small, and that the reported residuals in the North Atlantic budget are primarily
 679 driven by component errors at the regional scale.

680





Figure 12. Time series of North Atlantic sea level change components (top) and budget residuals (bottom). The left panels use a hybrid steric estimate (SIO for 0–2000m combined with CIGAR for 2000–6000m), while the right panels use the full-depth CIGAR steric estimate (0–6000m). In both left and right panels, GRACE manometric components are considered. All time series have applied three-month moving average filtering. The global mean trend of each component is removed.


Figure 13 displays the corresponding budget time series using CIGAR for the manometric component. Similarly, we evaluate the residuals for both steric configurations. When using the CIGAR manometric data, the mean residual trends amount to 0.88 ± 0.20 mm/yr for the hybrid steric case (SIO 0–2000 m plus CIGAR deep ocean) and 0.24 ± 0.22 mm/yr for the CIGAR full-depth case.

North Atlantic mean sea level budget



695

North Atlantic mean sea level budget



696

Figure 6+3. Time series of North Atlantic sea level change components (top) and budget residuals (bottom). The left panels use a hybrid steric estimate (SIO for 0–2000m combined with CIGAR for 2000–6000m), while the right panels use the full-depth CIGAR steric estimate (full-depth 0–6000m). In both left and right panels, CIGAR manometric components are considered. All time series have applied three-month moving average filtering. The global mean trend of each component is removed. Shaded areas represent data uncertainties.

~~Comparing the results from Figure 12 and Figure 13, we note that using CIGAR for the manometric component consistently leads to smaller regional residuals than using GRACE, regardless of the steric product considered. It is worth noting that the configuration using CIGAR for both manometric and steric components yields a mean residual trend of 0.24 ± 0.22 mm/yr, nearly four times smaller than in the case where the GRACE manometric component is used (0.90 ± 0.23 mm/yr). These results confirm our previous conclusions based on the residual trend maps. The trends of the residual time series for the different products of the sea level budget (global mean trends removed) are identical to those presented in Table 2 (gridded averages) and are not reproduced here.~~

64 Discussion

In this study, we have revisited the sea level budget of the North Atlantic over the 2004-2022 period, using a variety of different data sets. The objective was to further investigate the results from Bouih et al. (2025) who reported non-closure of the sea level budget in the North Atlantic over the same time span, and to identify which components of the budget (or which missing component) are responsible.

The main results of our study can be summarized as follows:

~~(4)~~—(1) By comparing the GRACE and CIGAR manometric trends with the ‘Altimetry minus steric full depth sea level’, we note a better agreement (same positive sign) between the latter and the CIGAR-based manometric component in the northeastern Atlantic, while the GRACE manometric trends have an opposite (negative) sign in this region.

~~(2) By comparing the SIO-based steric sea level trends with “Altimetry minus manometric sea level” in two cases (GRACE and CIGAR based manometric component) over the North Atlantic, the use of CIGAR manometric sea level provides higher correlation than when using GRACE. This comparison together with the one presented in (1) indicates potential errors in the GRACE and GRACE-FO data, particularly in the eastern region of the North Atlantic basin where the manometric component is strongly negative. The cause is so far unknown but possibly related to the [GIA](#) correction applied to GRACE data, e.g., the geocenter or GIA corrections. These needs dedicated investigation as shown in the recent study by Hightower et al. (2026).~~

~~(3) As of 2012, a negative trend is observed in the SIO halosteric component which suggests a physically real increase in salinity of the North Atlantic. Again, this needs further investigation.~~

~~(2) (4)~~ Our study shows that the deep ocean (>2000 m of depth) contribution to the North Atlantic sea level budget is not negligible and needs to be taken into account to obtain better closure of the sea level budget in this region. Based on CIGAR, the additional deep ocean steric contribution ~~(above the global mean deep ocean contribution)~~ to the North Atlantic sea level budget allows to almost close regional sea level budget amounts to 0.62 ± 0.04 mm/yr.

~~(5)~~ Accounting for the deep ocean steric contribution reduces by a factor of ~~330~~ 330% the residuals of the North Atlantic sea level budget when using the GRACE manometric component, and by a factor of ~~750~~ 750% when using CIGAR for the manometric component (and CIGAR full depth for the steric component). In the latter case, the mean residual trend over the North Atlantic is <1mm/yr, i.e., of the same order of magnitude as the regional trend error for altimetry gridded data according to Prandi et al. (2021). We can thus conclude that the closure of the sea level budget holds within the data uncertainties in this case.

~~(6) Our results also indicate that the main contribution to the mean North Atlantic residual trend comes from errors on the regional components (whatever the products used for the components) rather than from errors in the global mean trends.~~

~~(7) The North Atlantic sea level budget is closed within data uncertainties when using the CIGAR reanalysis for both the manometric and steric components.~~

~~(8) Our study also shows that the sea level budget of all other oceans (North Atlantic excluded) is almost closed when accounting for the deep ocean steric contribution based on the CIGAR ocean reanalysis~~

To conclude, these new findings represent a step further towards better understanding of the present-day sea level budget in the North Atlantic region. These investigations also support the need to better observe the deep ocean especially in the North Atlantic Ocean which is one of the objectives of the One Argo Project (Thierry et al., 2025). Nevertheless, some issues still merit deeper investigation. We can quote the followings, among others: Why GRACE manometric data give less good results in the Northeast Atlantic than the CIGAR-based manometric sea level? Which inaccurate correction applied to GRACE is involved? Is it related to the GIA ~~or the geocenter~~ corrections applied to GRACE? ~~What causes the slight salinity increase observed in the North Atlantic (full depth) with SIO data?~~ Can we explain the physical process causing the reported CIGAR-based positive deep ocean steric sea level? What is the impact on the CIGAR steric component estimates of the use of EN4 (with no salinity drift correction applied) in the assimilation procedure? What is the source of the (small) remaining residual trend of the North Atlantic sea level budget (after accounting for the deep ocean)? Clearly, future studies should be devoted to try answering these puzzling questions.

776 **Author contributions**

777 AC, WL and ZS designed the study. All analyses have been performed by ZS. ZS and AC
778 wrote a first version of the manuscript. All co-authors contributed to the discussion of the
779 results, editing and final writing of the manuscript.

780

781 **Competing interests**

782 The contact author has declared that none of the authors has any competing interests.

783

784 **Acknowledgements**

785 The authors thank two anonymous reviewers for helpful comments that significantly improved
786 the manuscript. Zhe Song is supported by China Scholarship Council. This study is a
787 contribution to the ongoing ESA (European Space Agency) CCI (Climate Change Initiative)
788 project entitled ‘Sea level budget closure CCI+ (SLBC_CCI+)’. This work is a contribution to
789 the GREAT project funded by CNES through the Ocean Surface Topography Science Team
790 (OSTST).

791

792 **References**

- 793 Ablain, M., Jugier, R., Zawadzki, L., Taburet, N., Cazenave, A., Meyssignac, B., and Picot,
794 N.: The TOPEX-A drift and impacts on GMSL time series, 2017.
- 795 Adhikari, S., Ivins, E. R., Frederikse, T., Landerer, F. W., and Caron, L.: Sea-level fingerprints
796 emergent from GRACE mission data, *Earth System Science Data*, 11, 629–646,
797 <https://doi.org/10.5194/essd-11-629-2019>, 2019.
- 798 Barnoud, A., Pfeffer, J., Guérou, A., Frery, M., Siméon, M., Cazenave, A., Chen, J., Llovel,
799 W., Thierry, V., Legeais, J., and Ablain, M.: Contributions of Altimetry and Argo to Non-
800 Closure of the Global Mean Sea Level Budget Since 2016, *Geophysical Research Letters*, 48,
801 <https://doi.org/10.1029/2021gl092824>, 2021.
- 802 Bouih, M., Barnoud, A., Yang, C., Storto, A., Blazquez, A., Llovel, W., Fraudeau, R., and
803 Cazenave, A.: Regional sea level trend budget over 2004–2022, *Ocean Sci.*, 21, 1425–1440,
804 <https://doi.org/10.5194/os-21-1425-2025>, 2025.
- 805 Brown, S., Willis, J., and Fournier, S.: Jason-3 Wet Path Delay Correction,
806 <https://doi.org/10.5067/J3L2G-PDCOR>, 2023.
- 807 Camargo, C. M. L., Riva, R. E. M., Hermans, T. H. J., Schütt, E. M., Marcos, M., Hernandez-
808 Carrasco, I., and Slangen, A. B. A.: Regionalizing the sea-level budget with machine learning
809 techniques, *Ocean Science*, 19, 17–41, <https://doi.org/10.5194/os-19-17-2023>, 2023.
- 810 Chen, J., Tapley, B., Seo, K., Wilson, C., and Ries, J.: Improved Quantification of Global Mean
811 Ocean Mass Change Using GRACE Satellite Gravimetry Measurements, *Geophys. Res. Lett.*,
812 46, 13984–13991, <https://doi.org/10.1029/2019GL085519>, 2019.
- 813 Chen, J., Tapley, B., Wilson, C., Cazenave, A., Seo, K.-W., and Kim, J.-S.: Global Ocean Mass
814 Change From GRACE and GRACE Follow-On and Altimeter and Argo Measurements,
815 *Geophysical Research Letters*, 47, e2020GL090656, <https://doi.org/10.1029/2020GL090656>,
816 2020.

- 817 Cheng, L., Zhu, J., Cowley, R., Boyer, T., and Wijffels, S.: Time, Probe Type, and Temperature
 818 Variable Bias Corrections to Historical Expendable Bathythermograph Observations, *Journal*
 819 *of Atmospheric and Oceanic Technology*, 31, 1793–1825, [https://doi.org/10.1175/JTECH-D-](https://doi.org/10.1175/JTECH-D-13-00197.1)
 820 13-00197.1, 2014.
- 821 Dieng, H. B., Cazenave, A., Meyssignac, B., and Ablain, M.: New estimate of the current rate
 822 of sea level rise from a sea level budget approach, *Geophysical Research Letters*, 44, 3744–
 823 3751, <https://doi.org/10.1002/2017GL073308>, 2017.
- 824 Dobslaw, H., Bergmann-Wolf, I., Dill, R., Poropat, L., Thomas, M., Dahle, C., Esselborn, S.,
 825 König, R., and Flechtner, F.: A new high-resolution model of non-tidal atmosphere and ocean
 826 mass variability for de-aliasing of satellite gravity observations: AOD1B RL06, *Geophysical*
 827 *Journal International*, 211, 263–269, <https://doi.org/10.1093/gji/ggx302>, 2017.
- 828 Flechtner, F., Dobslaw, H., and Fagiolini, E.: GRACE AOD1B product description document
 829 for product release 05, 2014.
- 830 Frederikse, T., Riva, R., Kleinherenbrink, M., Wada, Y., van den Broeke, M., and Marzeion,
 831 B.: Closing the sea level budget on a regional scale: Trends and variability on the Northwestern
 832 European continental shelf, *Geophysical Research Letters*, 43, 10,864–10,872,
 833 <https://doi.org/10.1002/2016GL070750>, 2016.
- 834 Gregory, J. M., Griffies, S. M., Hughes, C. W., Lowe, J. A., Church, J. A., Fukimori, I., Gomez,
 835 N., Kopp, R. E., Landerer, F., Cozannet, G. L., Ponte, R. M., Stammer, D., Tamisiea, M. E.,
 836 and van de Wal, R. S. W.: Concepts and Terminology for Sea Level: Mean, Variability and
 837 Change, *Both Local and Global*, *Surv Geophys*, 40, 1251–1289,
 838 <https://doi.org/10.1007/s10712-019-09525-z>, 2019.
- 839 Hamlington, B. D., Gardner, A. S., Ivins, E., Lenaerts, J. T. M., Reager, J. T., Trossman, D. S.,
 840 Zaron, E. D., Adhikari, S., Arendt, A., Aschwanden, A., Beckley, B. D., Bekaert, D. P. S.,
 841 Blewitt, G., Caron, L., Chambers, D. P., Chandanpurkar, H. A., Christianson, K., Csatho, B.,
 842 Cullather, R. I., DeConto, R. M., Fasullo, J. T., Frederikse, T., Freymueller, J. T., Gilford, D.

843 M., Giroto, M., Hammond, W. C., Hock, R., Holschuh, N., Kopp, R. E., Landerer, F., Larour,
844 E., Menemenlis, D., Merrifield, M., Mitrovica, J. X., Nerem, R. S., Nias, I. J., Nieves, V.,
845 Nowicki, S., Pangaluru, K., Piecuch, C. G., Ray, R. D., Rounce, D. R., Schlegel, N.-J.,
846 Seroussi, H., Shirzaei, M., Sweet, W. V., Velicogna, I., Vinogradova, N., Wahl, T., Wiese, D.
847 N., and Willis, M. J.: Understanding of Contemporary Regional Sea-Level Change and the
848 Implications for the Future, *Reviews of Geophysics*, 58, e2019RG000672,
849 <https://doi.org/10.1029/2019RG000672>, 2020.

850 Horwath, M., Gutknecht, B. D., Cazenave, A., Palanisamy, H. K., Marti, F., Marzeion, B.,
851 Paul, F., Le Bris, R., Hogg, A. E., Ootosaka, I., Shepherd, A., Döll, P., Cáceres, D., Müller
852 Schmied, H., Johannessen, J. A., Nilsen, J. E. Ø., Raj, R. P., Forsberg, R., Sandberg Sørensen,
853 L., Barletta, V. R., Simonsen, S. B., Knudsen, P., Andersen, O. B., Rannald, H., Rose, S. K.,
854 Merchant, C. J., Macintosh, C. R., von Schuckmann, K., Novotny, K., Groh, A., Restano, M.,
855 and Benveniste, J.: Global sea-level budget and ocean-mass budget, with a focus on advanced
856 data products and uncertainty characterisation, *Earth System Science Data*, 14, 411–447,
857 <https://doi.org/10.5194/essd-14-411-2022>, 2022.

858 Liu, C., Liang, X., Chambers, D., and Ponte, R.: Global Patterns of Spatial and Temporal
859 Variability in Salinity from Multiple Gridded Argo Products, *Journal of Climate*,
860 <https://doi.org/10.1175/JCLI-D-20-0053.1>, 2020.

861 Llovel, W., Balem, K., Tajouri, S., and Hochet, A.: Cause of Substantial Global Mean Sea
862 Level Rise Over 2014–2016, *Geophysical Research Letters*, 50, e2023GL104709,
863 <https://doi.org/10.1029/2023GL104709>, 2023.

864 Loomis, B. D., Rachlin, K. E., and Luthcke, S. B.: Improved Earth Oblateness Rate Reveals
865 Increased Ice Sheet Losses and Mass-Driven Sea Level Rise, *Geophys. Res. Lett.*, 46, 6910–
866 6917, <https://doi.org/10.1029/2019GL082929>, 2019.

867 McDougall, T. J., Barker, P. M., Marine, CSIRO., and Research, A.: Getting started with
868 TEOS-10 and the gibbs seawater (GSW) oceanographic toolbox, Trevor J. McDougall, 2011.

869 Mu, D., Church, J. A., King, M., Ludwigsen, C. B., and Xu, T.: Contrasting Discrepancy in the
 870 Sea Level Budget Between the North and South Atlantic Ocean Since 2016, *Earth and Space*
 871 *Science*, 11, e2023EA003133, <https://doi.org/10.1029/2023EA003133>, 2024.
 872 Nerem, R. S., Beckley, B. D., Fasullo, J. T., Hamlington, B. D., Masters, D., and Mitchum, G.
 873 T.: Climate-change–driven accelerated sea-level rise detected in the altimeter era, *Proceedings*
 874 *of the National Academy of Sciences*, 115, 2022–2025,
 875 <https://doi.org/10.1073/pnas.1717312115>, 2018.
 876 Pardaens, A. K., Gregory, J. M., and Lowe, J. A.: A model study of factors influencing
 877 projected changes in regional sea level over the twenty-first century, *Clim Dyn*, 36, 2015–2033,
 878 <https://doi.org/10.1007/s00382-009-0738-x>, 2011.
 879 Peltier, W. R., Argus, D. F., and Drummond, R.: Comment on “An Assessment of the ICE-
 880 6G_C (VM5a) Glacial Isostatic Adjustment Model” by Purcell et al.: The ICE-6G_C (VM5a)
 881 GIA model, *J. Geophys. Res. Solid Earth*, 123, 2019–2028,
 882 <https://doi.org/10.1002/2016JB013844>, 2018.
 883 Ponte, R. M., Sun, Q., Liu, C., and Liang, X.: How Salty Is the Global Ocean: Weighing It All
 884 or Tasting It a Sip at a Time?, *Geophysical Research Letters*, 48, e2021GL092935,
 885 <https://doi.org/10.1029/2021GL092935>, 2021.
 886 Roemmich, D. and Gilson, J.: The 2004–2008 mean and annual cycle of temperature, salinity,
 887 and steric height in the global ocean from the Argo Program, *Progress in Oceanography*, 82,
 888 81–100, <https://doi.org/10.1016/j.pocean.2009.03.004>, 2009.
 889 Royston, S., Dutt Vishwakarma, B., Westaway, R., Rougier, J., Sha, Z., and Bamber, J.: Can
 890 We Resolve the Basin-Scale Sea Level Trend Budget From GRACE Ocean Mass?, *Journal of*
 891 *Geophysical Research: Oceans*, 125, e2019JC015535, <https://doi.org/10.1029/2019JC015535>,
 892 2020.

893 Save, H., Bettadpur, S., and Tapley, B. D.: High-resolution CSR GRACE RL05 mascons:
 894 HIGH-RESOLUTION CSR GRACE RL05 MASCONS, *J. Geophys. Res. Solid Earth*, 121,
 895 7547–7569, <https://doi.org/10.1002/2016JB013007>, 2016.

896 Storto, A. and Yang, C.: Acceleration of the ocean warming from 1961 to 2022 unveiled by
 897 large-ensemble reanalyses, *Nat Commun*, 15, 545, [https://doi.org/10.1038/s41467-024-44749-](https://doi.org/10.1038/s41467-024-44749-7)
 898 7, 2024.

899 Sun, Y., Riva, R., and Ditmar, P.: Optimizing estimates of annual variations and trends in
 900 geocenter motion and J_2 from a combination of GRACE data and geophysical models, *J.*
 901 *Geophys. Res. Solid Earth*, 121, 8352–8370, <https://doi.org/10.1002/2016JB013073>, 2016.

902 Tapley, B., Watkins, M., Flechtner, F., Reigber, C., Bettadpur, S., Rodell, M., Sasgen, I.,
 903 Famiglietti, J., Landerer, F., Chambers, D., Reager, J., Gardner, A., Save, H., Ivins, E.,
 904 Swenson, S., Boening, C., Dahle, C., Wiese, D., Dobs law, H., and Velicogna, I.: Contributions
 905 of GRACE to understanding climate change, *Nature Climate Change*, 5,
 906 <https://doi.org/10.1038/s41558-019-0456-2>, 2019.

907 Thierry, V., Claustre, H., Pasqueron De Fommervault, O., Zilberman, N., Johnson, K. S., King,
 908 B. A., Wijffels, S. E., Bhaskar, U. T. V. S., Balmaseda, M. A., Belbeoch, M., Bollard, M.,
 909 Boutin, J., Boyd, P., Cancouët, R., Chai, F., Ciavatta, S., Crane, R., Cravatte, S., Dall’Olmo,
 910 G., Desbruyères, D., Durack, P. J., Fassbender, A. J., Fennel, K., Fujii, Y., Gasparin, F.,
 911 González-Santana, A., Gourcuff, C., Gray, A., Hewitt, H. T., Jayne, S. R., Johnson, G. C.,
 912 Kolodziejczyk, N., Le Boyer, A., Le Traon, P.-Y., Llovel, W., Lozier, M. S., Lyman, J. M.,
 913 McDonagh, E. L., Martin, A. P., Meyssignac, B., Mogensen, K. S., Morris, T., Oke, P. R.,
 914 Smith, W. O., Owens, B., Poffa, N., Post, J., Roemmich, D., Rykaczewski, R. R.,
 915 Sathyendranath, S., Scanderbeg, M., Scheurle, C., Schofield, O., Von Schuckmann, K.,
 916 Scourse, J., Sprintall, J., Suga, T., Tonani, M., Van Wijk, E., Xing, X., and Zuo, H.: Advancing
 917 ocean monitoring and knowledge for societal benefit: the urgency to expand Argo to OneArgo
 918 by 2030, *Front. Mar. Sci.*, 12, 1593904, <https://doi.org/10.3389/fmars.2025.1593904>, 2025.

919 Wang, C., Dong, S., and Munoz, E.: Seawater density variations in the North Atlantic and the
 920 Atlantic meridional overturning circulation, *Clim Dyn*, 34, 953–968,
 921 <https://doi.org/10.1007/s00382-009-0560-5>, 2010.
 922 Watkins, M. M., Wiese, D. N., Yuan, D.-N., Boening, C., and Landerer, F. W.: Improved
 923 methods for observing Earth’s time variable mass distribution with GRACE using spherical
 924 cap mascons, *Journal of Geophysical Research: Solid Earth*, 120, 2648–2671,
 925 <https://doi.org/10.1002/2014JB011547>, 2015.
 926 WCRP: Global sea-level budget 1993–present, *Earth System Science Data*, 10, 1551–1590,
 927 <https://doi.org/10.5194/essd-10-1551-2018>, 2018.
 928 Wong, A. P. S., Gilson, J., and Cabanes, C.: Argo salinity: bias and uncertainty evaluation,
 929 *Earth System Science Data*, 15, 383–393, <https://doi.org/10.5194/essd-15-383-2023>, 2023.
 930 Wunsch, C., Ponte, R. M., and Heimbach, P.: Decadal Trends in Sea Level Patterns: 1993–
 931 2004, *Journal of Climate*, 20, 5889–5911, <https://doi.org/10.1175/2007JCLI1840.1>, 2007.
 932
 933
 934