



The Mineral Aerosol Profiling from Infrared Radiances version 5.1 algorithm and its evaluation

Sophie Vandenbussche¹, Christodoulos Biskas², Maria-Elissavet Koukouli², Stelios Kazadzis³, and Martine De Mazière¹

¹Royal Belgian Institute for Space Aeronomy, Brussels, Belgium

²Laboratory of Atmospheric Physics, Aristotle University of Thessaloniki, Thessaloniki, Greece

³Physikalisch-Meteorologisches Observatorium Davos, World Radiation Centre, Davos, Switzerland

Correspondence: Sophie Vandenbussche (sophie.vandenbussche@aeronomie.be)

Abstract. Mineral (desert) dust aerosols are small sand/dust particles entrained by winds from bare areas and possibly transported over long distances. These aerosols are climate forcers and affect human health and many socio-economic sectors. They are therefore important to monitor both in near-real time and on the long term. In this work, the Infrared Atmospheric Sounding Interferometer (IASI) instrument is used to retrieve vertical profiles of mineral dust aerosols concentration, from which a 10 μm aerosol optical depth (AOD) and a mean aerosol altitude are obtained. More specifically, we present here the new version 5.1 of the Mineral Aerosol Profiling from Infrared Radiances (MAPIR) algorithm and its changes with respect to previous versions. MAPIR v5.1 was used to produce a consistent time series of dust profiles since the start of the IASI observations in 2007 and until now, using data from IASI onboard Metop-A and Metop-C. The capabilities of the instrument and retrieval are illustrated, showing good event detection, expected AOD seasonal cycles, good profiling capabilities and reasonable mean aerosol altitude, good time and cross-platform consistency. A true validation exercise is not possible as there exist no reference aerosols data from thermal infrared measurements (around 10 μm). Therefore, the absolute value of the obtained AOD can not be validated, although the best possible evaluation is provided using data obtained in the visible spectral range.

1 Introduction

Mineral dust, commonly referred to as desert dust, constitutes the most abundant atmospheric aerosol in dry mass (e.g. Textor et al., 2006), consisting of particles from arid and semi-arid regions. These particles are lofted into the atmosphere by strong winds and can travel vast distances across continents and oceans. Their size ranges from less than 1 nm to more than 100 μm (e.g. Mahowald et al., 2014). The main sources of mineral dust include the Sahara Desert in Africa, the Gobi Desert in Asia, and other dry regions worldwide. The global emission of mineral dust is estimated to be between 1 and 3 Gt yr^{-1} , and its average atmospheric dust load ranges from 8 to 35 Mt yr^{-1} (e.g. Zender et al., 2004).

The impact of mineral dust on Earth's climate is multifaceted. First, it influences the radiative balance by (back)scattering and absorbing sunlight in the short-wave spectral range, while absorbing, scattering, and emitting radiation in the infrared spectral range. This is called the direct radiative effect. Those interactions with light depend on different aerosol properties in different spectral ranges (e.g. Adebisi et al., 2023; Di Biagio et al., 2017, 2019; Koepke et al., 2015). They result in atmospheric



and/or surface heating or cooling, depending on the dust layer height, the atmospheric and surface temperatures, the surface
25 albedo and emissivity, the presence of clouds or other aerosols, the presence of sunlight, etc. Finally, these changes in surface
and atmospheric temperatures also affect surface energy exchanges and local wind speed. (e.g. Choobari et al., 2014). Mineral
dust can also modify cloud cover, cloud properties and precipitation processes by acting as cloud condensation and ice nuclei
(CCN and IN) (e.g. Choobari et al., 2014). These are called the semi-direct (cloud impact due to atmospheric heating) and
indirect radiative effects (cloud impact due to microphysics). The deposition of dust can stimulate biogeochemical activity in
30 ecosystems (both marine and terrestrial) by providing essential nutrients such as iron and phosphorus (e.g. Jickells et al., 2005).
This deposition may also locally modify the visible surface albedo resulting in increased snow melt (Xie et al., 2018).

Beyond its climatic effects, mineral dust has significant implications for human health, affecting the respiratory and circula-
tory systems (e.g. Zhang et al., 2016). Finally, dust aerosols also affect human activities in many ways (e.g. Middleton, 2017;
Monteiro et al., 2022; Mona et al., 2023): they disrupt transportation, damage infrastructure, reduce agricultural productivity,
35 reduce photovoltaic energy production, etc.

To monitor the dust impacts on the climate and analyse changes, one needs long-term consistent data sets to be able to extract
trends in different variables such as the dust load, vertical distribution, transport paths, source areas. To develop mitigation
strategies related to the dust impact on human health and activities, and to correctly forecast temperatures, clouds and rain,
near-real-time data of dust atmospheric distribution and properties are required. Those may be used as input in aerosol forecast
40 or weather models. For both types of analyses, data should be global (or almost), possibly cover day and night and, in the case
of forecast models, data should have the best feasible temporal resolution. This can be reached using satellite remote sensing.
Different types of instruments deliver different information on the dust aerosols, with different time resolutions and ground
coverage. A comprehensive recent review on the dust observation from satellites (and also all other observations) can be found
in Mona et al. (2023). Specificities of thermal infrared (TIR) observations versus observations in the visible spectral range
45 will be discussed in detail in Sect. 3.2, as they imply additional challenges for the validation of satellite observations. Those
specificities are also relevant for any data user, to correctly understand the typical properties of dust data retrieved from the
TIR spectral range, compared to the better known UV/Vis range.

The algorithm presented here falls within the effort to improve the current dust observation capabilities from TIR satellite
observations by the Infrared Atmospheric Sounding Interferometer (IASI) instrument. This algorithm is called Mineral Aerosol
50 Profiling from Infrared Radiances (MAPIR), and its previous version was described in detail in Callewaert et al. (2019). MAPIR
uses IASI radiance data, with additional information on the surface and atmospheric state, to retrieve vertical profiles of coarse
mineral dust aerosol concentrations. Those profiles may then be used to calculate the integrated Aerosol Optical Depth (AOD)
at a TIR wavelength (which is usually 10 μm) and converted to the visible wavelength most commonly used to report AOD,
550 nm. The profiles can also be used to calculate a mean altitude of the aerosol layer.

This manuscript describes the new version 5.1 of the algorithm, with several modifications such as the removal of all pre-
55 filters (which were responsible for missing some events), the increased vertical range to 10 km (to better cover rare special
high altitude events, with possible impact on aviation, or events above the Tibetan Plateau), the use of a different emissivity
database (while using it, it was made clear that the previously used one was biased low in areas/months of almost continuous



atmospheric dust presence). Furthermore, the uncertainty in the input atmospheric profiles of temperature and humidity is now propagated to the retrieval uncertainty, as such improving its estimation. Finally, the algorithm exists in two slightly different versions, depending on the need to produce long-term consistent data (climate version) or the best possible data in near-real-time (NRT version). MAPIR-climate uses as input the reprocessed IASI spectra and atmospheric data, and the corresponding variables available from the recently acquired IASI data (see section 2.1.1). This allows full consistency in that input data from the launch of the first IASI instrument in 2007. MAPIR-NRT uses the best possible data available in NRT, which is then evolving with time and changes in the EUMETSAT operational IASI processing chain. This work presents and evaluates the data produced with the MAPIR-climate version, while the MAPIR-NRT version will be assessed later on, when enough data will have been produced in NRT mode.

The manuscript is organised as follows. First, the MAPIR retrievals are fully described (Sect. 2), starting with the data it requires, continuing with the detailed algorithm description and changes with respect to the previous version, and ending with a general assessment of its performances. Section 3 demonstrates the capabilities of MAPIR and shows comparisons with data from other instruments. The section begins with the description of the data sets used in this evaluation, continues with a detailed discussion of the challenges of evaluating TIR-observed aerosol information, then shows the evaluation of the integrated dust AOD against reference measurements by the Aerosol Robotic Network (AERONET), discusses some properties of the profile data and shows example comparisons with Cloud–Aerosol lidar with Orthogonal Polarization (CALIOP) observations and against European Aerosol Research Lidar Network (EARLINET) data. The section ends with a discussion of MAPIR uncertainties, long-term stability and IASI-A versus C continuity. Finally, we wrap the paper up and propose some conclusions and future perspectives.

2 MAPIR retrievals

2.1 Instruments, models and data used in the MAPIR retrievals

2.1.1 IASI

The Infrared Atmospheric Sounding Interferometer (IASI) is a Michelson Interferometer developed by the Centre National d'Études Spatiales (CNES) in collaboration with the European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT). Mounted on the MetOp series of polar-orbiting satellites, IASI is designed to measure atmospheric temperature and humidity profiles with high accuracy and resolution, for use in weather forecast models. Its data has in addition been used to retrieve numerous other atmospheric gases such as carbon dioxide, methane, nitrous oxide, ozone, carbon monoxide and many other traces gases (e.g. Clerbaux et al., 2009; Clarisse et al., 2011; Hilton et al., 2012; Vandenbussche et al., 2022; De Wachter et al., 2017). Finally, IASI is also used to retrieve mineral aerosol (desert dust and volcanic ash) atmospheric load, mean altitude or vertical profile, particle composition and size (e.g. Callewaert et al., 2019; Clarisse et al., 2019; Capelle et al., 2018; Klüser et al., 2015; Maes et al., 2016).



90 The IASI instruments are deployed on the Metop series of satellites, which operate in a polar, sun-synchronous orbit at an altitude of approximately 817 km. The Equator crossing times are 09:30 (descending node) and 21:30 (ascending node) in mean local solar time, providing near-global coverage twice per day for each instrument. IASI is a nadir-viewing sensor with a swath width of about 2200 km and can observe up to 48.3° off-nadir on either side of the ground track. Each scan comprises 30 elementary fields of view, and each field contains four circular pixels with a diameter of 12 km at nadir. Toward the edges
95 of the swath, the footprint deforms into an ellipse of approximately 39×20 km. The instrument records radiances over the $645\text{--}2760\text{ cm}^{-1}$ spectral interval, with an apodized spectral resolution of 0.5 cm^{-1} . In the thermal infrared atmospheric window (used in MAPIR retrievals), the radiometric noise is on the order of 0.2 K (Clerbaux et al., 2009). Each spectrum is sampled every 0.25 cm^{-1} , yielding a total of 8461 radiance channels.

IASI data is perfectly suited for both long-term and NRT monitoring of the atmosphere, with continuous data since 2007
100 through three identical very stable IASI instruments launched in 2006, 2012 and 2018, one new generation instrument (IASI-NG) launched in summer 2025 and two others planned for further continuation of the observations until at least 2040 (CNES, 2026).

In this work, we use data from IASI/Metop-A (in short, IASI-A, data from July 2007 to September 2021) and IASI/Metop-C (in short, IASI-C, data since October 2019, including 2 years overlap with IASI-A). For computing time mitigation, we do
105 not use data from IASI/Metop-B which overlaps with the other instruments. For IASI-A we use level 1c radiance from the IASI Principal Components Scores (PCS) Fundamental Data Record Release 1 (EUMETSAT, 2022a), and surface temperature, atmospheric temperature and humidity profiles, land flag and surface elevation from the IASI All Sky Temperature and Humidity Profiles - Climate Data Record Release 1.1 (EUMETSAT, 2022b). Those will further be referred to as CDR (climate data record) for shortness. For IASI-C, we use the similar level 1c PCS from the NRT data (EUMETSAT, 2025a) and surface
110 temperature, atmospheric temperature and humidity profiles, land flag and surface elevation from the NRT Combined Sounding product, both in the "native" file format, delivered with a small delay through the EUMETSAT Data Store (EUMETSAT, 2025b). Those will further be referred to as OFL ("offline") for shortness. From that data, we select surface temperature and atmospheric temperature and humidity obtained with the Piece Wise Linear Regression version 3 (PWLR3) algorithm (and not the Optimal Estimation Method or OEM). This ensures consistency with IASI-A, as the PWLR3 algorithm was used for
115 reprocessing the IASI-A CDR. Table 1 summarizes all this information.

An update was implemented on 30 March 2023 by EUMETSAT. The construction of the PCS (from level 1c radiance data) now relies on a revised eigenvector basis. In parallel, the noise normalisation matrix has been enhanced: instead of a purely diagonal formulation, a full matrix is now applied. This modification enables noise decorrelation, thereby improving filtering performance compared with the previous approach, which was limited to variance homogenisation. This version of
120 the PCS also introduces a hybrid configuration that incorporates supplementary local PCS and associated eigenvectors. These additions are intended to better represent atypical local spectral features that may not be adequately captured by the global PCS framework. In the present work, these local PCS components are not employed, as the dust spectral signature is already satisfactorily described by the global eigenvector basis. Their exclusion avoids unnecessary complexity in processing. This



Table 1. IASI level 1 and level 2 input data for MAPIR v5.1 retrievals. FDR means Fundamental Data Record. CDR means climate data record. NRT means near-real-time

	IASI-A	IASI-C
Radiance	IASI PCS FDR release 1 (EUMETSAT, 2022a)	NRT IASI level 1 PCS (EUMETSAT, 2025a)
Atmosphere and surface	IASI all-sky T and H ₂ O CDR release 1.1 -> result of the PWLR3 algo (EUMETSAT, 2022b)	IASI level 2 NRT Combined Sounding Product -> selecting the PWLR3 data (EUMETSAT, 2025b)
Used abbreviation	RPRO (reprocessed)	OFL (offline)

update on 30 March 2023 represents the only discontinuity in the complete processing chain for the new MAPIR version 5.1,
 125 and bears no detectable impact (see Sect. 3.8).

The set-up described above refers to the MAPIR-climate setup, which is analysed and validated in this work. The MAPIR-NRT setup is based on exactly the same algorithm, uses the same level 1C data input but uses level 2 data surface temperature and atmospheric profiles from the IASI Optimal Estimation algorithm when available (and the same PWLR3 data when that OEM did not perform well). The MAPIR-NRT data are therefore slightly different from the MAPIR-climate data, and will
 130 contain inconsistencies when the IASI OEM is modified. The MAPIR-NRT data is planned for NRT distribution through the EUMETCast system, within work from the EUMETSAT Atmospheric Composition Satellite Application Facility (AC SAF), and will be validated in that context when enough data will have been generated.

2.1.2 Dust optical properties

The dust aerosol properties are fixed throughout time and space in MAPIR retrievals. The retrieval scheme assumes spherical
 135 particles characterised by the GEISA–HITRAN dust-like refractive index (RI) (Jacquinet-Husson et al., 2011). The particle size distribution is represented by a single-mode log-normal function with a mean radius of 0.6 μm and a geometric standard deviation of 2, yielding an effective radius of approximately 2 μm . Those properties are used in a Mie code to calculate the particle properties required for the radiative transfer code (see Sect. 2.1.4) and for conversion between visible and TIR wavelengths (see Sect. 2.1.3 and 3.2). The impact of the spherical assumption is also discussed in Sect. 3.2.

140 2.1.3 LIVAS

The LIVAS (Lidar climatology of Vertical Aerosol Structure for space-based lidar simulation studies) database utilizes data from the CALIOP (Cloud-Aerosol Lidar with Orthogonal Polarization) instrument aboard the CALIPSO (Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations) satellite, to provide a monthly average of high-resolution vertical profiles of dust extinction at visible wavelengths, globally on a $1^\circ \times 1^\circ$ horizontal grid, from 2007 to 2014 (Amiridis et al., 2013, 2015).



145 This data is used as *a priori* for our dust profile retrievals, after some pre-treatment. First, the high resolution extinction profiles are regridded to the low resolution MAPIR retrieval vertical grid. Second, the monthly data are averaged over the 8 years to obtain a mean monthly data set. Then, to assure having data in each grid cell and continuity between adjacent cells, a running mean of the data set is calculated along $5^\circ \times 5^\circ$. Finally, the extinction at 532 nm is converted to concentration (particles cm^{-3}) using a Mie code and the same dust particle optical properties used in the retrieval (see 2.1.2) and the obtained
150 concentrations in each layer are clipped to the minimum value of 2 particle cm^{-3} and a maximum value of 100 particles cm^{-3} . This minimal concentration is new in MAPIR version 5.1 and its justification is provided in Sect. 2.2.4.

2.1.4 RTTOV

RTTOV (Radiative Transfer for TOVS) is a fast radiative transfer model developed by the European Centre for Medium-Range Weather Forecasts (ECMWF) and the Satellite Application Facility on Numerical Weather Prediction (NWP SAF) (Saunders
155 et al., 2018). It is designed to simulate satellite radiances for a variety of atmospheric, surface, and cloud conditions in the infrared, visible, and microwave spectral regions. RTTOV uses pre-computed transmittance coefficients to efficiently calculate radiative transfer. The model also allows computing derivatives of the radiance with respect to some atmospheric gases. Finally, the model allows including aerosols with user-defined optical properties, and computing derivatives of the radiance with respect to the aerosol concentration.

160 In this work, RTTOV is used for all radiative transfer computations. We use RTTOV version 13 with the corresponding IASI version 13 predictor coefficients calculated on 101 levels (Saunders et al., 2020), with only water vapour as variable gas (mandatory), for which the vertical profiles originate from the IASI level 2 data described in Sect. 2.1.1 and interpolated on the internal RTTOV grid. All other gases are modelled with fixed concentrations using the RTTOV defaults values. The specific aerosol properties used in this work are described in Sect. 2.1.2. No solar sources are included (irrelevant in the TIR).
165 Scattering is computed using the Discrete Ordinate Method (DOM) with 4 streams. The plane parallel approximation is used, as it is mandatory for using the DOM. Surface is modelled as lambertian. Vertical profile interpolation, if needed, is done using the RTTOV default method. All input data are tested before running RTTOV. When needed, profiles are extrapolated up to top of atmosphere by maintaining the ratio to the default RTTOV profile instead of repeating the last known value. Concentrations are clipped to limits for the coefficient regression. All other RTTOV options are left to their default value.

170 2.1.5 Surface emissivity

In this study, surface emissivity is taken from the internal databases provided in RTTOV v13 (Saunders et al., 2020). Over oceanic regions, the IREMIS model is applied, with the surface wind speed fixed at 0 m s^{-1} due to the absence of wind information; this assumption has a negligible effect on dust retrievals. Over land, emissivity is prescribed from the Combined ASTER MODIS Emissivity over Land (CAMEL) climatology, version 2. The snow fraction is set to zero, as this parameter
175 is not available in the IASI observations. Within IREMIS, the automatic correction for satellite viewing angle is activated. RTTOV also offers an optional viewing angle correction for land surface emissivity; however, this adjustment originates from



a limited study and was found to generate spurious artefacts in the retrievals. Consequently, it is disabled in MAPIR. The classification between ocean and land surfaces relies on the dedicated flag provided in the IASI level 2 product.

2.2 The MAPIR algorithm

180 The MAPIR algorithm comprises pre-treatment / filtering of the input data (simple quality control), the retrieval itself, and post-treatment / quality control of the retrieved parameters. The retrieval algorithm is based on the optimal estimation method (OEM, (Rodgers, 2000)), which provides a maximum a posteriori solution by iteratively adjusting a state vector (the retrieved variables: here, the vertical profile of dust aerosol concentration and the surface temperature) in order to minimize a cost function. The latter contains both a measure of the difference between the observed (here, with the IASI instrument) and
185 simulated (with a radiative transfer model or RTM, here RTTOV v13) spectra, and a constraint on the possible values of the state vector (the "prior knowledge", usually a climatological mean and covariance of the retrieved parameters, here the LIVAS climatology with an assumed fixed 50% standard deviation). The need for an iterative method comes from the high non-linearity of the RTM with respect to the retrieved parameters. The OEM also allows estimating the information content of the retrieved data (the number of independent information pieces), and to propagate uncertainties from observation noise,
190 model and input variables uncertainties, to retrieved parameters uncertainties. The biggest (maybe only) drawback of the OEM is that it is computationally very demanding, due to its iterative nature and the subsequent need for multiple calls to the RTM.

The MAPIR version 5.1 algorithm is improved from MAPIR version 4.1 (Callewaert et al., 2019). The next subsections present the full description of the new version, and end with a description of the differences, in order to highlight them and explain their impact on the results. Fig. 1 shows a schematic representation of all data flow and algorithm parts, with the
195 reference to the section where each is discussed.

2.2.1 Pre-processing algorithm

The pre-processing of MAPIR v5.1 consists in reading the IASI radiances, surface temperature and atmospheric temperature and humidity profiles (described in Sect. 2.1.1), then screening them using the available quality indicators or RTTOV internal boundaries. The quality indicators are slightly different depending on the source of the data (CDR for IASI-A or OFL for
200 IASI-C). All criteria are listed in Table 2. All of them need to be met for a IASI observation to enter the retrieval. It is rare that observations are rejected due to these criteria (see further Sect. 2.3).

Surface temperature, surface pressure, and the atmospheric profiles (pressure, temperature, and water vapour) are required to lie within the validity ranges defined for RTTOV v13 (Saunders et al., 2020). Accordingly, surface temperature and the water vapour profile are internally constrained to the permissible intervals within RTTOV, as part of the input pre-processing.
205 In contrast, surface pressure as well as the atmospheric pressure and temperature profiles cannot be reliably adjusted. These parameters are not retrieved within MAPIR. Owing to their strong influence on the forward simulations and retrieval outcomes, any IASI observation for which one of these variables falls outside the RTTOV admissible limits is excluded from further processing.




Figure 1. Schematic representation of MAPIR v5.1 data flow and algorithm parts; the numbers in the grey circles refer to the section where each is discussed.

Table 2. IASI level 1 and level 2 quality control before use in MAPIR v5.1 retrievals. B1 and B2 refer to the respective spectral bands of the IASI instrument, amongst 3 in total. The MAPIR spectral windows are contained in B1 and B2.

CDR, IASI-A		OFL, IASI-C
Radiance, level 1c PCS	QFlag = 0 in B1 and B2	GQisQualIndex >= 95 GQisFlagQual = 0 in B1 and B2 ResidualRMS <= 1.05 in B1 and B2
T and H ₂ O profiles, level 2	qualityIndicatorForAtmosphericTemperature <= 2K qualityIndicatorForAtmosphericWaterVapour <= 5K	FG_QI_ATMOSPHERIC_TEMPERATURE <= 2K FG_QI_ATMOSPHERIC_WATER_VAPOUR <= 5K FLG_IASIBAD = 0
Product guides	(EUMETSAT, 2022a, b)	(EUMETSAT, 2019, 2017)

2.2.2 Retrieval scheme

210 The retrieval scheme is based on the optimal estimation method (OEM, Rodgers, 2000). The state vector, containing the retrieved parameters, is composed of the vertical profile of the logarithm of the dust aerosol concentration in 10 layers centred from 0.5 to 9.5 km altitude above sea level (by steps of 1 km) and of the surface temperature. Using the logarithmic space ensures that a negative concentration is never obtained, as such unphysical conditions would not be acceptable for RTTOV. The starting value of the state vector is the same as the *a priori*, which comes from the LIVAS climatology for the dust concentration

215 profile (see Sect. 2.1.3), and from the IASI PWLR3 level 2 data for the surface temperature (see Sect. 2.1.1).



MAPIR uses three spectral windows, on both sides of the well-known "V-shape" signature of minerals in the TIR (e.g. Sokolik, 2002). The first two spectral windows are highly sensitive to mineral aerosols, and also bear sensitivity to surface temperature. They range from 905 to 927 cm^{-1} and from 1098 to 1123 cm^{-1} . The third spectral window ranges from 1202.75 to 1204.75 cm^{-1} , and is mostly sensitive to the surface temperature, with still a small sensitivity to mineral aerosols. The spectral noise is set to 0.5 K in the first two windows and 0.2 K in the third window. This is a small noise inflation with respect to the instrument spectral noise reported to be 0.2 K in the first two windows, 0.03 K in the third by Clerbaux et al. (2009) or 0.25 K, 0.33 K and 0.07 K respectively by Hilton et al. (2012).

All radiative transfer computations are performed with RTTOV (see Sect. 2.1.4). The radiance derivatives with respect to the state parameters are calculated during each retrieval main loop, to account for their non-linearity. Those derivatives are used to compute the state vector for the next iteration, following the OEM formalism.

Due to the pronounced non-linearity of the radiative transfer with respect to the retrieved variables, the inversion relies on the Levenberg–Marquardt algorithm (Levenberg, 1944; Marquardt, 1963), which extends the Gauss–Newton iterative scheme. The damping parameter γ is initially set to 1. If the cost function increases during an iteration, γ is multiplied by 10 and the iteration is repeated with this updated value. Conversely, when the cost function decreases, γ is reduced by a factor of 2. Convergence is assessed within the standard Optimal Estimation Method (OEM) framework, applying a threshold of 0.1 in both state space and measurement space. In practice, convergence is typically achieved in fewer than six iterations, with an upper limit of 20 iterations imposed.

Finally, averaging kernels and uncertainties are provided, following the OEM formalism and the Levenberg–Marquardt modification, using equations from Ceccherini and Ridolfi (2010), as in the previous version of MAPIR (Callewaert et al., 2019). The result covariance matrix is computed separately for the uncertainties linked to the spectral noise, to the smoothing error, and to the input temperature and humidity profiles. The information content of the retrievals (degrees of freedom) is the trace of the averaging kernel matrix, and reports the number of independent pieces of information present in each single profile. In good cases (strong event and / or large temperature contrast between the surface and the dust layer), the retrievals may reach up to 2.5 degrees of freedom. Those degrees of freedom are distributed in the vertical range depending on the surface and atmospheric temperature, but also depend on the dust concentration profile because the retrieval is performed in the logarithmic space (e.g. Callewaert et al., 2019).

2.2.3 Post-processing algorithm: quality flag and additional variables

The post-processing consists in computing the dust AOD with associated uncertainty and the dust mean altitude, both from the full vertical profile. Uncertainty covariance matrices are computed, and finally quality and cloud flags are added.

The calculation of the dust AOD from the concentration profile C_{dust} is simply a vertical summation of the dust layers to a dust column, which is then multiplied by the cross-section obtained from the dust PSD and RI used for the retrieval (see Sect. 2.1.2). This can be formalised using the transformation operator L : a row of the same size as the vertical profile, containing a repetition of the cross-section value, multiplied by the layer height (which is currently constant in MAPIR). The AOD is then simply $C_{dust} * L$. The cross-section depends on the wavelength, and the dust AOD is reported both at $10\mu\text{m}$ (close to the



250 retrieval spectral windows, although $10\mu\text{m}$ is not in those windows) and at 550 nm (a standard wavelength for AOD reporting).
The ratio between the AOD at 550 nm and that at $10\mu\text{m}$ is 1.78. A discussion of that conversion takes place in Sect. 3.2.

The dust AOD uncertainty U_{AOD} is calculated from the OEM-obtained uncertainty covariance matrix S on the profile, using the transformation operator L : $U_{AOD} = \sqrt{L \times S \times L^T}$. In addition, an evaluation of the AOD uncertainty due to surface emissivity uncertainty is provided in Sect. 3.7, however this added uncertainty is very difficult to quantify for each retrieval
255 and is therefore not part of the pixel-wise reported uncertainty.

The mean dust altitude is defined as the height that partitions the dust column into two equal parts, with 50% of the particles located below and 50% above this level. This metric loses physical relevance in situations where several distinct dust layers coexist. Furthermore, it is only provided when the dust aerosol optical depth (AOD) exceeds 0.05 and the associated information content is greater than 1.25. Lower information content means that there is not enough information in the vertical range to
260 provide more than just an integrated vertical column, and the mean altitude usually ends up being the middle of the retrieval range, i.e. about 5 km.

The data used to build the cloud flag is again slightly different for the CDR IASI-A and the OFL IASI-C data. In the first case, only the cloud fraction is available (from the level 1 data) and the cloud-free criterion is that the fraction is less than 10%. In the second case, the same holds true and in addition, the level 2 cloud flag must be 1 or 2 (respectively clear sky and cloud
265 free with small contamination possible (EUMETSAT, 2017)).

The MAPIR data quality flag should be used to filter the retrieved data, before using them. The goal is to remove the spurious results, either due to non-convergence (reaching 20 iterations) or converging but with large spectral residuals (root-mean-square of the residuals larger than 0.75 K) or unreasonable values of the retrieved parameters (surface temperature below 200 K or above 350 K, dust AOD at $10\mu\text{m}$ larger than 5). Those underline a high probability that even though mathematically the
270 convergence is reached, something is wrong (which can be any of the input data, or the presence of a cloud), and the retrieval result does not reasonably reflect the reality. Finally, a test has been added to further remove thin clouds as they are not pre-filtered now. The dust aerosol retrieval converges on most cloudy observations, even though the spectral signature is different. If the cloud is thick, the aerosol retrieval will provide a very high concentration, leading to saturation of the signal in the TIR, together with a very low surface temperature. At that point, the dust signal would be similar to a cloud. This is easy to filter
275 out, as the retrieved parameters are unrealistic. If the cloud is thin, the retrieval provides a small dust concentration, together with a surface temperature slightly biased low. This is more complex to filter out. In the current setup, we exploit the fact that the EUMETSAT PWLR3 surface temperature (used as *a priori* in our retrievals, see Sect. 2.1.1) is an all-sky statistical product, with very decent quality even under a cloud or aerosols. The MAPIR-retrieved surface temperature will on the other hand be biased low in most cloud cases (as clouds reduce the observed radiance and are not modelled in MAPIR retrievals),
280 and should bear no bias in the dust aerosol cases (as those are correctly modelled in MAPIR retrievals). The criteria used for filtering thin clouds is therefore that the retrieved surface temperature can not be more than 2.5 K lower than the *a priori* surface temperature. Obviously, this filter is far from perfect, and it removes some thin dust observations, while retaining some thin cloud observations. However, it performs better in that discrimination than using the IASI cloud fraction (and flag), which were wrongly flagging intense dust plumes.



285 2.2.4 MAPIR v5.1 changes with respect to MAPIR v4.1

A first set of changes impacts the amount of data treated by the retrieval, through the removal of almost all prefilters. This also drives other changes to reduce single retrieval computational time and improve post-retrieval filtering. In MAPIR v5.1, we have removed the pre-filtering for the presence of dust outside the dust belt, which was previously performed by looking at brightness temperature differences to detect the dust V-shaped spectral signature. This filtering was shown to miss most of the
290 low intensity events, where the spectral slopes are low, as well as special conditions where the spectral V-shaped signature is inverted (when the dust layer is hotter than the surface). In addition, the cloud filtering prior to running the retrieval has been removed because the EUMETSAT IASI cloud flag is wrongly categorizing intense dust events as clouds. It was decided to run the retrieval on all IASI observations and remove the cloud-contaminated scenes as post-processing, as detailed in in Sect. 2.2.3. Finally, the retrieval is now running globally instead of only up to 60° latitude for version 4.1.

295 All these changes increase the number of retrievals to be run by an order of magnitude. To mitigate the impact on processing costs, ways to reduce the retrieval duration were investigated. The number of streams in the multiple scattering DOM was reduced from 8 to 4, gaining about a factor of 4 in computing time. This reduction also led to slight changes in the retrieved dust AOD, which is smaller by a maximum of 0.015 at 10 μm in the performed tests. An additional upgrade leading to a slight gain computation time is a technical improvement to the whole processing chain, in particular in the pre-processing part of
300 reading, colocating and quality filtering the IASI level 1 spectra and level 2 data. At the end, MAPIR v5.1 treats about 11 times more data, requiring only about 2.5 times more computing resources.

The new algorithm version extends the vertical range of the retrieval up to 10 km altitude (it was 7 km for version 4.1), allowing to better represent the dust distribution over high altitude areas such as the Tibetan Plateau, where dust is often found (e.g. Ge et al., 2014; Hu et al., 2020).

305 The retrieval *a priori* dust concentration was increased from 0.1 to 2 particles cm^{-3} , corresponding to a minimal *a priori* dust AOD of about 0.07 at 10 μm . This affects the areas/altitudes where dust is only rarely present, where the LIVAS climatology contains a very low dust extinction. Deviation from an *a priori* value of 0.1 particle cm^{-3} to a significant value expected during an event is very computationally costly (in the minimization cost function). With the previous setup (minimal *a priori* value of 0.1 particle cm^{-3}), it was seen that for rare dust events in unusual areas, the retrieval would either fail to converge or converge
310 but modify mostly the surface temperature. Those cases ended with a significant difference between the observed and modelled spectrum, and were then rejected during the quality control phase. It is also clear that this increased minimum *a priori* value has a tendency to push towards a small positive bias for low-load dust events, especially under cases with low sensitivity, where the retrieval output may be only slightly lower than the *a priori* value, leading to a dust AOD of about 0.05 at 10 μm . For the next algorithm version, a solution to this will be investigated, but it is certainly not straightforward, in particular if the computation
315 time can not be increased again.

The version of the radiative transfer tool used, RTTOV, was updated, from version 12.1 to version 13.0 (Saunders et al., 2020). This new RTTOV version also comes with a new version 13 of the RTTOV coefficients, containing updated gas line



parameters and a new optical depth parameterization. This has very minor impact on our simulations, leading to slightly lower spectral residuals in some cases.

320 The important new feature of RTTOV v13 for MAPIR is the availability of a new land surface emissivity database: the CAMEL climatology version 2 (Loveless et al., 2021), based on 16 years of data, which is now used in MAPIR v5.1. This is a significant change with respect to MAPIR v4.1, where the land surface emissivity from Zhou et al. (2011) was used. The latter data set has been used since the beginning of the MAPIR retrievals, as the only relevant data with the necessary spectral coverage and resolution. However, it suffers from a significant bias in the areas where dust is present almost every day during
325 some months (e.g. the Sahara during summer), due to the way the emissivity is obtained. The CAMEL climatology shows a more reasonable seasonal cycle, and is similar to the EUMETSAT IASI emissivity when available. Using the EUMETSAT IASI emissivity, distributed in NRT as 12 principal component scores, was also investigated. Its quality appears to be affected by the presence of clouds and dust, as one could expect. In addition, as this is a single spectrum retrieval, it contains an unreasonable variability (e.g. up to 5% over deserts within a month), with also unexplained year-to-year large differences that hinder the
330 production of a proper climatology. As a conclusion, the CAMEL v2.0 climatology seems the most adequate choice for land emissivity in MAPIR, at the current state-of-the-art. This modification leads to some significant changes in the retrieved dust AOD and vertical profiles, in some cases and areas.

The sea surface emissivity was also changed from the Newman et al. (2005) data set (constant everywhere over sea) to using the RTTOV internally calculated emissivity, using the IREMIS model, which is the default RTTOV sea emissivity (Hocking
335 et al., 2021). At nadir, it does not seem to make any difference with the previously used emissivity, but the viewing angle correction makes a small difference, which is expected to be an improvement even though not significant with respect to the uncertainty.

One additional improvement to the algorithm is the inclusion of propagation of temperature and water vapour profiles uncertainties, to the uncertainty in the aerosol concentration and AOD, through Rodger's formalism (Rodgers, 2000). This
340 requires the Jacobians (radiance derivatives) for those parameters, which comes with no additional computing costs in the case of RTTOV: the temperature and humidity Jacobians are always computed. The uncertainty in temperature and humidity input data are respectively set to 1 K and 10% (at all altitudes) which are the target accuracy levels for the IASI datasets.

The spectral noise in the retrieval has also been reduced. MAPIR v4.1 used to have a strong so-called noise inflation (Callewaert et al., 2019), meaning that the spectral noise used in the retrieval is much larger than the true instrumental spectral noise.
345 This was needed to allow convergence in many cases, and in general the noise inflation accounts for uncertainties that can not be modelled during the retrieval. The spectral noise in MAPIR v4.1 was set to 1 K in all retrieval windows. Now in MAPIR v5.1 it is set to 0.5 K in the first two windows, and 0.2 K in the third window. This new setup expectedly increases the information content of the retrieval, especially where the AOD is low, therefore allowing the retrieval to deviate more from the *a priori* value in those cases and to reduce the retrieved parameters' uncertainties. No convergence issues have been noticed with this
350 setup, but they arise if the spectral noise is further reduced.



Table 3. MAPIR version 5.1 changes with respect to version 4.1

Variable/parameter	MAPIR v4.1	MAPIR v5.1
Pre-filtering	Basic input data quality check, cloud free, spectral slopes outside the dust belt	Basic input data quality check
Latitude range	60° S to 60° N	global
DOM streams	8	4
Vertical grid	0:1:7 km	0:1:10 km
A priori min. conc.	0.1 part cm ⁻³	2 part cm ⁻³
RTTOV version	12.1	13.0
Land surface emissivity	Zhou et al. (2011)	CAMEL clim. v2, RTTOV internal
Sea surface emissivity	Newman et al. (2005)	IREMIS, RTTOV internal
Uncertainty	standard OEM	standard OEM, and propagated T and H ₂ O profile uncertainty
Spectral noise	1 K	0.5 K in win. 1 and 2; 0.2 K in win. 3
Retrieval win. 3	1020-1204 cm ⁻¹	1202.75-1204.75 cm ⁻¹
Land Ts <i>a priori</i> std	15 K	5 K

The third retrieval window, which was from 1202 to 1204 cm⁻¹ was slightly shifted to 1202.75 to 1204.75 cm⁻¹, to avoid some small methane absorption lines. This has no detectable impact on the retrieval results but is theoretically better and more robust.

Finally, the input IASI data has been selected to ensure consistency all along the IASI time series (for the climate version) or to ensure using the best available data (for the NRT version). This was already explained in details in Sect. 2.1.1. When doing that, the processing chain was also modified to be more efficient and more flexible with respect to quality control of the input data. One consequence of using the reprocessed IASI data is that the surface temperature *a priori* is now of better quality under dust storms, especially over deserts (where old IASI data versions produced highly biased data), allowing to reduce the surface temperature *a priori* standard deviation from 15 K to 5 K over land. That means that now in MAPIR v5.1 this standard deviation is set to 5 K everywhere.

2.3 Retrieval overall performance

Figure 2 shows the statistics of the IASI-A and IASI-C performance, in terms of pre-retrieval filtering, retrieval convergence rate and post-retrieval quality filtering (basic and including the cloud filtering).

For IASI-A, almost all observations are accepted for launching the retrieval (grey line in Fig. 2) and MAPIR v5.1 convergence rate is about 85% (green line), relatively low because the retrieval is run on all spectra, including those containing very




Figure 2. MAPIR v5.1 performance for IASI-A and IASI-C. In grey: the percentage of IASI observations for which the retrieval is run (passing all pre-filters), in green: the percentage of IASI observations for which the retrieval converges, in maximum 20 iterations; in blue: the percentage of retrievals passing the basic quality control criteria (i.e. RMS of spectral residuals below 0.75 K, $200\text{ K} < \text{surface temperature} < 350\text{ K}$ and dust AOD < 5 ; in pink: the percentage of retrievals passing all quality flags (i.e. also including those aiming to remove thin clouds, based on the difference between retrieved and *a priori* surface temperature))

thick clouds, which often lead to non-convergence. A total of about 70% cases pass the first set of quality filters (blue line, filtered on spectral residuals, reasonable surface temperature and AOD as detailed in Sect. 2.2.3). Finally, about 35% (of the total) pass the final quality test aimed at removing residual clouds (dark pink line). These numbers are stable along time, except a slight decreasing trend towards the end of the IASI-A life span. This reflects the long-term consistency of the EUMETSAT level 1 and level 2 IASI-A CDR, and most probably the small impact of the orbital drift towards the end of the instrument's life. The small spike of good quality data in April and May 2014 is unexplained.

For IASI-C, even though everything was expected to be consistent with IASI-A, after the processing a technical issue was discovered in the OFL IASI-C 1 data: the surface altitude is not provided for a significant portion of the pixels reported as cloudy (mostly those with a cloud flag value of 3 - partial cloud coverage in the pixel - or 4 - high or full cloud coverage of the pixel, but not all of those). This was not at all expected, as the surface altitude does not come from the retrieval itself. The absence of that parameter makes the MAPIR retrieval crash. Obviously, for the next (re)processing, the surface altitude in MAPIR will be provided from a separate digital elevation model to avoid such issues. This issue leads to running the retrieval only for 35 to 40% of the IASI observations, but then almost all those retrievals converge. After the full quality control, about 25 to 30% of the observation remain. This is significantly lower than for IASI-A, and a quick comparison of the data clearly shows that with IASI-C we are again missing some of the highly dusty pixels, falling in the cloudy category and missing the surface altitude. In addition, the statistics show at least one strong discontinuity in the quality control between 10 and 14 September 2022, that seems to correct for a decreasing trend in the year before, for which we could not find any explanation in the EUMETSAT IASI documentation. On the other hand, no discontinuity in the quality control is found due to the update in the PCS eigenvector basis on 30 March 2023.



385 3 Capability demonstration and evaluation

This section contains a demonstration of the capabilities of MAPIR v5.1, and an evaluation of the IASI MAPIR dust profiles and integrated AOD quality against different reference data sets. The latter reveals to be rather complex, because there are no appropriate reference data obtained from the TIR spectral range. The evaluation of the MAPIR data therefore has to be performed with an additional set of assumptions and approximations, which also contribute to the observed differences.

390 In most cases, aerosol information comes as a total column AOD. This will therefore be the main variable that we will evaluate. In addition, MAPIR provides a vertical distribution, and its validation is an even more difficult exercise, which will be briefly demonstrated here and fully provided in a separate publication (in preparation).

The subsections will first describe the data sets used in the evaluation, then further develop challenges for these comparisons, show the evaluation of the integrated dust AOD against AERONET reference observations, then discuss the information content
395 in the retrievals, the mean dust altitude and then showcase the profiling capability and validation against ground-based lidar observations from EARLINET. Finally, we will discuss uncertainties, long-term stability and cross-instrument consistency.

3.1 Instruments, models and data used for evaluation of the MAPIR retrievals

3.1.1 AERONET

AERONET (Aerosol Robotic Network) is a globally distributed network of ground-based remote sensing aerosol monitoring
400 stations established by NASA (National Aeronautics and Space Administration) and PHOTONS (PHOtométrie pour le Traitement Opérationnel de Normalisation Satellitaire) in collaboration with numerous international partners (e.g. Holben et al., 1998). Utilizing sun photometers, AERONET provides continuous, high-quality, and standardized measurements of aerosol optical properties, including aerosol optical depth (AOD), size distribution, and refractive index (e.g. Dubovik et al., 2002; Eck et al., 1999).

405 In this work, AERONET data is used as ground-based "truth" for comparison with the IASI MAPIR dust AOD. For that, we use the AERONET version 3 (Giles et al., 2019; Sinyuk et al., 2020) solar and lunar level 2 (cloud screened and quality assured) Spectral Deconvolution Algorithm (SDA) version 4.1 (O'Neill et al., 2003). From this data, we select the coarse mode total AOD at 500 nm (further referred to as CAOD, containing dust coarse mode aerosols and when present sea salt / marine aerosols) for comparison with the IASI dust AOD converted to 550nm. The challenges of these comparisons are discussed in
410 details in Sect. 3.2. All AERONET stations with median CAOD of at least 0.05 were considered in this analysis. This avoids comparing data from area with very little dust or with mostly marine aerosols, which would then mostly be comparing noise.

The AERONET data is filtered to remove events for which the reported uncertainty is large, following the method initially described in Capelle et al. (2018): we keep only data with a reported Root Mean Square Error (RMSE) lower than $0.05 + 0.15 * CAOD$. That filtering removes about 1/3 of the comparison points overall, especially in East Asia, but does not change the
415 comparison statistics globally (of IASI versus AERONET, which will be described later). The correlation between IASI and AERONET improves at Asian stations (where most of the "bad" data is removed), and slightly worsens at some "transport" stations, where the variability is low, because the filtering removes a significant number of high dust load events, lowering the




Figure 3. Reduction (in %) of AERONET - IASI-A collocation pairs due to AERONET quality filtering, including rejection of observations with fine mode fraction larger than 0.5.

variability even more. Finally, the AERONET data is filtered to remove mixed events of dust and fine aerosols such as those linked to biomass burning. Indeed, when two different aerosol types are mixed the partitioning between fine and coarse mode in the SDA algorithm is more difficult and the resulting CAOD is plausibly of lower quality. The filtering is done using the AERONET Fine Mode Fraction: when this fraction exceeds 0.5 the data is rejected. That results, for example, in about 45% of the IASI-A against AERONET solar comparison points being rejected, mostly of low AOD. Again the global statistics of the comparisons do not change significantly with this additional filtering. Figure 3 shows the impact of both data filters on the number of IASI-AERONET collocations per station, for the IASI-A full time series: from 4% data removed at Ragged Point (Barbados) to 90% data removed at Silpakorn University (Thailand).

Finally, stations were used only if at least 100 co-locations were found (the method is described in detail in Sect. 3.3.1), leading to using the 74 stations listed in Table A1.

3.1.2 EARLINET

Lidar (Light Detection and Ranging) is an active remote sensing technique widely used to study the vertical distribution of atmospheric aerosols. By measuring the backscattered signals from emitted laser pulses at certain wavelengths, it can acquire information on the shape, size and concentration of atmospheric particles. When ground-based, the signal may reach altitudes up to the lower levels of the stratosphere. Lidar is the most accurate means of studying the vertical resolution of aerosols, making it an excellent tool in mineral dust (e.g. De Rosa et al., 2025), satellite instrument evaluation (e.g. Michailidis et al., 2023) and aerosol classification studies (e.g. Voudouri et al., 2019). The European lidar Network (EARLINET) consists of more than 30 ground-based lidar stations distributed across Europe (e.g. Pappalardo et al., 2014) providing harmonized and quality-assured datasets. The network ensures data consistency through the application of the Single Calculus Chain (SCC) algorithm (e.g. D'Amico et al., 2015), an automated processing framework that standardizes lidar data analysis. In the present study, Level 1 data from the EARLINET station of Limassol, Cyprus (CYC, 34.68° N, 33.04° E, 11 m) (e.g. Mamouri et al., 2016) were used, a site strategically located in the Eastern Mediterranean, frequently affected by long-range transported mineral dust originating from the Sahara Desert and the Middle-East. The CYC station consists of a multiwavelength PollyXT lidar (e.g.



Engelmann et al., 2016) which performs continuous automated measurements under diverse atmospheric conditions, rendering it optimal for evaluation of polar-orbiting satellite data.

3.1.3 CALIOP

The Cloud–Aerosol Lidar with Orthogonal Polarization (CALIOP) is a nadir-pointing, dual-wavelength elastic backscatter lidar onboard the CALIPSO satellite, launched in April 2006 as part of the A-Train constellation with equator crossing time of 13:30 and 01:30. CALIOP provides global, vertically resolved observations of aerosols by measuring backscattered laser radiation at 532 and 1064 nm (Winker et al., 2009). At 532 nm, CALIOP measures both parallel and perpendicular polarized backscatter, enabling the computation of particulate depolarization ratios and supporting aerosol type discrimination, while the 1064 nm channel measures total attenuated backscatter. The instrument has a nominal horizontal sampling of 333 m along track and a vertical resolution of 30–60 m in the troposphere.

This study uses CALIOP Version 4.2 Level 2 aerosol profile products. Aerosol backscatter and extinction coefficients are retrieved from calibrated attenuated backscatter profiles by removing the molecular contribution and applying aerosol-type-dependent lidar ratios. Aerosol subtyping (e.g. dust, smoke, polluted continental) in Version 4.2 is based on an improved classification algorithm that exploits layer-integrated backscatter, depolarization ratio, spectral dependence, and layer altitude (Kim et al., 2018). We use the CALIOP aerosol type and 532 nm extinction profiles, to qualitatively compare with IASI MAPIR dust concentration profiles. The data is quality-controlled as advised in Winker et al. (2013): only data with Cloud Aerosol Discrimination (CAD) score between -100 and -20 (avoiding detection artefacts) and with extinction quality flag pertaining to successful retrievals (value of 0,1,16 or 18) are kept (e.g. Song et al., 2021; Ridley et al., 2012; Mielonen et al., 2009; Gkikas et al., 2016).

3.2 Challenges of the evaluation

The main challenge arises from the absence of reference aerosol data in the TIR. That means that the comparison of the MAPIR data has to be done with data gathered in the visible (VIS) or near-infrared (NIR) spectral ranges, and from very different types of instruments. In the case of aerosols, the implications are even wider because the spectral sensitivity to aerosols is largely different in the VIS and TIR.

In a nutshell, TIR allows observations of (almost) only coarse mode mineral aerosols, while VIS bears the radiative impact of all aerosol types and all sizes. The aerosol effect on radiation is the combination of absorption and emission in the TIR, and scattering. The first depends on the particle size, composition and absorption features, while the latter depends on the ratio between the particle size and the wavelength. Both are significantly different in VIS and TIR.

Light absorption depends on the particle composition and on the particle surface (hence size). Absorption in the TIR is due only to mineral aerosols, while VIS absorption is mostly due to black carbon, with some contribution of dust and organic carbon (e.g. Li et al., 2022). Even within the dust absorption, the sensitivity differs from TIR to VIS: in VIS the absorption mostly depends on the iron oxide content, while in TIR absorption is due to clays, quartz, calcite, kaolinite and feldspar (e.g.,



Di Biagio et al., 2019, 2017; Li et al., 2021). The coarse mode particles obviously absorb significantly more than fine particles (for the same number of particles).

475 As for the scattering, in TIR it is due to coarse particles only, while in the VIS, there is also a significant fine-mode scattering contribution (e.g. Dufresne et al., 2002; Li et al., 2022).

There are however ways to mitigate those differences and perform meaningful comparisons between TIR aerosol information and VIS reference aerosol data. In the VIS spectral range, it is possible to separate the fine and coarse AOD (e.g. for AERONET O'Neill et al., 2003). The main assumption here is that the spectral variation of the coarse mode is neutral (within the VIS
480 range). The VIS coarse mode AOD obtained this way also contains thin cloud contamination, for which a removal is attempted based on the temporal variability, and does not contain only the dust aerosol contribution: sea salt particles are also included, and volcanic ash when present. This VIS coarse mode AOD is, to our knowledge, the best parameter to be compared with the TIR-obtained dust AOD. However, obviously, the separation is not perfect, in particular when there is a mixture of aerosols.

Finally, in addition to selecting the VIS AOD that best matches the types of particles observed in the TIR (i.e. coarse mode
485 AOD, as explained above), there is a need to convert the AOD obtained in the TIR to a VIS AOD. Here, we convert from the TIR AOD (reported at 10 μm) to the standard VIS wavelength of 550 nm. Theoretically, the conversion is simple, using the particle cross-section at the two wavelengths (Eq. 1). The cross-sections are obtained using the particle optical properties used in the retrieval (see Sect. 2.1.2) and a Mie code (spherical assumption, as in MAPIR retrievals).

$$AOD_{550nm} = AOD_{10\mu m} * \frac{X_{sect550nm}}{X_{sect10\mu m}} \quad (1)$$

490 After conversion, the reported AOD at 550 nm is the coarse mode dust AOD, representing particles that are observable in the TIR. The conversion adds a significant uncertainty linked to the difference in sensitivity at both wavelengths (already discussed above) and to the fact that the cross-sections are calculated using a Mie code, with the approximation of spherical particles. It is well-known that mineral dust particles are not spherical (e.g. Kandler et al., 2007). The shape of the particles impacts scattering phase functions, with only a small effect in the TIR where the particle size is smaller than the light wavelength (e.g.
495 Koepke et al., 2015). However, non-sphericity has a significant impact in the VIS and especially in conversions between VIS and TIR (e.g. Song et al., 2018).

Clarisse et al. (2019) made a literature review of the different conversion factors for mineral dust AOD, from 10 μm to 550 nm and highlighted that they range from 0.9 to 3.5, the VIS AOD being larger than the TIR AOD. Our conversion factor of 1.78 falls in that range, a bit lower than the average value of 2.2.

500 Finally, in TIR particle thermal emission occurs, with the same spectral dependency as the absorption and with an intensity depending on the particle temperature, hence on its altitude. The AOD, on the other hand, does not contain any altitude information, and is simply the vertically integrated aerosol number concentration multiplied by the extinction coefficient at the desired wavelength. That means that the same TIR AOD with a different underlying vertical profile actually produces a different impact on radiance. It also means that in a TIR aerosol retrieval the altitude of the aerosols is highly important and
505 any bias in it directly leads to a bias in the retrieved TIR AOD.



Table 4. Literature survey of co-location methods between different satellite instruments and AERONET.

Instrument	Co-location criteria	Reference
CALIOP	80 km CALIOP averages, 1 h AERONET averages	Proestakis et al. (2024)
IASI	0.25°, 1 h, both averaged	Capelle et al. (2018)
IASI	30 km, 30 min, both averaged	Clarisse et al. (2019)
IASI	0.25°, 1 h, single-point comparisons	Callewaert et al. (2019)
SEVIRI	single pixel over the station, 10 min	Bernard et al. (2011)
MIDAS	25 km, 2 h, both averaged	Gkikas et al. (2021)

All these considerations are highly important and need to be kept in mind when comparing those to data obtained from UV/VIS instruments. Failing to do so would lead to erroneous conclusions and a wrong evaluation of the TIR dust observation and capabilities.

3.3 Integrated AOD comparison against AERONET

510 3.3.1 Co-location method

The method used to co-locate the ground-based reference and satellite observations may have a strong influence on the comparison results. Ideally, one would wish for the smallest time and space differences, but one also requires sufficient comparison data for a reasonable statistical analysis. Finally, one can compare each single pair of co-located data (maximising the number of comparisons) or averaged data (reducing the noise in the comparisons). A short literature survey shows (Table 4) different approaches in different publications / analyses.

In this work, we have tested different criteria and compared the results: co-locations distances from 25 to 150 km and times from 15 min to 1 h, single point comparisons or averages within the co-location criteria, with and without outlier removal. General conclusions remain very similar for the different co-locations criteria. Differences are seen at some stations, with usually an obvious explanation. First, at stations located close to sources, the correlation coefficient tends to decrease when loosening the co-location constraints, attributed to the high horizontal variability of the atmospheric dust in those areas. Second, at stations with only few data, loosening the co-location constraints allows more data to enter the comparisons, leading to different statistics (sometimes better and sometimes worse). Third, at stations with little variability in the dust / coarse aerosol load, far from sources, loosening the co-location constraints improves the statistics as more events are included in the comparisons, enhancing slightly the variability and leading to a slightly better correlation coefficient.

Using mean values in the comparisons instead of comparing single observations smooths the local variability (in time and space), and might lead to misinterpretation when using loose co-location criteria. Indeed, single observation comparisons with loose criteria show a large spread of both IASI and AERONET data for a single event (especially close to sources where the variability is larger), while using averaged values only shows a slight correlation coefficient reduction. Having performed



Table 5. Statistics of IASI-MAPIR AOD converted to 550nm versus AERONET SDA coarse mode AOD

Parameter	IASI-A vs solar	IASI-A vs lunar	IASI-C vs solar	IASI-C vs lunar
N points	47 310	1 024	12 167	959
Correlation coeff. (all stations)	0.78	0.67	0.79	0.7
Mean absolute bias (std dev)	-0.002 (0.11)	-0.011 (0.18)	0.002 (0.09)	-0.001 (0.09)
Median absolute bias	0.012	-0.010	0.011	0.008
Median relative bias	9%	-5%	10%	6.2%
Comp. with bias lower than 0.1	76%	58%	81%	78%

comparisons with all types of co-location criteria allows a better understanding of both data sets, and therefore also a better interpretation of the comparison results.

The results shown and discussed here are for maximum 25 km distance between the IASI pixel centre and the AERONET station, maximum 15 min between the observations, and single comparisons (no time or space average). This assumes that the dust aerosol fields is homogenous within 25km and 15 min. Finally, the worst 3% of the comparisons are removed from the statistics for each station, as probable outliers or cloud misclassification. This only slightly changes the statistics. The same comparisons are performed using both solar and lunar AERONET data, but statistics are computed separately as the AERONET data quality might differ between solar and lunar measurements acquired under very different circumstances and with different sensitivity. Stations for which there is less than 100 colocations are removed from the analysis.

3.3.2 Global statistics

In this section, we analyse the comparisons globally, all co-locations from all-stations grouped together before extracting statistics.

Table 5 provides the comparison statistics for IASI-A and IASI-C separately (dust AOD converted from 10 μm to 550 nm), and for solar and lunar AERONET observations separately (SDA coarse mode AOD). The overall correlation coefficient is almost 0.8 between both IASI and AERONET solar observations, and almost 0.7 for comparisons with AERONET lunar observations, which shows very good capability of IASI and MAPIR to observe dust events and follow their seasonal cycles. The mean absolute bias is close to 0 with a standard deviation of about 0.1, except for IASI-A against lunar AERONET where it is larger (0.011) and with about twice the standard deviation. The median absolute bias is of about 0.01, positive for all groups except again for IASI-A against AERONET lunar where it is negative. The median relative bias is about 10% for both IASI instruments against AERONET solar, and about 5% against AERONET lunar. The median absolute bias is positive in all cases except IASI-A against AERONET lunar. The ratio of comparisons with a bias lower than 0.1 lies between 76 and 81% except again for IASI-A against AERONET lunar, where it is only 58%. We recall here the important considerations from Sect. 3.2, which have a strong impact on absolute value comparisons while mostly not affecting correlation.



Figures 4 and 5 show graphically the comparison results of IASI (A and C separately) versus AERONET (solar and lunar separately) observations. Panels (a,d) show the density plot of the IASI AOD against the AERONET AOD, with its linear regression line and equation. Panels (b,e) show scatter plots of the MAPIR bias versus the AERONET AOD and data bins with constant number of points (median and interquartile (IQR) are shown). Finally, panels (c,f) show histograms of the MAPIR bias, together with its median and quartiles. These figures show that even though the correlation, mean and median bias highlight very good performance of MAPIR v5.1, there is a clear overestimation by MAPIR v5.1 of the low 550 nm AOD values, and a linear dependence of the MAPIR 550 nm AOD bias with the AOD, the bias being positive for small AOD and becoming negative from AOD of about 0.1. The histograms also show almost Gaussian distributions of the bias, with a maximum skewed towards positive values while the median is close to zero. The overestimation of low AOD is clearly linked to the lack of IASI sensitivity for very low AOD and therefore also exists in the 10 μm AOD data. In those cases, the deviation from the *a priori* during the retrieval is difficult. The *a priori* minimal value was set to 2 particles cm^{-3} all over the retrieval range, corresponding to a dust AOD of about 0.08 at 10 μm , or about 0.14 at 550 nm. The retrieval manages to deviate to lower values in absence of dust, but as will be seen more precisely in Sect. 3.3.3 there is a lower threshold that the retrieval does not seem to be able to pass. The underestimation of high 550 nm AOD, could be linked to an underestimated AOD conversion factor from TIR to VIS (see Sect. 3.2), and the linear dependence of the bias with the AOD aligns with such a possibility. Indeed, using a TIR to VIS conversion factor of 2.85 (instead of the physically calculated 1.78) leads to a constant bias (0.1 +/- 0.05) of MAPIR 550 nm AOD with respect to AERONET solar CAOD. To validate the absolute value of the IASI dust TIR AOD and conversion factor to VIS, it would be necessary to have some reference data directly at 10 μm , which is currently not available. Therefore, these AOD comparisons are interesting as they help understand the product (and are requested by users), but the absolute value of the bias is not truly a product quality assessment. The correlation is however relevant, as it assesses relative changes and not the absolute values.

Comparisons with lunar AERONET data show similar results as for solar for IASI-C (although with much less points, and at much less stations) but much more noise for IASI-A, especially for high AOD. This is most probably due to unfiltered thin clouds, leading to artificially high dust MAPIR AOD in some cases even though these should not appear in the collocations with AERONET data as the latter also undergo a cloud filtering. The absence of such issues for IASI-C comparisons with lunar AERONET highlights that the IASI-C data was partially cloud-filtered before the retrieval, due to the absence of the surface altitude data for most of the cloudy pixels (see Sect. 2.3). This unwanted filtering however also clearly removed some of the high AOD dust cases, as seen on both IASI-C comparison plots for which the last equal number bin (black, in panel (c)) occurs for a significantly lower AOD than for IASI-A.

3.3.3 Stations

Figure 6 shows the Pearson correlation coefficient between the IASI-A dust AOD converted to 550 nm and solar AERONET SDA coarse mode AOD at 500 nm, for the 68 stations passing the criteria detailed in Sect. 3.1.1. The corresponding map for IASI-C comparisons is very similar, with some different stations due to the different time frame, and a total of 41 stations. The comparisons with AERONET lunar data contain only 8 stations for IASI-A and 7 for IASI-C and the correlation coefficients




Figure 4. MAPIR v5.1 comparisons for IASI-A (a,b,c) and IASI-C (d,e,f) dust AOD converted to 550 nm against AERONET solar data SDA coarse mode AOD at 500 nm. (a,d) Density plot of AOD, IASI-A versus AERONET, with its linear regression and equation; (b,e) Scatter plot of the MAPIR AOD bias versus the AERONET AOD, with bins of equal number of points. (c,f) Histograms of the MAPIR AOD bias.




Figure 5. Same as Fig. 4 but for comparisons with AERONET lunar data.




Figure 6. Correlation coefficient between MAPIR v5.1 comparisons for IASI-A dust AOD converted to 550 nm and AERONET solar data SDA coarse mode AOD at 500 nm, for all stations with at least 100 colocations

are again similar, slightly lower for IASI-A as already seen in the global statistics. The very limited number of lunar stations in these comparisons arises from the strong constraint of at least 100 colocations at a station to keep it in the analysis. Table A1 summarizes all these statistics by station.

For most stations, the correlation coefficient is about 0.8, including close to sources and after long-range transport towards the Caribbean. This shows the good capability of MAPIR / IASI to identify the dust events, their relative intensity and seasonal cycles. The stations with the worst correlation coefficient (below 0.5, e.g. in Ascension Island, Arica or Manila) show very little AOD variability, and highly discontinuous time series, rendering the statistical analysis unfit in terms of correlation. The same occurs for Zinder airport lying in the Sahara but reporting only very episodic data, only for strong events, making it again statistically unfit for extraction of a correlation coefficient. There is also a group of stations with a correlation coefficient around 0.6-0.7, lying mostly in the Middle-East. Most of them lack full seasonal cycle observations, and therefore again lack a part of the natural variability, reducing the capability to get a good correlation coefficient from noisy data.

Figure 7 shows the time series of AERONET solar and lunar data, and of IASI-A and IASI-C data split between daytime and night-time, for two AERONET stations: Dakar Belair and Ragged Point. Those stations were selected from the few stations containing solar and lunar AERONET observations, and during the IASI-A and IASI-C overlap time, one relatively close to sources and the other after long-range transport. We emphasize that in those plots, all data are plotted and not only the colocations. This provides a lot of useful information. First, one can see a continuity and consistency between AERONET solar and lunar data, between IASI daytime and nighttime data, and between IASI-A and IASI-C. Second, MAPIR gets the seasonal cycles very well, even after long range transport. Third, there is a minimal AOD (about 0.05 at 550 nm) under which the MAPIR algorithm does not seem to be able to go, as seen in Sect. 3.3.2. Fourth, for each dust event, the AERONET data contains a lot of observations, spanning a range of AODs in a short time range (the vertical lines of grey/black points). This obviously renders noisy the comparisons with the twice-per-day satellite data, unless a very strong time constraint is applied - and this was observed when testing the different colocation criteria. And finally, one can also see in those plots some plausible remaining unfiltered cloudy cases in the IASI data: the isolated high IASI AOD when AERONET does not show an event.

Figure 8 shows the time series of colocations for the two same stations, separately for IASI-A and IASI-C. For Ragged Point, there are not enough lunar colocations for IASI-A, therefore only IASI-C is shown. Those plots clearly show again




Figure 7. Time series of all AERONET (solar and lunar) SDA coarse mode AOD at 500 nm and of IASI-A and IASI-C dust AOD converted to 550 nm (separated day and night) for the time span with available AERONET data, at (a) the Dakar Belair AERONET station and (b) the Ragged Point station. Note that these are full data series, not only collocations.

the nice continuity in all data sets, and the good capability of IASI-MAPIR to observe seasonal cycles. They also show the MAPIR underestimation of the 550 nm AOD larger than 0.1 and increasing underestimation when the AOD increases, as already discussed in Sect. 3.3.2. However, we do not see here any possible unfiltered clouds in the IASI data. This is due to the strong collocation criteria and the fact that AERONET data is also cloud-cleared, therefore the remaining cloudy IASI should not have any colocated AERONET data.

3.4 Degrees of freedom

The number of degrees of freedom (or DOF) are the number of distinct information pieces one can extract from the vertical profile. The DOF considers the instrument / retrieval pair sensitivity to the true atmospheric state, and is extracted from the averaging kernel matrix as its diagonal. For the specific case of TIR observations, the DOF will largely depend on the difference between the surface temperature and the target species' temperature, which relates to its altitude. With MAPIR, the DOF reaches about 2.5 for the best cases (large AOD, large temperature contrast). Figure 9 shows the dust AOD at 10 μm (a) and the DOF (b) for 19 June 2020, during a huge Saharan dust outflow towards America. The figure shows the MAPIR IASI-A data separately for the morning and evening overpasses. One can clearly see the DOF being around 1 or even lower in areas without dust, while the DOF may reach up to 2.5 where there is a large dust AOD and expected large temperature contrast, such as above the desert. As was detailed for the previous MAPIR version 4.1 in (Callewaert et al., 2019), when performing the retrieval in the logarithmic mode the averaging kernels tend to peak at altitudes and locations where there is dust, explaining partly the low DOF in absence of dust.




Figure 8. Time series of collocated AERONET (solar and lunar) SDA coarse mode AOD at 500 nm and of (a) IASI-A and (b) IASI-C dust AOD converted to 550 nm (separated day and night), at the Dakar Belair AERONET station, and (c) IASI-C at the Ragged Point station.




Figure 9. Dust $10\ \mu\text{m}$ AOD (a) and number of degrees of freedom (DOF, b) for MAPIR IASI-A, 19 June 2020. Morning and evening overpass data are plotted separately



3.5 Mean altitude

As detailed in Sect. 2.3, the MAPIR data set also contains a mean altitude (above sea level), provided when the dust AOD is greater than 0.05 and the information content larger than 1.25, the latter being the most strict criterion. We do not aim here at validating the mean altitude values. This was done for the previous MAPIR version (Callewaert et al., 2019) while in this work we will provide some direct profile validation and preliminary mean altitude validation in Sect. 3.6.2.

It is important for the user to understand that even though the mean altitude is provided only when there is a reasonable vertical sensitivity, there still remain cases where the mean altitude is simply about half the total vertical range (i.e. about 5 km). Those cases are difficult to filter out without extremely strict filters, which would then end up in only very little cases where the mean altitude is provided, reducing its usefulness. Figure 10 shows the mean altitude from MAPIR IASI-A dust profiles for 19 June 2020 (the same day as in Fig. 9 showing AOD and DOF), separately for morning and evening overpass. One can see there that for this huge event the transport towards America occurred (as often) at about 2.5 to 3 km altitude, and one can spot the plausible source area as the location in the central Sahara with very low mean altitude. There were also other smaller dust events that day throughout the dust belt as well as in Australia. The highest mean altitudes (of about 5 km) at the edges of the Saharan part of the plume occur in parts of the plume with lower sensitivity, at the limit of the criterion to provide the mean altitude, and are close to the middle of the profile vertical range. These kind of "artefacts" occur at different places, for a limited number of IASI observations during the day. When aggregating the data at monthly (Fig. 11 a) or yearly (Fig. 11 b) scale, one can see some areas popping out with an unreasonable high mean altitude (even above 6 km), mostly in South Africa and South America in areas where there usually is no dust. Therefore it is safe to assume that these averages result from a limited number of artefact events. On the other hand, one can nicely see for example a mean dust altitude of more than 5 km over the Tibetan Plateau, which is expected.

3.6 Profiles

3.6.1 Qualitative comparisons with CALIOP

This section shows qualitative illustrations of the IASI MAPIR dust concentration profiles compared to the CALIOP high resolution extinction profiles. The IASI MAPIR data is averaged around every 5 CALIOP observations along its track, with a maximum 50 km distance and 6 h time difference. The second criterion ensures comparing day-time or night-time observations separately. IASI and CALIOP observations always bear a time difference of 3 to 5 h due to their respectively mid-morning (9:30) and mid-afternoon (13:30) orbits, and depending on the IASI pixel position within the cross-path swath.

On Figs. 12 to 14, the CALIOP data is plotted along its track for selected relevant orbit parts, first as aerosol type (all types included), then as extinction at 532 nm for only the "dust", "dusty marine" and "polluted dust" types. Then the IASI MAPIR data is also plotted as concentration profiles (the main retrieval output), and a daily IASI 10 μm AOD map is provided with the CALIOP track plotted in teal.

For the first example (Fig. 12), we look at a night-time Middle-East dust plume on 28 October 2018, with IASI-A. One can see a significant dust plume between 50 and 20°N along the selected CALIOP track, with the most intense dust extinction




Figure 10. Mean altitude (km above sea level) for MAPIR IASI-A, 19 June 2020.




Figure 11. Mean altitude for MAPIR IASI-A averaged over the month of June 2020 (a) and the whole year 2020 (b). Morning and evening overpasses are included.

and concentration at about 30° N (Afghanistan / Iran border)), between 1 and 2 km altitude, with smaller dust amounts up to 4 km altitude. The IASI MAPIR dust profiles place the dust at the same location as observed with CALIOP, both horizontally and vertically. With IASI, dust is also observed right above the surface at about 45° N along that track, where CALIOP typing determined the presence of continental aerosols. As IASI observations are not sensitive to other aerosol types, it is well plausible that the observed aerosols are a small amount of dust, maybe with different properties than the usual desert dust.

The second example (Fig. 13) is the next orbit, on the same day, showing dust over Syria, Jordan, then Egypt and the Sudan and finally in the South of the African continent. A first plume is detected with both CALIOP and IASI as most intense around 30° N (in Jordan), again at 1 to 2 km altitude. The plume extension towards 20° N (towards the Red Sea then Egypt and the Sudan) is more intensely observed with IASI than with CALIOP, which could be due to the time difference and plausibly



670 looking at the source area. There is a second dust plume detected with IASI at 20-30° S (in Botswana and South Africa), for which CALIOP type is "polluted dust", with a very low extinction, while IASI shows significant dust amounts.

The third example (Fig. 14) is selected during an unusually intense Atlantic dust transport case, the Godzilla event in June 2020 (the same event already used for some other illustrations). The figure shows that during the event, even after long-range transport, the dust column is very intense and reaches high altitudes with respect to the usual transport altitude. Indeed, CALIOP
675 and IASI concur in placing large amounts of dust up to 3-4 km altitude all across the plume's head, close to the Antillas. The CALIOP signal is even saturated and does not provide information below 2 km altitude.

These examples all show the good dust profiling capabilities with IASI data and the MAPIR algorithm, even over land or in unusual situations.

3.6.2 Assessment against ground-based lidars

680 To evaluate the MAPIR dust concentration profile and altitude, two approaches were used. During the first approach, the one-step POLIPHON (POLarization LIdar PHOtometer Networking) (Mamouri and Ansmann, 2014) method is implemented, in order to separate the contribution of dust particles to a lidar signal at 532 nm. The threshold particle depolarization values which are an essential component of the technique are derived from bibliography (Teschke et al., 2009). The retrieved lidar dust backscatter profiles were then regridded from the instrument's native vertical resolution of 7.5 m to 1 km, in order to ensure
685 the consistency with the IASI dust profile vertical grid. Lastly, the dust concentration (particles cm⁻³) is calculated using the following equation, proposed by (Mamouri and Ansmann, 2014):

$$M = \frac{v}{\tau} \frac{\beta_d S_d}{V_d}$$

where $\frac{v}{\tau}$ is the global POLIPHON conversion factor, v is the volume concentration of dust, τ is the aerosol optical depth at 532 nm, β_d is the dust backscatter coefficient, S_d is the lidar ratio, and V_d is the volume of a spherical dust particle with an effective radius of 0.61 μm , a value derived from AERONET (Aerosol Robotic Network) inversion data for the Limassol site
690 (Ansmann et al., 2019). For the second approach, the same procedure is implemented with an extra step added at the end. The final dust concentration lidar signal is processed using the following equation (Rodgers and Connor, 2003):

$$x'_L = x_a + A(x_L - x_a)$$

where x_a is the *a priori* profile provided by LIVAS Section 2.1.3, x_L is the lidar dust concentration profile, A denotes the averaging kernels, and x'_L is the convolved lidar dust concentration profile under the effect of MAPIR parameters. In order to fully assess the capabilities and limitations of the validation process, it is performed both with and without applying the
695 averaging kernel correction. The dust mean altitude from the lidar observations is calculated in both cases as the median of the profile; that is, it corresponds to the altitude at which half of the dust particles are above and the other half below, in the same way as performed within the MAPIR product.




Figure 12. Comparison of CALIOP and IASI-A MAPIR for 28 October 2018, night-time. (a) CALIOP aerosol type; (b) CALIOP 532 nm extinction for the aerosol types "dust", "polluted dust" and "dusty marine"; (c) IASI-A MAPIR dust concentration profile along the selected CALIOP orbit part; (d) IASI-A MAPIR dust AOD (daily average), with the track of CALIOP for the selected orbit part.

In total, 12 dust transport cases over Limassol, Cyprus, occurring between March and May 2024, were used for the MAPIR evaluation against ground-based EARLINET lidar observations. Each event was identified through a synergistic approach combining lidar and AERONET observations with forecast and back-trajectory models such as the NOAA hybrid single particle Lagrangian integrated trajectory (HYSPPLIT) model (e.g. Draxler and Hess, 1998). For ten of the cases, lidar profiles were temporally averaged over 60 minutes, while two were averaged over 40 minutes. Since the PollyXT system operates continuously, both lidar and IASI were measuring simultaneously in almost all instances. For this study, only the closest satellite pixel to




Figure 13. Same as Fig. 12, for a different CALIOP orbit of the same day.

the lidar station is used in the comparison. The mean distance between the IASI pixel and the lidar site for the selected cases was 19.32 ± 13.27 km, which is considered acceptable given that the satellite instrument has a pixel diameter of approximately 12 km.

In Fig.15, the dust concentration differences between lidar and IASI are shown for each vertical layer, for both the regridded (a) and the convolved (b) lidar profiles. For each selected case, the difference between lidar and IASI is calculated for each layer, and the average over all 12 identified dust cases is computed per layer. When compared with the regridded lidar profile, IASI tends to underestimate concentrations below 6 km, with maximum differences reaching approximately 25 particles cm⁻³ (corresponding to a 1 km layer AOD of about 0.1 at 10 µm) between 2.5 and 4.5 km. Negative differences at higher altitudes




Figure 14. Same as Fig. 12, for an orbit of 19 June 2020 day-time and using IASI-C MAPIR data.

indicate that MAPIR/IASI detects dust where lidar provides slightly smaller values. When the lidar profiles are processed using the MAPIR averaging kernels and the associated *a priori* profiles, the correlation improves significantly. In the layers where the first approach shows maximum differences, between 2.5 and 4.5 km, discrepancies are reduced by up to 80%. It is worth noting that the first layer provided by MAPIR, from 0 to 1 km, is not included in the evaluation, since the CYC lidar overlap altitude is around 600 m. These preliminary findings highlight the importance of applying the averaging kernels to the validation process in the case of dust profiles. They also show the good profiling capability of MAPIR/IASI, especially considering that the absolute value of the concentration in the MAPIR retrievals highly depends on the dust optical properties used in the retrieval (while the 10 μm AOD depends much less on those properties).




Figure 15. Dust concentration differences at each layer, regridded (a) and convolved (b) lidar vs IASI/MAPIR, for 12 dust cases over Limassol, Cyprus.

In Fig. 16, the comparison of the dust mean altitude derived from the 12 colocated lidar and MAPIR profiles is presented. The correlation between the convolved lidar profile and the MAPIR retrieval is substantially higher ($r = 0.86$) than that obtained with the simply regridded lidar profile ($r = 0.19$). Once again, the implementation of the averaging kernels and the *a priori* profiles proves to be of immense importance in the final products. The difference between the regridded lidar dust altitude and the MAPIR retrieval has a mean value of 0.18 ± 1.47 km, while the application of the convolution approach reduces the mean difference to 0.04 ± 1.11 km. It is worth noting that in only 2 out of the selected 12 cases, the difference in mean dust altitude between IASI and MAPIR exceeds 2 km.

Overall, the MAPIR profiles slightly underestimate dust particle concentrations at mid-altitudes, between 2.5 and 4.5 km, but show substantially better performance at higher altitudes. Applying the MAPIR averaging kernels and the LIVAS *a priori* profile to the lidar signal significantly reduces the discrepancies between the instruments. Regarding the dust mean altitude, both approaches yield satisfactory agreement between lidar and IASI, with mean differences below 0.2 km. Finally, it is worth noting that these results provide an initial assessment of the product's performance at a single site, analysing 12 dust transport cases. A full, multi-station validation effort is currently in preparation and will extend the findings presented here.

3.7 Uncertainties

As described respectively in Sect. 2.2.2 and 2.2.3, the retrieval provides the uncertainty on the dust concentration profile, which is then used to compute the uncertainty on the dust AOD. That uncertainty, based on the OEM formalism, contains separately the uncertainty linked to the (inflated) spectral noise, to the vertical smoothing, and to the propagated uncertainty of the vertical profiles of temperature and humidity. Table 6 gives an indication of the median and inter-percentile (IP) of 68 (i.e. ± 1 -sigma)




Figure 16. Dust mean altitude comparison, regridded (blue) and convolved (red) lidar vs IASI/MAPIR, for 12 dust cases over Limassol, Cyprus.

Table 6. Overview of MAPIR dust AOD $10\ \mu\text{m}$ uncertainty

Cause	Input	Median AOD unc.	IP68 AOD unc.
Spectral noise	0.5 K or 0.2 K	0.005 (clear-sky) - 0.008 (Tropical summer)	Up to 0.02 (Tropical summer and Antarctica)
Vertical smoothing	Averaging kernels, Sa matrix	0.005 to 0.008	0.005
Atmospheric T	1 K, T Jacobians	Negligible	Non-negligible only in Antarctica: up to 0.004
Atmospheric H ₂ O	10%, H ₂ O Jacobians	<0.002	Very small except in the Tropical summer: up to 0.01
Total OEM propagation	All components above	0.005 to 0.01	Low variability except in the North Tropics and Antarctica: up to 0.03

for the different uncertainty components. The statistics were calculated for the dust $10\ \mu\text{m}$ AOD, by latitude bands and by month. Here only a summarized overview is provided, to simplify the complex information.



740 In addition, a short sensitivity study was undertaken to evaluate the impact of the surface emissivity uncertainty on the dust retrieval results. It is not possible to properly propagate such an uncertainty through the OEM, because RTTOV does not provide emissivity Jacobians. The study was therefore undertaken by modifying the surface emissivity manually (5% reduction), and analysing the impact it has on the retrieval results. This impact depends, among others, on the dust AOD, on the satellite viewing angle, on the surface type and temperature. The only clear conclusion is that 5% uncertainty in the surface
745 emissivity leads to minimum 0.015 AOD uncertainty in absence of dust, and maximum 0.05 AOD uncertainty in presence of dust. It also leads to a mean altitude uncertainty of about 0.5 km.

Combining both analyses leads to an important conclusion for the users: any dust AOD $10\ \mu\text{m}$ below 0.02 should be considered as noise, and any dust AOD $10\ \mu\text{m}$ below 0.06 should be considered with a highly critical eye, especially over land where the surface emissivity uncertainty is larger. As the uncertainty clearly depends on the precise case, the user is
750 advised to read the calculated OEM uncertainty for each retrieval, provided in the data (only the total OEM uncertainty is provided in the data, in order to spare storage space), and to keep in mind the additional plausible uncertainty due to the surface emissivity uncertainty, of up to 0.05.

3.8 Long-term stability

The long-term stability was assessed with 2 different exercises. First, the AERONET comparisons were used, looking for
755 potential drifts in the MAPIR IASI bias. Second, the full IASI MAPIR data (both instruments) was used to plot time series of daily averages within large areas, looking for any discontinuity or unexpected feature.

For the first analysis, a simple linear regression analysis was performed on time series of MAPIR bias versus AERONET solar data. We kept only stations for which the AERONET data spans at least 5 years, and contains at least 50 points per year on average (meaning there can be gaps - otherwise almost all stations are rejected). The same analysis was attempted
760 using AERONET lunar data but no station passed those criteria. The p-value was used to consider only the trends/drifts that are significant at the 95% confidence interval. Fig. 17 shows the results of this analysis for IASI-A, with in grey the stations for which the analysis was run without significant trend/drift (representing 19 over the 33 stations analysed). It shows that only very small significant drifts are detected, below the absolute value of 0.006 (for the 550 nm AOD bias) per year except at Medenine where it reaches 0.009. Medenine provides data only for a year from mid 2014 to mid 2015, and 1.5 year from
765 end 2018 to early 2020. The other stations close to it show either a non-significant drift (Tunis-Carthage) or a negative drift (Gozo). In general, the sign of the drifts is inconsistent between the different stations. We performed the same analysis using the Theil-Senn robust linear fit and, although some stations shift between significant and non-significant drift (both ways), the general conclusions remain the same. We consider these results as indication that there is no drift in the MAPIR IASI-A data, as would be expected considering that all inputs are consistent in time, to the best of our knowledge.

770 The same analysis was run for IASI-C, and shows only one significant drift over the 8 stations analysed: at Medenine, with -0.01 550 nm AOD per year (while it was positive for IASI-A). The data at this station has again a long gap between early 2020 and early 2022, meaning that the 5 years time span constraint is met, but at the beginning of the series there are only a few months of data, followed by 2 years without data. In addition, for the 2 stations very close to Medenine (Gozo and Ben




Figure 17. Linear trend in the bias of MAPIR IASI-A versus AERONET solar observations (AOD 550 nm per year). Stations plotted in grey are those included in the analysis but without significant trend at the 95% confidence (p value larger than 0.05).

Salem), no significant drift is detected in MAPIR. Therefore we conclude that there is currently no indication of a drift in the
 775 MAPIR IASI-C data. There does not seem to be any impact of the discontinuities pinpointed in the quality control (Sect. 2), or
 of the change of the PCS eigenvector basis on 30 March 2023.

For the second exercise, we plotted MAPIR IASI-A and C dust AOD $10\ \mu\text{m}$ daily averages (morning and evening data
 combined) for selected large areas, looking for any discontinuity or unexpected feature. Figure 18 shows those for the North
 Atlantic area (latitude 15 to 35° N, longitude 20 to 60° W), the large Saharan desert (latitude 9.5 to 36.5° N, longitude 17.5° W
 780 to 37.5° E) and the large Asia (latitude 20 to 45° N, longitude 70 to 120° E) respectively. All time series (including for different
 areas not shown here) look perfectly reasonable and do not bear any clear discontinuity. The consistency between IASI-A and
 C will be discussed in the next Sect. 3.9.

3.9 IASI-A vs IASI-C

In this section, we will discuss the potential and definite differences between IASI-A and IASI-C MAPIR dust products. First,
 785 it was shown in Sect. 2.3 that there is an issue in the OFL data used for IASI-C, in which the surface altitude was not provided
 in the files for a significant part of the cloudy observations. That results in less retrievals run for IASI-C, most of them being
 cloudy (which would then be excluded by the QC after the retrieval anyway) but some of them are mis-classified dusty cases.
 This leads to a smaller coverage of some events with IASI-C, but it does not seem to lead to an inconsistency between IASI-A
 and IASI-C in general. Fig. 19 illustrates this for one day during the huge Godzilla dust event. One can very clearly observe a
 790 significant number of missing pixels in the IASI-C retrieval results, with respect to IASI-A. This does not occur specifically for
 the highest AODs only, therefore it would seem that it impacts mostly the coverage, and not so much the representativeness of
 the daily AOD. Fig. 20 illustrates this with the 2020 yearly average dust $10\ \mu\text{m}$ AOD. There are some small differences, mostly
 for the Sahara and Australia. This will be discussed and illustrated further with the second point of the current discussion.

Secondly, even though both Metop-A and C are in the mid-morning orbit 9:30 equator crossing time (descending node), this
 795 is only valid during their operational time. Metop-A has started drifting from this reference orbit in August 2016, after its last
 manoeuvre. This drift remains very small until 2018 (equator crossing time is about 9:15 at the end of that year), but starts
 to be significant during the year 2019, with the equator crossing time of about 8:50 at the end of the year, then about 8:25 at




Figure 18. Time series of MAPIR IASI-A and IASI-C dust AOD at 10 μm , averaged daily (combined morning and evening overpasses) for (a) the North Atlantic area, latitude 15 to 35° N, longitude 20 to 60° W, (b) the Sahara area, latitude 9.5 to 36.5° N, longitude 17.5° W to 37.5° E, (c) for Asia, latitude 20 to 45° N, longitude 70 to 120° E.




Figure 19. MAPIR IASI-A (a) and IASI-C (b) dust AOD 10 μm for 19 June 2020, morning and overpass separately.

the end of 2020 and about 8:00 when IASI-A was decommissioned. This may have an impact on the IASI-A versus IASI-C consistency, especially in some dust source area with diurnal cycle. We have analysed timeseries of IASI-A and IASI-C dust
800 10 μm AOD, similarly as for Fig. 18 but zooming on the years with data from both instruments, for different large source




Figure 20. Yearly average of MAPIR IASI-A (a) and IASI-C (b) dust AOD $10\mu\text{m}$ for 2020.




Figure 21. Same as Fig. 18, zoomed on the 2019–2021 years, for (a) the Sahara area, latitude 9.5 to 36.5° N, longitude 17.5° W to 37.5° E, and (b) the Australian area, latitude 35 to 15° S, longitude 115 to 150° E.

(Sahara, Taklamakan, Gobi, Middle East, Australia) or transport area (North Atlantic or Indian oceans). This analysis was done for morning observations, evening observations, or both combined. From all this, the only area where there is a clear difference between IASI-A and IASI-C are the Sahara and Australia (Fig. 21). In both cases the differences mostly arise from the morning observations, which highlights the plausible cause of the time difference and the morning emission mechanisms. Indeed, part of the dust emitted right after sunrise (6:00–7:00 in the Tropics) and observed with IASI-A could have settled down when IASI-C passes, explaining slightly lower dust AOD with the latter.

Considering the existence of these shifts in the time series, which seem to be due to the Metop-A drift, we recommend for long term analyses to use IASI-A only until IASI-C data comes up in October 2019. At that moment there is already about 30 minutes time difference between both satellites, but the dust data seems to be reasonably continuous.



810 4 Conclusions

Mineral dust is a dominant tropospheric aerosol, with impacts on climate, health and many socio-economical spheres. It is therefore important to monitor it both on the long term and in near-real-time. The Mineral Aerosol Profiling from Infrared Radiances (MAPIR v5.1) algorithm and its application to IASI observations since their beginning in 2007 provide an important contribution to this monitoring.

815 In this manuscript, the MAPIR version 5.1 was presented in detail, clearly highlighting the improvements and changes since the last published version 4.1. MAPIR v5.1 retrieves dust aerosol concentration profiles from ground to 10 km altitude (with respect to sea level), in 1 km thick layers, from which the integrated AOD is obtained together with a mean dust altitude. MAPIR is based on the optimal estimation method with Levenberg-Marquardt optimisation, and uses RTTOV for all radiative transfer computations, with the discrete ordinate method for scattering. The algorithm was used to produce a fully consistent
820 time series of IASI-A dust profile data (July 2007 to September 2021), using the reprocessed IASI spectra and atmospheric profiles, and a IASI-C dust profile time series (since October 2019) as consistent as possible with the IASI-A data. The data product also contains the uncertainties, both as the full profile covariance matrix and as AOD uncertainty. Finally, averaging kernels and a quality flag are also provided.

The retrieval shows a very good capability to retrieve dust aerosols distributions, correctly reproducing seasonal cycles at
825 all analysed locations, and able to detect unusual events. The vertically integrated AOD, converted from 10 μm to 550 nm, was compared against AERONET sun/moon-photometer observations, showing overall a very low median bias of about 0.01, and a good correlation of almost 0.8 when comparing to solar data, and 0.7 for lunar comparisons. The MAPIR bias however varies with the 550 nm AOD (positive bias for low AOD, decreasing towards a negative bias for high AOD) in a quasi linear relationship that suggests the underestimation of the current 10 μm to 550 nm conversion. This conversion and also the chal-
830 lenges of the comparison of thermal infrared based retrievals with visible observations are discussed in detail in the manuscript, highlighting the need for reference data directly in the thermal infrared and the fact that the absolute thermal infrared retrieved AOD can currently not be validated. On the other hand, the good correlation between the compared data sets shows the good capabilities of MAPIR to reproduce the variability of the dust AOD, and does not depend on the absolute AOD values. The low AOD positive bias arises from the low sensitivity in the presence of only small amounts of aerosols, rendering the deviation
835 from the a priori very costly for the retrieval. This bias is also seen at 10 μm : the minimal dust AOD in absence of dust events is about 0.06.

In addition to the good capabilities to reproduce the total column dust load, MAPIR also shows good capabilities to provide vertical information. The information content is however relatively low (maximum about 2.5 degrees of freedom, when the dust load is relatively high and the thermal contrast with the surface is clear), and the 10 retrieved layers are not at all independent.
840 Examples of comparisons with CALIOP show the good capability of MAPIR to retrieve the dust in the correct vertical range(s) even when multiple layers are present. A short demonstration exercise of profile comparisons between MAPIR/IASI and EARLINET lidar data at the Limassol station also highlights the good potential of MAPIR for providing vertical information. This exercise will be expanded and will be the subject of a separate publication.



The mean altitude obtained from the profiles also shows to be reasonable. It is provided only when the sensitivity is acceptable (1.25 degrees of freedom, 0.05 AOD at 10 μm). When the sensitivity is too low, the middle of the vertical range is mostly obtained as mean altitude, which is not physically meaningful. Those cases are mostly filtered out (and the mean altitude not provided), but some still pass the filters and appear as artefacts in the data. This effect stands out when averaging for long time periods, in areas where there is typically only rarely dust.

Finally, the MAPIR data shows long-term stability along both IASI-A and C time series, and a very good potential for merging if taking into account the orbital drift of IASI-A. Indeed, we show that for merging the best moment to switch from IASI-A to IASI-C is at the start of the IASI-C data series, reducing to the minimum the time difference between both instruments. This is especially important close to source areas, where a time difference between instruments also means a potentially large difference in the observed scene. In the current IASI-C time series, some observations are rejected from the data because they lack the surface altitude information. This occurs mostly for cloudy cases, but also for some dusty cases. That issue will be solved in the next processing, using another source for the surface altitude. Our analysis shows that it does not seem to impact long-term analyses, or large area averages.

The cloud filtering remains a challenging step. In the current version, no filtering is done prior to the retrieval, to avoid the removal of some dust events (which occurs with every currently existing cloud flag, as far as we are aware and have analysed). The cloud filtering forms part of the post-retrieval quality control, and leads to much better dust/cloud discrimination than any attempted pre-filtering. However, it is obviously still not perfect and it happens that a thin cloud is kept in the dust data, as an artificial relatively large AOD.

MAPIR v5.1 presented here was used to produce full data time series, which are analysed in this manuscript, but is also in preparation for delivering near-real-time (NRT) observations through the EUMETSAT AC-SAF and the EumetCast system. The two data streams will be maintained in parallel, the "climate" version aiming at the most consistent long-term data series, and the "NRT" version aiming at the best data at production time, for use in forecast models.

Looking towards the future, the MAPIR algorithm will be implemented on data from the newly launched IASI-NG (new generation) instrument onboard Metop-SG (second generation) platform. This is expected to be relatively straightforward, given the similar instrument properties and spectral range. No particular improvement is expected for MAPIR retrievals from the increased spectral resolution and reduced spectral noise: the dust aerosol signature is broad. However, the atmospheric temperature profile is expected to be of improved quality with IASI-NG, which will indirectly reflect on the quality of the dust profiles. Another new instrument has been recently launched, with TIR hyperspectral capabilities: the InfraRed Sounder (IRS) onboard Meteosat Third Generation (MTG), a geostationary platform. Application of a MAPIR-like algorithm to that instrument is planned, and bears serious challenges such as the enormous data amount and the large viewing angles. It is however expected to bring extremely useful information with the high temporal resolution, allowing to better follow the diurnal cycle and to provide more regular input to weather and aerosol forecast models.



Data availability. The MAPIR v5.1 climate time series received a DOI (<https://doi.org/10.18758/f7el2zbr>), and is also available through the iasi.aeronomie.be website, with data completion at least twice per year. Furthermore, the data is produced as level 3 daily and monthly files, containing only the dust AOD and mean altitude. This level 3 data is distributed through the Copernicus Climate Data Store (cds.climate.copernicus.eu, doi: <https://doi.org/10.24381/cds.239d815c>, together with a series of other aerosol data from different satellite instruments.

880 Appendix A: List of used AERONET stations and correlation coefficient with MAPIR IASI

Table A1: AERONET SDA coarse AOD 550nm vs MAPIR IASI dust AOD converted to 550nm correlations by station (sorted by latitude, North to South)

Station	Lat	Lon	Alt (m)	$R_{\text{sol-A}}$	$R_{\text{sol-C}}$	$R_{\text{lun-A}}$	$R_{\text{lun-C}}$
Beijing	39.98	116.38	92	0.87			
Beijing CAMS	39.93	116.32	106	0.81	0.90		
XiangHe	39.75	116.96	36	0.86	0.87		
Dushanbe	38.55	68.86	821	0.66	0.71		
Tunis Carthage	36.84	10.20	10	0.87			
Tizi Ouzou	36.70	4.06	133	0.89			
IMS METU ERDEMLI	36.56	34.26	3	0.73	0.83		
Blida	36.51	2.88	230	0.89			
Gozo	36.03	14.26	111	0.75	0.71		
SACOL	35.95	104.14	1965.8	0.86			
Ben Salem	35.55	9.91	130	0.89	0.92		
Lampedusa	35.52	12.63	45	0.81	0.82		0.75
CUT TEPAK	34.67	33.04	22	0.73	0.57		
Oujda	34.65	-1.90	620	0.91			
Medenine IRA	33.50	10.64	33.5	0.51	0.49		
Migal	33.24	35.58	200	0.77	0.83		
Nes Ziona	31.92	34.79	40	0.76			
Weizmann Institute	31.91	34.81	73	0.67	0.56		
Saada	31.63	-8.16	420	0.87	0.91		
Lahore	31.48	74.26	209	0.77	0.80		
SEDE BOKER	30.86	34.78	480	0.76	0.79		
Cairo EMA 2	30.08	31.29	70	0.57	0.69		
Eilat	29.50	34.92	15	0.60	0.72		
Kuwait University	29.32	47.97	42	0.61	0.53	0.64	
Shagaya Park	29.21	47.06	242	0.54		0.69	

Continued on next page



Station	Lat	Lon	Alt (m)	$R_{\text{sol-A}}$	$R_{\text{sol-C}}$	$R_{\text{lun-A}}$	$R_{\text{lun-C}}$
IIT Delhi	28.55	77.19	15		0.87		
La Laguna	28.48	-16.32	568	0.85	0.78		
Santa Cruz Tenerife	28.47	-16.25	52	0.81	0.68	0.58	0.70
Gual Pahari	28.43	77.15	250	0.89			
Amity Univ Gurgaon	28.32	76.92	285	0.84	0.86		
PLOCAN Tower	28.04	-15.39	12		0.76		
Tenerife	28.03	-16.63	52	0.77	0.69	0.56	0.79
El Farafra	27.06	27.99	92	0.52			
Jaipur	26.91	75.81	450	0.85			
Kanpur	26.51	80.23	123	0.80	0.81	0.59	
Qena SVU	26.20	32.75	75.7	0.68			
Gandhi College	25.87	84.13	60	0.80			
Dhadnah	25.51	56.32	81	0.74			
Abu Al Bukhoosh	25.50	53.15	24.5	0.73			
Karachi	24.95	67.14	49	0.85	0.57		
Solar Village	24.91	46.40	764	0.85			
DEWA ResearchCentre	24.77	55.37	86.8	0.65			
Masdar Institute	24.44	54.62	4	0.72			
Mussafa	24.37	54.47	10	0.80			
Mezaira	23.10	53.75	201	0.66			
University of Nizwa	22.91	57.67	564		0.70		0.59
Tamanrasset INM	22.79	5.53	1377	0.78	0.80		
KAUST Campus	22.30	39.10	11.2	0.72	0.69		
Hada El Sham	21.80	39.73	254	0.73			
Cape San Juan	18.38	-65.62	15	0.85	0.75		
NEON GUAN	17.97	-66.87	128	0.82	0.77		
La Parguera	17.97	-67.05	12.4	0.88	0.86		
Mindelo OSCM	16.88	-25.00	10		0.79		
Calhau	16.86	-24.87	40	0.83			
Capo Verde	16.73	-22.94	60	0.84	0.78		
Guadeloup	16.22	-61.53	39	0.83			
Dakar Belair	14.70	-17.43	15	0.73	0.76	0.64	0.53
Manila Observatory	14.64	121.08	63	0.43			
Dakar	14.39	-16.96	21	0.77		0.51	
Silpakorn Univ	13.82	100.04	72	0.69			
Zinder Airport	13.78	8.99	456	0.26			

Continued on next page



Station	Lat	Lon	Alt (m)	$R_{\text{sol-A}}$	$R_{\text{sol-C}}$	$R_{\text{lun-A}}$	$R_{\text{lun-C}}$
IER Cinzana	13.28	-5.93	285	0.70	0.73		
Ragged Point	13.17	-59.43	40	0.84	0.75		0.69
Bac Lieu	9.28	105.73	10	0.67			
Ilorin	8.48	4.67	400	0.75	0.75	0.48	
Songkhla Met Sta	7.18	100.60	15	0.42			
MCO Hanimaadhoo	6.78	73.18	13	0.63			
LAMTO STATION	6.22	-5.03	105		0.80		
Koforidua ANUC	6.11	-0.30	205	0.80			
ICIPE Mbita	-0.43	34.21	1152	0.52			
San Marco Platform	-2.94	40.21	20		0.64		
ASI Malindi	-3.00	40.19	12	0.46	0.58		0.41
Ascension Island	-7.98	-14.41	30	0.43	0.27		
Arica	-18.47	-70.31	25	0.37			

Author contributions. SV performed the algorithm work and most of data analysis and evaluation, and is the main writer of the paper. CB performed the comparisons with the EARLINET lidar data and wrote the corresponding sections. MEK provided guidance for the validation in general and a thorough review of the whole manuscript. SK contributed to the AERONET validation setup and discussion. MDM participated in all scientific discussions, providing guidance along the project, properly secured the funding and thoroughly reviewed the manuscript.

Competing interests. At least one of the (co-)authors serves as editor for the special issue to which this paper belongs.

Acknowledgements. Funding for MAPIR algorithm development has been provided (including historical first development) by Belspo Supplem Researcher grant nos. WE/35/Q07, R07 and S07, European Space Agency Climate Change Initiative (ESA CCI) aerosols phase 2 (2013-2016) grant 4000109874/14/I-NB and Belspo / ESA PRODEX IASI.flow (2016-2020) PEA 4000111402 and HIRS (2021-2026) PEAs 4000134591 and 4000144088.

Additional funding for current version data processing and delivery to Copernicus Climate Change Services was provided by European Center for Medium-Range Weather Forecast (ECMWF) Copernicus Climate Change Services 2 (C3S2) 312a and 313a Lot2 Atmospheric Composition.

The authors thank the EUMETSAT for operating the IASI instrument and providing its data, including consistent reprocessing. The authors thank the AERONET PI(s) and Co-I(s) and their staff for establishing and maintaining the 74 sites used in this investigation. The authors also thank the CALIOP team at National Aeronautics and Space Administration (NASA) for the instrument exploitation and data provision. We also thank the LIVAS team for provision of the CALIOP pure dust profile data, and the EUMETSAT Numerical Weather



Prediction (NWP) SAF for providing the RTTOV tool. The authors acknowledge EARLINET for providing aerosol lidar profiles available from the EARLINET database.

900 SV thanks James Hocking for useful discussions regarding correct configuration and use of RTTOV. SK and SV acknowledges Harmonia Cost Action CA21119 supported by the European Cooperation in Science and Technology (<https://harmonia-cost.eu/>).

Part of the results presented in this work have been produced using the Aristotle University of Thessaloniki (AUTH) High Performance Computing Infrastructure and Resources. The authors would like to acknowledge the support provided by the Digital Governance Unit of the Aristotle University of Thessaloniki (AUTH) throughout the progress of this research work.



905 References

- Adebisi, A., Kok, J. F., Murray, B. J., Ryder, C. L., Stuut, J.-B. W., Kahn, R. A., Knippertz, P., Formenti, P., Mahowald, N. M., Pérez García-Pando, C., Klose, M., Ansmann, A., Samset, B. H., Ito, A., Balkanski, Y., Di Biagio, C., Romanias, M. N., Huang, Y., and Meng, J.: A review of coarse mineral dust in the Earth system, *Aeolian Research*, 60, 100 849, <https://doi.org/10.1016/j.aeolia.2022.100849>, 2023.
- Amiridis, V., Wandinger, U., Marinou, E., Giannakaki, E., Tsekeri, A., Basart, S., Kazadzis, S., Gkikas, A., Taylor, M., Baldasano, J., and Ansmann, A.: Optimizing CALIPSO Saharan dust retrievals, *Atmospheric Chemistry and Physics*, 13, 12 089–12 106, <https://doi.org/10.5194/acp-13-12089-2013>, 2013.
- Amiridis, V., Marinou, E., Tsekeri, A., Wandinger, U., Schwarz, A., Giannakaki, E., Mamouri, R., Kokkalis, P., Biniotoglou, I., Solomos, S., Herekakis, T., Kazadzis, S., Gerasopoulos, E., Proestakis, E., Kottas, M., Balis, D., Papayannis, A., Kontoes, C., Kourtidis, K., Pappagiannopoulos, N., Mona, L., Pappalardo, G., Le Rille, O., and Ansmann, A.: LIVAS: a 3-D multi-wavelength aerosol/cloud database based on CALIPSO and EARLINET, *Atmospheric Chemistry and Physics*, 15, 7127–7153, <https://doi.org/10.5194/acp-15-7127-2015>, 2015.
- Ansmann, A., Mamouri, R.-E., Hofer, J., Baars, H., Althausen, D., and Abdullaev, S. F.: Dust mass, cloud condensation nuclei, and ice-nucleating particle profiling with polarization lidar: updated POLIPHON conversion factors from global AERONET analysis, *Atmospheric Measurement Techniques*, 12, 4849–4865, <https://doi.org/10.5194/amt-12-4849-2019>, 2019.
- Bernard, E., Moulin, C., Ramon, D., Jolivet, D., Riedi, J., and Nicolas, J.-M.: Description and validation of an AOT product over land at the 0.6 μm channel of the SEVIRI sensor onboard MSG, *Atmospheric Measurement Techniques*, 4, 2543–2565, <https://doi.org/10.5194/amt-4-2543-2011>, 2011.
- Callewaert, S., Vandenbussche, S., Kumps, N., Kylling, A., Shang, X., Komppula, M., Goloub, P., and Mazière, M. D.: The Mineral Aerosol Profiling from Infrared Radiances (MAPIR) algorithm: version 4.1 description and evaluation, *Atmos. Meas. Tech.*, p. 26, <https://doi.org/10.5194/amt-12-3673-2019>, 2019.
- Capelle, V., Chédin, A., Pondrom, M., Crevoisier, C., Armante, R., Crepeau, L., and Scott, N.: Infrared dust aerosol optical depth retrieved daily from IASI and comparison with AERONET over the period 2007–2016, *Remote Sensing of Environment*, 206, 15–32, <https://doi.org/10.1016/j.rse.2017.12.008>, 2018.
- Ceccherini, S. and Ridolfi, M.: Technical Note: Variance-covariance matrix and averaging kernels for the Levenberg-Marquardt solution of the retrieval of atmospheric vertical profiles, *Atmos. Chem. Phys.*, p. 9, <https://doi.org/10.5194/acp-10-3131-2010>, 2010.
- Choobari, O. A., Zawar-Reza, P., and Sturman, A.: The global distribution of mineral dust and its impacts on the climate system: A review, *Atmospheric Research*, 138, 152–165, <https://doi.org/10.1016/j.atmosres.2013.11.007>, 2014.
- Clarisse, L., R'Honi, Y., Coheur, P.-F., Hurtmans, D., and Clerbaux, C.: Thermal infrared nadir observations of 24 atmospheric gases, *Geophysical Research Letters*, 38, n/a–n/a, <https://doi.org/10.1029/2011GL047271>, 2011.
- Clarisse, L., Clerbaux, C., Franco, B., Hadji-Lazaro, J., Whitburn, S., Kopp, A. K., Hurtmans, D., and Coheur, P.-F.: A Decadal Data Set of Global Atmospheric Dust Retrieved From IASI Satellite Measurements, *Journal of Geophysical Research: Atmospheres*, 124, 1618–1647, <https://doi.org/10.1029/2018JD029701>, 2019.
- Clerbaux, C., Boynard, A., Clarisse, L., George, M., Hadji-Lazaro, J., Herbin, H., Hurtmans, D., Pommier, M., Razavi, A., Turquety, S., and Wespes, C.: Monitoring of atmospheric composition using the thermal infrared IASI/MetOp sounder, *Atmos. Chem. Phys.*, p. 14, 2009.
- CNES: IASI-NG, <https://cnes.fr/projets/iasi-ng>, accessed 2026-01-20, 2026.



- 940 D'Amico, G., Amodeo, A., Baars, H., Biniotoglou, I., Freudenthaler, V., Mattis, I., Wandinger, U., and Pappalardo, G.: EAR-LINET Single Calculus Chain – overview on methodology and strategy, *Atmospheric Measurement Techniques*, 8, 4891–4916, <https://doi.org/10.5194/amt-8-4891-2015>, 2015.
- De Rosa, B., Mytilinaios, M., Amodeo, A., Colangelo, C., D'Amico, G., Demma, C., Gandolfi, I., Giunta, A., Gumà-Claramunt, P., Laurita, T., Lolli, S., Papagiannopoulos, N., Papanikolaou, C.-A., Rosoldi, M., Summa, D., and Mona, L.: Observations of Saharan Dust Intrusions
945 over Potenza, Southern Italy, During 13 Years of Lidar Measurements: Seasonal Variability of Optical Properties and Radiative Impact, *Remote Sensing*, 17, <https://doi.org/10.3390/rs17030453>, 2025.
- De Wachter, E., Kumps, N., Vandaele, A. C., Langerock, B., and De Mazière, M.: Retrieval and validation of MetOp/IASI methane, *Atmospheric Measurement Techniques*, 10, 4623–4638, <https://doi.org/10.5194/amt-10-4623-2017>, 2017.
- Di Biagio, C., Formenti, P., Balkanski, Y., Caponi, L., Cazaunau, M., Pangui, E., Journet, E., Nowak, S., Caquineau, S., Andreae, M. O.,
950 Kandler, K., Saeed, T., Piketh, S., Seibert, D., Williams, E., and Doussin, J.-F.: Global scale variability of the mineral dust long-wave refractive index: a new dataset of in situ measurements for climate modeling and remote sensing, *Atmospheric Chemistry and Physics*, 17, 1901–1929, <https://doi.org/10.5194/acp-17-1901-2017>, 2017.
- Di Biagio, C., Formenti, P., Balkanski, Y., Caponi, L., Cazaunau, M., Pangui, E., Journet, E., Nowak, S., Andreae, M. O., Kandler, K.,
Saeed, T., Piketh, S., Seibert, D., Williams, E., and Doussin, J.-F.: Complex refractive indices and single-scattering albedo of global dust
955 aerosols in the shortwave spectrum and relationship to size and iron content, *Atmospheric Chemistry and Physics*, 19, 15 503–15 531, <https://doi.org/10.5194/acp-19-15503-2019>, publisher: Copernicus GmbH, 2019.
- Draxler, R. and Hess, G.: An overview of the HYSPLIT₄ modelling system for trajectories, *Australian Meteorological Magazine*, 47, 295–308, 1998.
- Dubovik, O., Holben, B., Eck, T., Smirnov, A., Kaufman, Y., King, M., Tanré, D., and Slutsker, I.: Variability of Absorption and Optical Properties of Key Aerosol Types Observed in Worldwide Locations, *Journal of the Atmospheric Sciences*, 59, 590–608, [https://doi.org/10.1175/1520-0469\(2002\)059<0590:VOAAOP>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<0590:VOAAOP>2.0.CO;2), 2002.
- 960 Dufresne, J.-L., Gautier, C., Ricchiazzi, P., and Fouquart, Y.: Longwave Scattering Effects of Mineral Aerosols, *Journal of the Atmospheric Sciences*, 59, 1959–1966, [https://doi.org/10.1175/1520-0469\(2002\)059<1959:LSEOMA>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<1959:LSEOMA>2.0.CO;2), 2002.
- Eck, T., Holben, b., Reid, J., Dubovik, O., Smirnov, A., Neill, Slutsker, I., and Kinne, S.: Wavelength dependence of the optical depth of biomass burning, urban, and desert dust aerosols, *Journal of Geophysical Research*, 104, 31 333–31 349, <https://doi.org/10.1029/1999JD900923>, 1999.
- Engelmann, R., Kanitz, T., Baars, H., Heese, B., Althausen, D., Skupin, A., Wandinger, U., Komppula, M., Stachlewska, I. S., Amiridis, V.,
965 Marinou, E., Mattis, I., Linné, H., and Ansmann, A.: The automated multiwavelength Raman polarization and water-vapor lidar Polly^{XT}: the neXT generation, *Atmospheric Measurement Techniques*, 9, 1767–1784, <https://doi.org/10.5194/amt-9-1767-2016>, 2016.
- EUMETSAT: IASI Level 2 Product Guide v3E, 2017.
- EUMETSAT: IASI Level 1 Product Guide v5, 2019.
- EUMETSAT: IASI Principal Components Scores Fundamental Data Record Release 1 - Metop-A and -B,
970 https://doi.org/10.15770/EUM_SEC_CLM_0084, 2022a.
- EUMETSAT: IASI All Sky Temperature and Humidity Profiles - Climate Data Record Release 1.1 - Metop-A and -B, https://doi.org/10.15770/EUM_SEC_CLM_0063, 2022b.
- EUMETSAT: CDR IASI Level 1 PCC release 1 Product User Guide | EUMETSAT, 2022a.
- EUMETSAT: CDR IASI Level 2 TS, T, Q release 1.1 Product User Guide – | EUMETSAT, 2022b.



- 975EUMETSAT: IASI Level 1 Principal Component Scores - Metop - Global - collection id EO:EUM:DAT:METOP:IASPCS01, <https://navigator.eumetsat.int/product/EO:EUM:DAT:METOP:IASPCS01>, 2025a.
- EUMETSAT: IASI Combined Sounding Products - Metop - collection id EO:EUM:DAT:METOP:IASND02, <https://navigator.eumetsat.int/product/EO:EUM:DAT:METOP:IASND02>, 2025b.
- Ge, J. M., Huang, J. P., Xu, C. P., Qi, Y. L., and Liu, H. Y.: Characteristics of Taklimakan dust emission and distribution: A satellite
980 and reanalysis field perspective: Taklimakan dust Characteristics, *Journal of Geophysical Research: Atmospheres*, 119, 11,772–11,783, <https://doi.org/10.1002/2014JD022280>, 2014.
- Giles, D. M., Sinyuk, A., Sorokin, M. G., Schafer, J. S., Smirnov, A., Slutsker, I., Eck, T. F., Holben, B. N., Lewis, J. R., Campbell, J. R., Welton, E. J., Korkin, S. V., and Lyapustin, A. I.: Advancements in the Aerosol Robotic Network (AERONET) Version 3 database – automated near-real-time quality control algorithm with improved cloud screening for Sun photometer aerosol optical depth (AOD) measurements,
985 *Atmospheric Measurement Techniques*, 12, 169–209, <https://doi.org/10.5194/amt-12-169-2019>, 2019.
- Gkikas, A., Basart, S., Hatzianastassiou, N., Marinou, E., Amiridis, V., Kazadzis, S., Pey, J., Querol, X., Jorba, O., Gassó, S., and Baldasano, J.: Mediterranean intense desert dust outbreaks and their vertical structure based on remote sensing data, *Atmospheric Chemistry and Physics*, 16, <https://doi.org/10.5194/acp-16-8609-2016>, 2016.
- Gkikas, A., Proestakis, E., Amiridis, V., Kazadzis, S., Di Tomaso, E., Tsekeri, A., Marinou, E., Hatzianastassiou, N., and Pérez García-Pando, C.: ModIs Dust AeroSol (MIDAS): a global fine-resolution dust optical depth data set, *Atmospheric Measurement Techniques*, 14, 309–334, <https://doi.org/10.5194/amt-14-309-2021>, publisher: Copernicus GmbH, 2021.
- Hilton, F., Armante, R., August, T., Barnet, C., Bouchard, A., Camy-Peyret, C., Capelle, V., Clarisse, L., Clerbaux, C., Coheur, P.-F., Collard, A., Crevoisier, C., Dufour, G., Edwards, D., Faijan, F., Fourrié, N., Gambacorta, A., Goldberg, M., Guidard, V., Hurtmans, D., Illingworth, S.,
995 S., Péquignot, E., Peyridieu, S., Phulpin, T., Remedios, J., Schlüssel, P., Serio, C., Strow, L., Stubenrauch, C., Taylor, J., Tobin, D., Wolf, W., and Zhou, D.: Hyperspectral Earth Observation from IASI: Five Years of Accomplishments, *Bulletin of the American Meteorological Society*, 93, 347–370, <https://doi.org/10.1175/BAMS-D-11-00027.1>, 2012.
- Hocking, J., Saunders, R., Geer, A., and Vidot, J.: RTTOV v13 Users Guide, Tech. rep., European Organisation for the Exploitation of Meteorological Satellites Numerical Weather Prediction Satellite Application Facility, 2021.
- 1000Holben, B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, T., Lavenu, F., Jankowiak, I., and Smirnov, A.: AERONET—A Federated Instrument Network and Data Archive for Aerosol Characterization, *Remote Sensing of Environment*, 66, 1–16, [https://doi.org/10.1016/S0034-4257\(98\)00031-5](https://doi.org/10.1016/S0034-4257(98)00031-5), 1998.
- Hu, Z., Huang, J., Zhao, C., Jin, Q., Ma, Y., and Yang, B.: Modeling dust sources, transport, and radiative effects at different altitudes over the Tibetan Plateau, *Atmospheric Chemistry and Physics*, 20, 1507–1529, <https://doi.org/10.5194/acp-20-1507-2020>, publisher: Copernicus
1005 GmbH, 2020.
- Jacquinet-Husson, N., Crepeau, L., Armante, R., Boutammine, C., Chédin, A., Scott, N., Crevoisier, C., Capelle, V., Boone, C., Poulet-Crovisier, N., Barbe, A., Campargue, A., Chris Benner, D., Benilan, Y., Bézard, B., Boudon, V., Brown, L., Coudert, L., Coustenis, A., Dana, V., Devi, V., Fally, S., Fayt, A., Flaud, J.-M., Goldman, A., Herman, M., Harris, G., Jacquemart, D., Jolly, A., Kleiner, I., Kleinböhl, A., Kwabia-Tchana, F., Lavrentieva, N., Lacombe, N., Xu, L.-H., Lyulin, O., Mandin, J.-Y., Maki, A., Mikhailenko, S., Miller, C., Mishina, T., Moazzen-
1010 Ahmadi, N., Müller, H., Nikitin, A., Orphal, J., Perevalov, V., Perrin, A., Petkie, D., Predoi-Cross, A., Rinsland, C., Remedios, J., Rotger, M., Smith, M., Sung, K., Tashkun, S., Tennyson, J., Toth, R., Vandaele, A.-C., and Vander Auwera, J.: The 2009 edition of the GEISA spectro-



- scopic database, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 112, 2395–2445, <https://doi.org/10.1016/j.jqsrt.2011.06.004>, 2011.
- Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, J. J., Boyd, P. W., Duce, R. A., Hunter, K. A.,
1015 Kawahata, H., Kubilay, N., laRoche, J., Liss, P. S., Mahowald, N., Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global Iron
Connections Between Desert Dust, Ocean Biogeochemistry, and Climate, *Science*, 308, 67–71, <https://doi.org/10.1126/science.1105959>,
publisher: American Association for the Advancement of Science, 2005.
- Kandler, K., Benker, N., Bundke, U., Cuevas, E., Ebert, M., Knippertz, P., Rodríguez, S., Schütz, L., and Weinbruch, S.: Chemical composition
and complex refractive index of Saharan Mineral Dust at Izaña, Tenerife (Spain) derived by electron microscopy, *Atmospheric Environment*,
1020 41, 8058–8074, <https://doi.org/10.1016/j.atmosenv.2007.06.047>, 2007.
- Kim, M.-H., Omar, A. H., Tackett, J. L., Vaughan, M. A., Winker, D. M., Hostetler, C. A., Liu, Z., and Hu, Y.: The CALIPSO version 4 automated
aerosol classification and lidar ratio selection algorithm, *Atmospheric Measurement Techniques*, 11, 6107–6135, <https://doi.org/10.5194/amt-11-6107-2018>, 2018.
- Klüser, L., Banks, J., Martynenko, D., Bergemann, C., Brindley, H., and Holzer-Popp, T.: Information content of space-borne
1025 hyperspectral infrared observations with respect to mineral dust properties, *Remote Sensing of Environment*, 156, 294–309,
<https://doi.org/10.1016/j.rse.2014.09.036>, 2015.
- Koepke, P., Gasteiger, J., and Hess, M.: Technical Note: Optical properties of desert aerosol with non-spherical mineral particles: data incorpo-
rated to OPAC, *Atmospheric Chemistry and Physics*, 15, 5947–5956, <https://doi.org/10.5194/acp-15-5947-2015>, 2015.
- Levenberg, K.: A Method for the Solution of Certain Non-Linear Problems in Least Squares, *Quarterly of Applied Mathematics*, 2, 164–168,
1030 1944.
- Li, J., Carlson, B. E., Yung, Y. L., Lv, D., Hansen, J., Penner, J. E., Liao, H., Ramaswamy, V., Kahn, R. A., Zhang, P., Dubovik, O., Ding, A.,
Lacis, A. A., Zhang, L., and Dong, Y.: Scattering and absorbing aerosols in the climate system, *Nature Reviews Earth & Environment*, 3,
363–379, <https://doi.org/10.1038/s43017-022-00296-7>, 2022.
- Li, L., Mahowald, N. M., Miller, R. L., Pérez García-Pando, C., Klose, M., Hamilton, D. S., Gonçalves Ageitos, M., Ginoux, P., Balkanski, Y.,
1035 Green, R. O., Kalashnikova, O., Kok, J. F., Obiso, V., Paynter, D., and Thompson, D. R.: Quantifying the range of the dust direct radiative
effect due to source mineralogy uncertainty, *Atmospheric Chemistry and Physics*, 21, 3973–4005, <https://doi.org/10.5194/acp-21-3973-2021>,
2021.
- Loveless, M., Borbas, E. E., Knuteson, R., Cawse-Nicholson, K., Hulley, G., and Hook, S.: Climatology of the Combined ASTER MODIS
Emissivity over Land (CAMEL) Version 2, *Remote Sensing*, 13, 111, <https://doi.org/10.3390/rs13010111>, 2021.
- 1040 Maes, K., Vandenbussche, S., Klüser, L., Kumps, N., and de Mazière, M.: Vertical Profiling of Volcanic Ash from the 2011 Puyehue Cordón
Caulle Eruption Using IASI, *Remote Sensing*, 8, 103, <https://doi.org/10.3390/rs8020103>, 2016.
- Mahowald, N., Albani, S., Kok, J. F., Engelstaeder, S., Scanza, R., Ward, D. S., and Flanner, M. G.: The size distribution of desert dust aerosols
and its impact on the Earth system, *Aeolian Research*, 15, 53–71, <https://doi.org/10.1016/j.aeolia.2013.09.002>, 2014.
- Mamouri, R. E. and Ansmann, A.: Fine and coarse dust separation with polarization lidar, *Atmospheric Measurement Techniques*, 7, 3717–
1045 3735, <https://doi.org/10.5194/amt-7-3717-2014>, 2014.
- Mamouri, R.-E., Ansmann, A., Nisantzi, A., Solomos, S., Kallos, G., and Hadjimitsis, D. G.: Extreme dust storm over the eastern Mediterranean
in September 2015: satellite, lidar, and surface observations in the Cyprus region, *Atmospheric Chemistry and Physics*, 16, 13 711–13 724,
<https://doi.org/10.5194/acp-16-13711-2016>, 2016.



- Marquardt, D. W.: An Algorithm for Least-Squares Estimation of Nonlinear Parameters, *Journal of the Society for Industrial and Applied Mathematics*, 11, 431–441, <https://doi.org/10.1137/0111030>, 1963.
- Michailidis, K., Koukouli, M.-E., Balis, D., Veefkind, J. P., de Graaf, M., Mona, L., Papagianopoulos, N., Pappalardo, G., Tsikoudi, I., Amiridis, V., Marinou, E., Gialitaki, A., Mamouri, R.-E., Nisantzi, A., Bortoli, D., João Costa, M., Salgueiro, V., Papayannis, A., Mylonaki, M., Alados-Arboledas, L., Romano, S., Perrone, M. R., and Baars, H.: Validation of the TROPOMI/S5P aerosol layer height using EARLINET lidars, *Atmospheric Chemistry and Physics*, 23, 1919–1940, <https://doi.org/10.5194/acp-23-1919-2023>, 2023.
- Middleton, N.: Desert dust hazards: A global review, *Aeolian Research*, 24, 53–63, <https://doi.org/10.1016/j.aeolia.2016.12.001>, 2017.
- Mielonen, T., Arola, A., Komppula, M., Kukkonen, J., Koskinen, J., de Leeuw, G., and Lehtinen, K.: Comparison of CALIOP Level 2 aerosol subtypes to aerosol types derived from AERONET inversion data, *Geophysical Research Letters*, 38, <https://doi.org/10.1029/2009GL039609>, 2009.
- Mona, L., Amiridis, V., Cuevas, E., Gkikas, A., Trippetta, S., Vandenbussche, S., Benedetti, A., Dagsson-Waldhauserova, P., Formenti, P., Haefele, A., Kazadzis, S., Knippertz, P., Laurent, B., Madonna, F., Nickovic, S., Papagiannopoulos, N., Pappalardo, G., García-Pando, C. P., Popp, T., Rodríguez, S., Sealy, A., Sugimoto, N., Terradellas, E., Vimic, A. V., Weinzierl, B., and Basart, S.: Observing Mineral Dust in Northern Africa, the Middle East, and Europe: Current Capabilities and Challenges ahead for the Development of Dust Services, *Bulletin of the American Meteorological Society*, 104, E2223–E2264, <https://doi.org/10.1175/BAMS-D-23-0005.1>, 2023.
- Monteiro, A., Basart, S., Kazadzis, S., Votsis, A., Gkikas, A., Vandenbussche, S., Tobias, A., Gama, C., García-Pando, C. P., Terradellas, E., Notas, G., Middleton, N., Kushta, J., Amiridis, V., Lagouvardos, K., Kosmopoulos, P., Kotroni, V., Kanakidou, M., Mihalopoulos, N., Kalivitis, N., Dagsson-Waldhauserová, P., El-Askary, H., Sievers, K., Giannaros, T., Mona, L., Hirtl, M., Skomorowski, P., Virtanen, T. H., Christoudias, T., Di Mauro, B., Trippetta, S., Kutuzov, S., Meinander, O., and Nickovic, S.: Multi-sectoral impact assessment of an extreme African dust episode in the Eastern Mediterranean in March 2018, *Science of The Total Environment*, 843, 156 861, <https://doi.org/10.1016/j.scitotenv.2022.156861>, 2022.
- Newman, S. M., Smith, J. A., Glew, M. D., Rogers, S. M., and Taylor, J. P.: Temperature and salinity dependence of sea surface emissivity in the thermal infrared, *Quarterly Journal of the Royal Meteorological Society*, 131, 2539–2557, <https://doi.org/10.1256/qj.04.150>, 2005.
- O’Neill, N. T., Eck, T. F., Smirnov, A., Holben, B. N., and Thulasiraman, S.: Spectral discrimination of coarse and fine mode optical depth, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD002975>, 2003.
- Pappalardo, G., Amodeo, A., Apituley, A., Comeron, A., Freudenthaler, V., Linné, H., Ansmann, A., Bösenberg, J., D’Amico, G., Mattis, I., Mona, L., Wandinger, U., Amiridis, V., Alados-Arboledas, L., Nicolae, D., and Wiegner, M.: EARLINET: towards an advanced sustainable European aerosol lidar network, *Atmospheric Measurement Techniques*, 7, 2389–2409, <https://doi.org/10.5194/amt-7-2389-2014>, 2014.
- Proestakis, E., Gkikas, A., Georgiou, T., Kampouri, A., Drakaki, E., Ryder, C. L., Marengo, F., Marinou, E., and Amiridis, V.: A near-global multiyear climate data record of the fine-mode and coarse-mode components of atmospheric pure dust, *Atmospheric Measurement Techniques*, 17, 3625–3667, <https://doi.org/10.5194/amt-17-3625-2024>, 2024.
- Ridley, D., Heald, C., and Ford, B.: North African dust export and deposition: A satellite and model perspective, *Journal of Geophysical Research*, 117, D02 202, <https://doi.org/10.1029/2011JD016794>, 2012.
- Rodgers, C. D.: Inverse Methods for Atmospheric Sounding: Theory and Practice, vol. 2 of *Series on Atmospheric, Oceanic and Planetary Physics*, WORLD SCIENTIFIC, ISBN 978-981-02-2740-1 978-981-281-371-8, <https://doi.org/10.1142/3171>, 2000.
- Rodgers, C. D. and Connor, B. J.: Intercomparison of remote sounding instruments, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD002299>, 2003.



- Saunders, R., Hocking, J., Turner, E., Rayer, P., Rundle, D., Brunel, P., Vidot, J., Roquet, P., Matricardi, M., Geer, A., Bormann, N., and Lupu, C.: An update on the RTTOV fast radiative transfer model (currently at version 12), *Geoscientific Model Development*, 11, 2717–2737, <https://doi.org/10.5194/gmd-11-2717-2018>, 2018.
- Saunders, R., Hocking, J., Turner, E., Geer, A., Lupu, C., and Vidot, J.: RTTOV-13 SCIENCE AND VALIDATION REPORT, Tech. rep., 1090 EUMETSAT Satellite Application Facility on Numerical Weather Prediction (NWP SAF), https://nwp-saf.eumetsat.int/site/download/documentation/rtm/docs_rtov13/rttov13_svr.pdf, 2020.
- Sinyuk, A., Holben, B. N., Eck, T. F., Giles, D. M., Slutsker, I., Korkin, S., Schafer, J. S., Smirnov, A., Sorokin, M., and Lyapustin, A.: The AERONET Version 3 aerosol retrieval algorithm, associated uncertainties and comparisons to Version 2, *Atmospheric Measurement Techniques*, 13, 3375–3411, <https://doi.org/10.5194/amt-13-3375-2020>, 2020.
- 1095 Sokolik, I. N.: The spectral radiative signature of wind-blown mineral dust: Implications for remote sensing in the thermal IR region, *Geophysical Research Letters*, 29, 7–1–7–4, <https://doi.org/10.1029/2002GL015910>, 2002.
- Song, Q., Zhang, Z., Yu, H., Kato, S., Yang, P., Colarco, P., Remer, L. A., and Ryder, C. L.: Net radiative effects of dust in the tropical North Atlantic based on integrated satellite observations and in situ measurements, *Atmospheric Chemistry and Physics*, 18, 11 303–11 322, <https://doi.org/10.5194/acp-18-11303-2018>, 2018.
- 1100 Song, Q., Zhang, Z., Yu, H., Ginoux, P., and Shen, J.: Global dust optical depth climatology derived from CALIOP and MODIS aerosol retrievals on decadal timescales: regional and interannual variability, *Atmospheric Chemistry and Physics*, 21, <https://doi.org/10.5194/acp-21-13369-2021>, 2021.
- Tesche, M., Ansmann, A., Müller, D., Althausen, D., Engelmann, R., Freudenthaler, V., and Groß, S.: Vertically resolved separation of dust and smoke over Cape Verde using multiwavelength Raman and polarization lidars during Saharan Mineral Dust Experiment 2008, *Journal of Geophysical Research: Atmospheres*, 114, <https://doi.org/10.1029/2009JD011862>, 2009.
- 1105 Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Feichter, H., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Horowitz, L., Huang, P., Isaksen, I., Iversen, T., Kloster, S., Koch, D., Kirkevåg, A., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, O., Stier, P., Takemura, T., and Tie, X.: Analysis and quantification of the diversities of aerosol life cycles within AeroCom, *Atmos. Chem. Phys.*, p. 37, 2006.
- Vandenbussche, S., Langerock, B., Vigouroux, C., Buschmann, M., Deutscher, N. M., Feist, D. G., García, O., Hannigan, J. W., Hase, F., Kivi, R., Kumps, N., Makarova, M., Millet, D. B., Morino, I., Nagahama, T., Notholt, J., Ohyama, H., Ortega, I., Petri, C., Rettinger, M., Schneider, M., Servais, C. P., Sha, M. K., Shiomi, K., Smale, D., Strong, K., Sussmann, R., Té, Y., Velasco, V. A., Vrekoussis, M., Warneke, T., Wells, K. C., Wunch, D., Zhou, M., and Mazière, M. D.: Nitrous Oxide Profiling from Infrared Radiances (NOPIR): Algorithm Description, 1115 Application to 10 Years of IASI Observations and Quality Assessment, p. 30, <https://doi.org/https://doi.org/10.3390/rs14081810>, 2022.
- Voudouri, K., Siomos, N., Michailidis, K., Papagiannopoulos, N., Mona, L., Nicolae, D., and Balis, D.: Comparison of two automated aerosol typing methods and their application to an EARLINET station, *Atmospheric Chemistry and Physics*, 19, 10 961–10 980, <https://doi.org/10.5194/acp-19-10961-2019>, 2019.
- Winker, D. M., Vaughan, M. A., Omar, A., Hu, Y., Powell, K. A., Liu, Z., Hunt, W. H., and Young, S. A.: Overview of the 1120 CALIPSO Mission and CALIOP Data Processing Algorithms, *Journal of Atmospheric and Oceanic Technology*, 26, 2310–2323, <https://doi.org/10.1175/2009JTECHA1281.1>, 2009.
- Winker, D. M., Tackett, J. L., Getzewich, B. J., Liu, Z., Vaughan, M. A., and Rogers, R. R.: The global 3-D distribution of tropospheric aerosols as characterized by CALIOP, *Atmospheric Chemistry and Physics*, 13, 3345–3361, <https://doi.org/10.5194/acp-13-3345-2013>, 2013.



- Xie, X., Liu, X., Che, H., Xie, X., Li, X., Shi, Z., Wang, H., Zhao, T., and Liu, Y.: Radiative feedbacks of dust in snow over eastern Asia in CAM4-BAM, *Atmospheric Chemistry and Physics*, 18, 12 683–12 698, <https://doi.org/10.5194/acp-18-12683-2018>, 2018.
- Zender, C. S., Miller, R. L. L., and Tegen, I.: Quantifying mineral dust mass budgets: Terminology, constraints, and current estimates, *Eos, Transactions American Geophysical Union*, 85, 509–512, <https://doi.org/10.1029/2004EO480002>, 2004.
- Zhang, X., Zhao, L., Tong, D., Wu, G., Dan, M., and Teng, B.: A Systematic Review of Global Desert Dust and Associated Human Health Effects, *Atmosphere*, 7, 158, <https://doi.org/10.3390/atmos7120158>, 2016.
- 1125 Zhou, D. K., Larar, A. M., Liu, X., Smith, W. L., Strow, L. L., Yang, P., Schlusser, P., and Calbet, X.: Global Land Surface Emissivity Retrieved From Satellite Ultraspectral IR Measurements, *IEEE Transactions on Geoscience and Remote Sensing*, 49, 1277–1290, <https://doi.org/10.1109/TGRS.2010.2051036>, 2011.
- 1130