

The ~~i~~mpact of NaOH, CaO, and ~~[Ca²⁺] + [HCO₃⁻]~~ ~~Ca²⁺ + HCO₃⁻~~ ~~a~~ Additions on PIC and POC ~~f~~Formation in Los Angeles Harbor Waters

Rucha P. Wani¹, Devan L. Roper¹, Ria Agrawal², Esther J. Lim¹, Nick E. Rollins¹, and William M. Berelson¹

¹Department of Earth Sciences, University of Southern California, Los Angeles, 90089, United States

²Rosenstiel School of Marine, Atmospheric & Earth Science, University of Miami, Miami, 33149, United States

Correspondence: Rucha P. Wani (rwani@usc.edu)

Abstract

Negative CO₂ emission technologies such as ocean alkalinity enhancement, in tandem with emissions reduction are necessary to keep the climate system below a critical tipping point. While the biogeochemical consequences of alkalinity enhancement remain poorly constrained, field deployment is accelerating in the commercial sector, leaving questions of ecosystem impact in the wake. In this study we conduct alkalinity perturbation experiments to capture the resultant impact to the organic carbon and calcium carbonate pools. We quantify shifts in dissolved/particulate inorganic and organic carbon after the addition of three alkalinity sources – NaOH, CaO, and CaCl₂ + NaHCO₃ (to simulate accelerated weathering of limestone). These experiments are conducted with coastal sea water with enhancements of $\pm \sim 500$ and $\pm \sim 1000$ $\mu\text{mol kg}^{-1}$ alkalinity, and incubated in situ, to elucidate the impact over 0–4 days. ¹³C isotope spikes were added to trace carbon partitioning throughout the ~~experiment~~~~experiment~~, and a few bottle experiments were completely isolated from the light via bottle shading to assess impacts on ~~a majority heterotrophic ecosystem~~~~non-photosynthesizers~~. Despite a wide range in the initial ambient particulate carbon (both organic and inorganic) there was no statistically significant calcium carbonate precipitation or change in the organic carbon partitioning after a perturbation of $\pm \sim 500$ $\mu\text{mol kg}^{-1}$ alkalinity in the form of NaOH, CaO or CaCl₂ + NaHCO₃. An increase of alkalinity by ~ 1000 $\mu\text{mol kg}^{-1}$ after CaO addition resulted in ~~the a~~ statistically significant decrease of particulate organic carbon and dissolved oxygen relative to the control. To understand how a decrease in the particulate organic carbon may impact downstream trophic transfers, future studies should resolve the observed changes in particulate organic carbon by

Formatted: Font: 14 pt, Bold

Formatted: Font: 14 pt

Formatted: Font: 14 pt

Formatted: Font: 14 pt

Formatted: Font: 14 pt

Formatted: Font: 14 pt

Formatted: Font: 14 pt, Bold


characterizing biomass within various particle class sizes with further filtration, microscopy, flow cytometry, and 'omics analyses.

1. Introduction

To mitigate the impact of excess atmospheric carbon dioxide (CO₂), humans need to reduce CO₂ emissions and remove CO₂ from the atmosphere. According to the Intergovernmental Panel on Climate Change, ~~in the 8 years since the Paris Agreement,~~ global efforts to reduce greenhouse gas emissions are still not on track to avoid surpassing the 1.5 °C threshold of temperature increase by the end of the century (de Coninck et al., 2018; (UNFCCC) United Nations Framework Convention on Climate Change, 2022). Since reducing CO₂ emissions may not be sufficient to meet the 2015 Paris Agreement Goals (National Academies of Sciences, 2022), strategies for marine Carbon Dioxide Removal (mCDR) are being investigated to aid in the uptake and storage of CO₂. The Earth has balanced its own carbon storage and acidity on geological timescales (i.e., 100-1000 ka) through mineral weathering and mineral precipitation (Bernier et al., 1983; Bernier and Bernier, 1997; Holland, 1978; Penman et al., 2020; Walker et al., 1981). While there may be more than enough calcium carbonate (CaCO₃) in ocean sediments to neutralize all anthropogenic CO₂ (Archer, 1996; Sulpis et al., 2018), this recovery could take >10,000 years (Zachos et al., 2005) given slow dissolution kinetics and ocean mixing rates. Ocean Alkalinity Enhancement (OAE) is a mCDR method that can potentially speed up the process of oceanic CO₂ uptake for human relevant timescales. A thorough assessment of the scientific background and considerations for various forms of OAE has been presented in Renforth & Henderson (2017). The proposed risks and co-benefits of increasing surface ocean alkalinity based on different deployment strategies have been theorized in Bach et al. (2019).

There are still many technical challenges to overcome before OAE can be scaled to remove gigatons of CO₂ (Eisaman et al., 2023; Renforth, 2025). The impacts of different alkaline feedstocks and concentrations on ocean chemistry and the ecosystem are yet to be completely understood. The successful long-term sequestration of CO₂ relies on many factors – the rate of air-sea gas exchange, the loss of alkalinity to re-precipitation of carbonate minerals or even the loss of an unequilibrated plume as it subducts below the surface (Lunstrum et al., 2026). Increasing alkalinity could change the cycling of inorganic carbon in the ocean by increasing the

likelihood of CaCO_3 precipitation ~~or decreasing natural CaCO_3 dissolution~~ and thereby affecting the associated organic carbon cycling pumps (Bach, 2024; Falkowski et al., 2000; Ridgwell and Zeebe, 2005; Riebesell et al., 2000; Zondervan et al., 2001). Limiting secondary precipitation of CaCO_3 is important for OAE because there is release of CO_2 from the precipitation of CaCO_3 (Eq. 1) – which reduces the efficiency of OAE.



Another potential outcome of OAE is for organic matter to be ballasted to the deep ocean by aggregating on newly formed particulate inorganic carbon (PIC) (Riebesell et al., 2009). The natural global production of particulate organic carbon (POC) is known to be 4-10 times higher than PIC production (Barker et al., 2006); although this ratio is highly variable in time and space (Berelson et al., 2007; Dunne et al., 2007). Renforth & Henderson, (2017) hypothesized that if the POC production remains higher than PIC production even under OAE, then the CO_2 emissions from CaCO_3 production by marine calcifiers could be offset. This highlights the need for us to assess whether enhanced calcification after OAE will contribute to CO_2 production or if it might further sequester organic carbon by enhancing POC production or by aggregating and ballasting POC via enhanced PIC (Bach et al., 2019).

Given the variety of compounds available to increase alkalinity (Renforth and Henderson, 2017), their composition and deployment style will determine the magnitude of change in Omega (Ω = the saturation state) and likelihood of CaCO_3 precipitation. This applies to both the near field impacts at the point of injection and the far field impacts after the alkalinity has been dispersed. Hereafter, we shall reference Ω in terms of $\Omega_{\text{aragonite}}$ as this is the morphotype that precipitates more readily in present-day seawater (Morse et al., 2007). Many studies have observed increased carbonate mineral precipitation, as demonstrated by the loss of alkalinity (TA) and dissolved inorganic carbon (DIC) after the addition of sodium hydroxide (NaOH), quicklime/slaked lime ($\text{CaO}/\text{Ca}(\text{OH})_2$), and even sodium carbonate (Na_2CO_3). Precipitation was observed in both short term (<5 days) (Hartmann et al., 2023; Hashim et al., 2025; Moras et al., 2022; Subhas et al., 2022), and longer term (~20 days) perturbation experiments at a range of mineral saturation states Ω (Moras et al., 2022, 2024; Paul et al., 2024; Ringham et al., 2024; Suitner et al., 2024). Yet, OAE experiments conducted with cultures of coccolithophores have noted no change to calcification rates after additions of ~~bicarbonate and calcium~~ calcium and

Formatted: Subscript

Formatted: Centered

bicarbonate ($[\text{Ca}^{2+}] + [\text{HCO}_3^-] - [\text{Ca}^{2+}] - [\text{HCO}_3^-]$) (Gately et al., 2023). The experimental conditions of observed CaCO_3 precipitation will be discussed to provide further context to the results presented in this study.

The examples of existing perturbation experiments come from natural waters around Spain, North Pacific, North Atlantic gyres, and Australian coastal waters. To this rapidly growing body of research, we present the changes of organic and inorganic carbon inventory and partitioning during 4-day OAE experiments conducted using coastal California waters from Los Angeles (LA) Harbor. The waters offshore from Los Angeles are richly characterized by decades of regional studies (<https://dornsife.usc.edu/spot/publications-and-projects/>, <https://calcofi.org/publications/peer-reviewed-publications/>). Yet waters from within LA Harbor contain blends of offshore ecosystems modified by nearshore processes. In our study, we use three different approaches to elevate $\Omega_{\text{aragonite}}$ by: (i) increasing pH (using two different bases) and (ii) adding $[\text{Ca}^{2+}]$ and $[\text{HCO}_3^-]$ to simulate accelerated weathering of limestone. In constraining the resultant changes to the amount of PIC, POC and DOC production, we conduct a carbon budget analysis that helps define how alkalinity may change the natural cycling of organic and inorganic carbon.

2. ~~Study Area~~Materials & Methods

2.1 Study ~~a~~Area

With the exception of some short-term experiments conducted on Santa Catalina Island at the USC Dornsife Wrigley Marine Science Center (33° 26' 40.52" N, 118° 28' 47.787" W), all other experiments were conducted with water collected from ~1 m ~~water~~-depth in Los Angeles Harbor, at AltaSea Marine Center (33° 43' 33.64" N, 118° 16' 11.999" W). This region exchanges seawater with the Southern California Bight, which is a part of the California Current System. The microbial community structure in the Port of Los Angeles in the summer is dominated by diatoms, pico-eukaryotes, and heterotrophic bacteria. Dinoflagellates, Synechococcus, and photosynthetic nanoplankton are in less abundance. In the fall, the carbon biomass shifts to the following in order of abundance: Bacteria > Picoeukaryotes > Synechococcus > Diatoms ~ ~~Pico~~ ~~N~~nanoplankton > Dinoflagellates; these proportionally decrease in the winter when Diatoms become the dominant species again (Bialonski et al., 2016; Caron et al., 2017; Connell et al., 2017). Regardless of the season, the carbon biomass is higher in the Port compared to the costal

Formatted: Not Superscript/ Subscript

San Pedro Ocean Time series station (~17 km offshore) or in the marine protected area near Santa Catalina Island (~45 km offshore) (Connell et al., 2017).

Alkalinity perturbation experiments were conducted ~~with-in~~ 5L, Thermo Scientific Nalgene PETG bottles (total volume is ~6L with no headspace) suspended 1 m below the water surface off a dock at AltaSea ~~Marine Center~~. Surface seawater conditions (temperature and sunlight exposure) during each experiment are recorded in Table 1. Temperature (°C) and Light Intensity (lux) were measured with HOBO™ Temperature and Light Loggers, and the lux was converted to photosynthetic photon flux density ($\mu\text{mol photons m}^{-2} \text{s}^{-1}$) by multiplying lux with the standard conversion factor of sunlight (0.0185). Salinity of the control waters from all experiments were roughly 35 ± 0.5 ppt; salinity was measured with a refractometer.

Table 1. Ambient ~~s~~Sea ~~s~~Surface ~~e~~Conditions during each ~~e~~Experimental ~~d~~Deployment.

<u>Target Treatment</u>	Start Date	Sea Surface Temperature (°C)	Light ($\mu\text{mol m}^{-2} \text{s}^{-1}$)
CaCl₂ + NaHCO₃ (~ +500 $\mu\text{mol kg}^{-1}$)	06-24-24	16.4 – 22.3	0 – 540
	07-15-24	17.4 – 24.5	0 – 605
NaOH (~ +500 $\mu\text{mol kg}^{-1}$)	06-17-24	17.2* - 21.1*	0 – 640
	07-22-24	15.8 – 20.8	0 – 605
CaO (~ +500 $\mu\text{mol kg}^{-1}$)	11-18-24	13.7* - 14.7*	0 – 400
	01-27-25	13.5 – 20.8	0 – 470
CaO (~ +1000 $\mu\text{mol kg}^{-1}$)	11-04-24	14.1 – 22.01	0 – 620

*Data taken from *NOAA Tides & Currents* from Station #9410840 Santa Monica, CA. All other data measured from the AltaSea dock side water at ~~~~~1 m water depth.

2.2 Alkalinity ~~t~~Types

In this study we induce an alkalinity perturbation that simulates two different methods of OAE at two target enrichments, (1) Base Addition: where a strong base is delivered to the surface of the ocean and is intended to increase oceanic uptake of atmospheric CO₂ by disrupting the pH balance. To simulate this, we use two common bases, NaOH (J. T. Baker, New Jersey. ACS

Formatted: Centered

Formatted Table


Formatted: Centered

Formatted: Centered

Formatted: Centered

Formatted: Centered

Grade, Lot. No.: 99057), and CaO (Sigma-Aldrich, Missouri. Reagent Grade, Batch No.: MKCS6828). The other method considered here is (2) accelerated weathering of limestone (AWL) (Dong et al., 2025; Kheshgi, 1995; Rau and Caldeira, 1999) where atmospheric CO₂ and water are pumped through a limestone reactor to neutralize the CO₂. The outflow water is enriched in the byproducts of this reaction (Eq. 2) – Ca²⁺ and HCO₃⁻ ions. In this method the CO₂ is neutralized before it enters the seawater, as opposed to being neutralized in-situ during air-sea gas exchange with the base addition method. We mimic AWL with the addition of Calcium Chloride (CaCl₂, Sigma-Aldrich, Missouri. Assay ≥ 97%, Batch No.: SLCL2754) and Sodium Bicarbonate (NaHCO₃, Sigma-Aldrich, Missouri. ACS Grade, Batch No.: SLCM3421) with a molar ratio of 1:2 for Ca²⁺:HCO₃⁻ (Eq. 3).



To achieve specific dosages/concentrations, the alkalinity needed to be fully dissolved, thus we avoided introducing any particles into the incubations. Also, aqueous alkalinity delivery is currently considered a favorable delivery method for OAE projects (Cai and Jiao, 2022; Savoie et al., 2025). To compare our results with other published regional studies (Bach et al., 2024; Ferderer et al., 2022; Kousoulas et al., 2025; Subhas et al., 2022) and evaluate impacts above current instrument detection limits (See Section 3.1.1), we chose target enrichments of ~500 and ~1000 μmol kg⁻¹. We created dissolved stock solutions of each alkalinity type (NaOH, CaO, CaCl₂, NaHCO₃) with minimal exposure to the atmosphere. A solid salt or base was precisely weighed, transferred to a pre-weighed volumetric flask, the flask was filled with Milli-Q water, and no headspace remained. After all solids were dissolved, the solution was siphoned through a 0.2 μm GF/F filter into an evacuated gas impermeable foil bag (Supelco, Pennsylvania) for storage. We took subsamples for TA and DIC from the stock foil bags during preparation and while dosing in the field (measurement details in Section 2.5) to back-calculate the final concentrations of the stock solutions. ~~and precisely dosed our incubations according to these measurements to achieve the desired alkalinity enrichments~~ The target dosages, the predicted changes calculated through a dilution equation, and the actual measured changes in TA within bottle replicates (n = 3) are reported in (Table 2) ~~30 mL of 0.94M NaOH (+ 500 μmol kg⁻¹), 30 mL 0.05M CaCl₂ + 0.12M NaHCO₃ (+ 500 μmol kg⁻¹), 100mL of 0.016M (+ 500 μmol kg⁻¹),~~

and 160 mL of 0.018M CaO ($+1000 \mu\text{mol kg}^{-1}$). Alkalinity was measured with an open-system Gran titration method at 25 °C (details in (Lunstrum and Berelson, 2022; Subhas et al., 2015)) and no precipitation was observed in the storage containers (stored at 4 °C until analysis). As the solubility of CaO in water is less than that of the other chemicals used in this study, a larger volume of a weaker CaO stock was required to make the alkalinity additions. Hence, an immediate drop in DIC (by <2%) due to the expected dilution by the stock solution is observed in all CaO experiments.

To increase the sensitivity of our measurements for the carbon budget analysis, we added 1-2mL of a ^{13}C tracer in the form of $\text{NaH}^{13}\text{CO}_3$ (Sigma Aldrich, Missouri. CAS No.: 87081-58-1, Assay = 99%, Lot No.: MBBC0208V) to all controls and treatments in five experiments: 07-15-24, 07-22-24, 11-04-24, 11-18-24, and 01-27-25. The tracer solution ($\sim 0.015\text{M}$) was prepared similarly to the stock solutions discussed above. The tracer only changed the alkalinity by $<52 \mu\text{mol kg}^{-1}$ but increased the $\delta^{13}\text{DIC}$ signal to 80-200 ‰ (precise start values are included in the associated dataset (Wani et al., 2026)).

Table 2. Summary of alkalinity additions

Target Treatment	Start Date	Stock Alkalinity (mol kg ⁻¹) and Dose (mL)	Predicted ΔTA ($\mu\text{mol kg}^{-1}$) ($\pm$ stdev)	Measured ΔTA ($\mu\text{mol kg}^{-1}$) ($\pm$ error propagated)
CaCl₂ + NaHCO₃ ($\sim 500 \mu\text{mol kg}^{-1}$)	06-24-24	0.1 & 30 mL	474 $\pm$ 9	486 $\pm$ 11
	07-15-24	0.1 & 28 mL	445 $\pm$ 2	441 $\pm$ 10
NaOH ($\sim 500 \mu\text{mol kg}^{-1}$)	06-17-24	0.107 & 30 mL	524 $\pm$ 1	520 $\pm$ 4
	07-22-24	0.094 & 35 mL	532 $\pm$ 9	527 $\pm$ 11
CaO ($\sim 500 \mu\text{mol kg}^{-1}$)	11-18-24	0.016 & 100 mL	509 $\pm$ 0	533 $\pm$ 18
	01-27-25	0.019 & 100 mL	616 $\pm$ 6	612 $\pm$ 13
CaO ($\sim 1000 \mu\text{mol kg}^{-1}$)	11-04-24	0.018 & 160 mL	919 $\pm$ 2	923 $\pm$ 12

Predicted error comes from the standard deviation of the seawater volume in n = 3 bottles. Measured error comes from the standard deviation of the replicate bottles and the error was propagated after values were corrected with a certified reference material (Dickson Seawater).

2.42.3 Experimental Setup

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Not Bold

Formatted Table

Formatted: Font: Not Bold

Formatted: Font: 11 pt

Formatted: Font: 11 pt

Formatted: Font: 11 pt

Formatted: Font: 11 pt

Formatted: Font: 10 pt, Font color: Auto

Formatted: Font: 10 pt, Font color: Auto


Formatted: Font: 10 pt

Formatted: Line spacing: single

There is long standing discourse on whether bottle experiments mimic natural conditions enough to be useful to answer questions of “ecosystem impact” (Daehler and Strong, 1996; Fraser and Keddy, 1997; Giesy, 1980; Ives et al., 1996; Lawton et al., 1993; Stachowski-Haberkorn et al., 2008). There are many experimental design factors that can influence the community assemblage, such as: material used for the structure, microcosm volume, filling mechanism, mixing, replication, carbonate chemistry, exclusion of larger organisms, and wall surface growth (Riebesell et al., 2010). In perturbation experiments conducted with small volume microcosms (< 10L) it is recommended to limit the timescale of study to days-weeks if the goal is to understand the impact of the perturbation on microbial or plankton assemblages (Parsons, 1982; Riebesell et al., 2010). The purpose is to limit unwanted effects such as biofouling on the walls of the enclosure, nutrient limitation, excess grazing or changes due to lack of predators- these are commonly referred to as “bottle effects”. Our aim was to compare multiple alkalinity sources; thus we opted for a replication method that incorporated sacrificial bottle timepoints over the span of four days (Iglesias-Rodríguez et al. 2023). We maintained a closed system experiment (with no headspace) to constrain the carbon budget by eliminating gas exchange as a potential variable for this study. This can be an analog for water that may temporarily lose contact with the atmosphere, e.g. when waters mix deeper into surface ocean or for times when gas exchange is minimal.

We collected seawater for each experimental treatment by pumping from 1 m below the surface from a dock at AltaSea, in the Los Angeles Harbor. The water passed through a 250 µm mesh (to remove large grazers) (Carter et al., 2005; de la Broise and Palenik, 2007) and 22 bottles were filled in a ‘round robin’ manner to promote homogeneity between bottles filled at different times (filling occurred in < 20 mins). Once all bottles were full, with no headspace, half were randomly selected to receive a dosage of alkalinity (NaOH, CaO, or CaCl₂ + NaHCO₃). Bottles were opened, 50-150 mL of seawater was removed; stock water was pulled out of the stock solution foil bags into volumetric syringes ~~and injected into the bottles. and then pushed~~ through 0.2 µm PES syringe filters into the bottles. Any remaining headspace was once again filled with seawater; bottles were sealed and then inverted five times before being tied to the dock. All treatment bottles were given a dose of alkalinity on Day 0, some were immediately sampled, others left in the water column to experience natural changes in temperature and light conditions until their sampling day. Only one treatment experiment was conducted per week;

hence, we can only compare each alkalinity type to its respective control (no addition of alkalinity). Sampling occurred on days 0, 2, and 4 with three bottle replicates per time point. We also filled and added alkalinity to 2 bottles that were incubated devoid of light as well as 2 dark controls (wrapped in black tape, See Fig. 1). Results and discussion of all dark bottle experiments appear in the [Appendix A Supplemental Information](#).



Formatted: Font: Not Bold

2.5.2.4 Sample Processing

Bottles pulled out of the ocean were transported back to the lab (on ice) on their sampling day and weighed to track water that was used for sub-sampling. Bottles were gently inverted 5 times and then opened to measure temperature ($^{\circ}\text{C}$) with a probe and dissolved oxygen ($\mu\text{mol kg}^{-1}$) with PreSens Optical Oxygen Sensor Spots that were attached to each bottle. The oxygen measurements were corrected to the mean temperature recorded by the in situ HOBOTM Loggers. To avoid sampling from the surface layer of water in contact with the air, a ~6 inch piece of Tygon[®] F-4040-A attached to a 50 mL syringe via a one-way valve was inserted into bottom of the sample bottle. The first 15 mL of sample water was used to rinse the syringe, filter and needle that would be used to collect carbonate chemistry samples, ensuring no bubbles entered the sample storage containers as recommended by standard operating procedures (Dickson et al., 2007). Through subsequent sampling, 250 mL of sample water was filtered through a 25 mm 0.7 μm glass fiber filter (Whatmann GF/F) (~15 mins/bottle for filtration and subsampling). A portion of the filtrate was collected for 5mL duplicates of DIC (injected into septum-top of pre-

evacuated exetainers), alkalinity (15 mL Falcon tubes with no headspace), nutrients (15 mL with no headspace), dissolved organic carbon – DOC (40 mL pre-combusted amber glass septa vials), and 50 mL for archival purposes. We note that there could be some contamination in DOC samples because we used Tygon® to pull the sample into the syringe (Dickson et al., 2007); however, this would affect the controls and treatments in the same manner. The GF/F filter was wrapped in aluminum foil and placed in a freezer (-20 °C) for later Chl a analysis (< a month) (using a Turner Trilogy Fluorometer and the EPA SOP LG405.v11 for ‘Chlorophyll-A Non-Acidification Method’ (U.S. EPA, 2021)). One liter was filtered for future genomic analyses not discussed in this paper, on Supor 0.2 µm 47 mm PES filters, and stored at -80 °C. The remaining (4.5 L) seawater was filtered (detailed in Section 2.5.2) for particulate organic and inorganic carbon.

2.6.2.5 Measurements for ~~c~~Carbon ~~b~~Budget

A carbon budget for each bottle was created by assessing the total concentration of carbon in the dissolved and particulate pools. Carbonate system parameters were derived from TA and DIC measurements (discussed below) using CO2Sys.v3 (Pierrot et al., 2021) (Parameters: pH Scale = Total Scale (~~mol per kg SW~~), CO₂ Constants = (Dickson and Millero, 1987), KSO₄ = (Dickson, 1990), KF = (Perez and Fraga, 1987), Boron = (Lee et al., 2010)). Ω_{aragonite} values for treatments with added Ca²⁺ (CaO & [Ca²⁺] + [HCO₃⁻] Experiments) were calculated by multiplying the [CO₃²⁻] output from CO2Sys and [Ca²⁺] accounting for the increased Ca²⁺. All calculations use K_{sp}' for Aragonite in seawater (K_{sp}' = 10^{-6.19}). All other Ω_{aragonite} values derived from CO2Sys scale [Ca²⁺] to salinity.

~~2.6.12.5.1~~ Dissolved ~~c~~Carbon ~~p~~Phases

Samples for Dissolved Inorganic Carbon (DIC) were collected in pre-evacuated 12 mL borosilicate glass Labco Exetainers® and immediately placed at 4 °C for storage until analysis (< 1 week). For analysis, samples were acidified with 2 mL of 10 % H₃PO₄ (Spectrum Chemicals, Gardena, California) and the evolved CO₂ was analyzed by a Picarro G2121-i Cavity Ring Down Spectrometer (as detailed in (Subhas et al., 2015)). Optical grade calcite, with a known ¹³C value (2.47 ‰) was used to create an eight-point standard curve (ppmCO₂ per mg of CaCO₃) for each

run and a certified reference material (CRM-Batch 214) generated by Andrew Dickson's Lab at SIO, UCSD was used to correct all the final DIC concentrations (4 CRM replicates included per run). The total CO₂ standard curve produced with the Automate Sample Delivery System + Picarro CRDS had an $R^2 = 0.9999 \pm 0.0001$ and slope variability = 1.5 % for all runs in this dataset. Replicate CRM samples provided a measure of precision, $\pm 20 \mu\text{mol kg}^{-1}$. The isotopic values of replicate standards were within $\pm 0.2 \text{ ‰}$.

Samples for Dissolved Organic Carbon (DOC) were collected in 40 mL pre-combusted amber glass vials, acidified to 0.1 % v/v with concentrated hydrochloric acid (ACS grade, Supelco, Massachusetts), and placed in cold storage (4 °C) until analysis. The samples were run on a Shimadzu TOC-L Analyzer (NPOC method; Injection volume = 150 μL ; No. of Injections: Best 4 of 5; Max Variance of Replicates: 2.00 %). Within a run the standard curve had an $R^2 = 0.9998 \pm 0.0002$, between runs the reproducibility of the slope had 16.5 % error. Samples were initially purged with CO₂-free compressed air to remove inorganic carbon, combusted in a furnace via catalytic oxidation, and the generated CO₂ was measured by an infrared gas analysis. Standards for organic carbon were made with Oxalic Acid (dried at 70-°C for 2 hr), and CRMs from the Hansell Lab (University of Miami, RSMAS) were used to correct final concentrations (DSR Batch 22 Lot # 10-22) with a replicate precision of $\pm 10 \mu\text{mol kg}^{-1}$.

2.6.22.5.2 Particulate Carbon Phases

Roughly 4.5 L of seawater was filtered on 1-2 filters (0.7 μm 47mm Whatman GF/F) for particulate inorganic carbon (PIC) and particulate organic carbon (POC) analyses. After filtration, ~10 mL of buffered DI water (NaOH added to Milli-Q, pH 9) was used to rinse any residual seawater, and then filters were placed in the oven to dry overnight (60 °C). Filters were hole punched (diameter = duplicates of 6-12.5 mm); the hole punched fractions (2-8 % of total filtered area) were wrapped in tin foil capsules (9 x 5 mm) for combustion in a CosTech Elemental Analyzer – which is connected to the Picarro for CO₂ and ¹²C/¹³C analysis. This analysis of Total Carbon had a reproducibility of $\pm 1 \text{ ‰}$. The remaining fraction of the filters were rolled and placed into Labco Exetainers® – these samples were acidified with 5 mL 10 % H₃PO₄ and the evolved CO₂ measured similarly to DIC described above. Particulate Organic Carbon (POC) was calculated by subtracting the PIC ($\mu\text{gC L}^{-1}$) concentration from the Total Carbon ($\mu\text{gC L}^{-1}$) calculated for the whole filtered area.

2.6 Data Visualization and Statistical Analysis

Statistical analysis and data visualization was conducted in R and R Studio version 4.5.1. Means and standard deviations for each treatment by time point were generated using the `data_summary` function. Triplicate values were averaged, but data points that were found outside of three standard deviations from the mean were flagged as outliers and not included in the statistical analysis (Pukelsheim, 1994). The dataset was analyzed with two statistical approaches: (i) To test whether the effect of alkalinity addition had a statistically significant impact on PIC, POC, DOC and Chl *a* over time, we used a linear model with an interaction term and assessed if the p-value on the interaction term (i.e., the difference between the slopes of the control and the treatment) was statistically significant (significance threshold $\alpha = 0.05$) (results summarized Table 54). (ii) Since the previous method assumes linear relationships and normally distributed data (which may apply to some but not all variables) we also used a nonparametric two-way ANOVA (Analysis of Variance) – Scheirer Ray Hare test – to assess the impact of alkalinity addition on the Control and Treatment groups (significance threshold $\alpha = 0.05$) (results summarized in Table 56). The implications of these tests and any statistically significant responses are summarized in the context of ecological variability in the Results and Discussion sections.

3. Results

3.1 Simulating AWL with $[\text{Ca}^{2+}]$ and $[\text{HCO}_3^-]$ Addition

3.1.1 Short Term Impacts of Extremely Low Alkalinity Additions with high dilution

We intended to recreate an alkalinity impact condition that resembles extremely-high dilution of AWL after a few hours; an example outflow of alkalinity from a shipping vessel fitted with an AWL reactor would be about 3000-5000 $\mu\text{mol kg}^{-1}$ (Dong et al., 2025). As the ship's wake would dilute the outflow, we chose to dilute a stock ($\sim 4754 \mu\text{mol kg}^{-1}$) by different proportion; 1:150 (40 mL of stock), 1:300 (20 mL of stock), and 1:600 (10 mL of stock) (Table 32) based on dilution calculations presented by Dong et al., (2025) (inspired by (Caserini et al., 2021; Chou, 1996)). These incubations that lasted 3 hours were conducted on Santa Catalina Island.

As seen in Table 32, additions at what could be realistic deployment concentrations after several hours of mixing result in changes that are extremely close to replicate error and instrument measurement error. The control seawater had an alkalinity = $2223 \pm 6 \mu\text{mol kg}^{-1}$, DIC = $2015 \pm 21 \mu\text{mol kg}^{-1}$, $\Omega_{\text{aragonite}} = 2.41 \pm 0.02$, $\text{pCO}_2 = 536 \pm 4 \mu\text{atm}$, $\text{PIC} = 7.1 \pm 0.2 \mu\text{g L}^{-1}$, and $\text{POC} = 68.4 \pm 7.7 \mu\text{g L}^{-1}$. In the least diluted sample, the seawater alkalinity changes from $2223 \pm 6 \mu\text{mol kg}^{-1}$ to $2233 \pm 5 \mu\text{mol kg}^{-1}$ (the expected increase would be $2248 \mu\text{mol kg}^{-1}$) but there is no statistically significant impact on the concentrations of PIC or POC ($p > 0.05$). Since we saw no impact on PIC or POC at such small changes in alkalinity, and these additions were at the lower limit of our measurement capabilities, we focused the rest of the paper on experiments that perturb the alkalinity by larger amounts (25-50 %) and for longer durations (2-4 days). All the following experiments were conducted with water collected and incubated at AltaSea in LA Harbor.

Table 23. Summary of how concentration alkalinity amendments.

Treatment	Target ΔTA ($\mu\text{mol kg}^{-1}$)	Measured ΔTA ($\mu\text{mol kg}^{-1}$)	Target ΔDIC ($\mu\text{mol kg}^{-1}$)	Measured ΔDIC ($\mu\text{mol kg}^{-1}$)	Δ $\Omega_{\text{aragonite}}$	ΔpCO ₂ (μatm)	ΔPIC ($\mu\text{g L}^{-1}$)	ΔPOC ($\mu\text{g L}^{-1}$)
1:600	12	9.3 ± 8	12	11.1 ± 20	-0.01 ± 0.02	9 ± 8	-0.3 ± 0.2	0.6 ± 9.7
1:300	16	11.5 ± 5	16	21.8 ± 17	-0.09 ± 0.02	38 ± 8	-0.4 ± 0	-1.5 ± 6.7
1:150	25	10.4 ± 5	25	24.9 ± 22	-0.12 ± 0.06	51 ± 21	-0.1 ± 0.3	12.2 ± 20.5

The control seawater had an alkalinity = $2223 \pm 6 \mu\text{mol kg}^{-1}$, DIC = $2015 \pm 21 \mu\text{mol kg}^{-1}$, $\Omega_{\text{aragonite}} = 2.41 \pm 0.02$, $\text{pCO}_2 = 536 \pm 4 \mu\text{atm}$, $\text{PIC} = 7.1 \pm 0.2 \mu\text{g L}^{-1}$, and $\text{POC} = 68.4 \pm 7.7 \mu\text{g L}^{-1}$. The treatments are shown as dilution ratios. Expected TA and DIC have been calculated by only assuming $[\text{HCO}_3^-]$ has increased. The reported error is standard deviations between triplicate bottles ($n = 3$).

3.2 Impacts of higher alkalinity amendments after 0-4 days

All the following experiments with a 25-50% alkalinity amendment above the control seawater were conducted with water collected and incubated at AltaSea in LA Harbor. Table 4 summarizes the concentrations of PIC, POC, Chl a, DOC, $\Omega_{\text{aragonite}}$ values of the control seawater on Day 0, and the change in $\Omega_{\text{aragonite}}$ values after treatment.

Table 34. Initial concentrations of biogeochemical parameters at Day 0 (Control) of 4-day experiments

Formatted: Line spacing: single

Formatted: Font: Bold, Not Italic

Formatted: Font: Bold, Not Italic

Formatted: Font: Bold, Not Italic

Formatted: Font: Bold, Not Italic

Formatted: Font: Bold, Not Italic

Formatted: Indent: First line: 0.5", Line spacing: 1.5 lines

Target Treatment	Start Date	PIC $\mu\text{gC L}^{-1}$	POC $\mu\text{gC L}^{-1}$	CHL a $\mu\text{g L}^{-1}$	DOC $\mu\text{mol kg}^{-1}$	Control $\Omega_{\text{aragonite}}$	Treatment $\Omega_{\text{aragonite}}$
CaCl₂ + NaHCO₃ (~500 $\mu\text{mol kg}^{-1}$)	06-24-24	3.90 ± 1.15	278 ± 20	2.37 ± 0.25	N/A	2.02 ± 0.06	2.28 ± 0.05
	07-15-24	2.51 ± 0.30	209 ± 18	0.60 ± 0.21	117 ± 15	1.88 ± 0.07	2.12 ± 0.05
NaOH (~500 $\mu\text{mol kg}^{-1}$)	06-17-24	0.95 ± 0.21	366 ± 33	1.82 ± 0.32	N/A	2.28 ± 0.02	8.21 ± 0.04
	07-22-24	1.16 ± 1.02	369 ± 26	1.91 ± 0.40	96.1 ± 12.8	1.92 ± 0.04	7.63 ± 0.07
CaO (~500 $\mu\text{mol kg}^{-1}$)	11-18-24	100.0 ± 49.3	171 ± 48	0.09 ± 0.02	29.2 ± 1.15	1.47 ± 0.02	7.97 ± 0.23
	01-27-25	28.4 ± 4.53	279 ± 83	0.56 ± 0.07	66.2 ± 2.16	1.23 ± 0.14	8.63 ± 0.16
CaO (~1000 $\mu\text{mol kg}^{-1}$)	11-04-24	4.83 ± 0.79	218 ± 25	0.76 ± 0.07	65.2 ± 1.59	1.81 ± 0.04	13.73 ± 0.09

Table 3. Initial Concentrations of Biogeochemical Parameters at Day 0 (Control) of 4 day Experiments

	Start Date	PIC $\mu\text{gC L}^{-1}$	POC $\mu\text{gC L}^{-1}$	CHL $\mu\text{g L}^{-1}$	DOC $\mu\text{mol kg}^{-1}$	Control $\Omega_{\text{aragonite}}$	Treatment $\Omega_{\text{aragonite}}$
CaCl₂ + NaHCO₃ (~500 $\mu\text{mol kg}^{-1}$)	06-24-24	3.90 ± 1.15	278 ± 20	2.37 ± 0.25	N/A	2.02 ± 0.06	2.28 ± 0.05
	07-15-24	2.51 ± 0.30	209 ± 18	0.60 ± 0.21	117 ± 15	1.88 ± 0.07	2.12 ± 0.05
NaOH (~500 $\mu\text{mol kg}^{-1}$)	06-17-24	0.95 ± 0.21	366 ± 33	1.82 ± 0.32	N/A	2.28 ± 0.02	8.21 ± 0.04
	07-22-24	1.16 ± 1.02	369 ± 26	1.91 ± 0.40	96.1 ± 12.8	1.92 ± 0.04	7.63 ± 0.07
CaO (~500 $\mu\text{mol kg}^{-1}$)	11-18-24	100.0 ± 49.3	171 ± 48	0.09 ± 0.02	29.2 ± 1.15	1.47 ± 0.02	7.97 ± 0.23
	01-27-25	28.4 ± 4.53	279 ± 83	0.56 ± 0.07	66.2 ± 2.16	1.23 ± 0.14	8.63 ± 0.16
CaO (~1000 $\mu\text{mol kg}^{-1}$)	11-04-24	4.83 ± 0.79	218 ± 25	0.76 ± 0.07	65.2 ± 1.59	1.81 ± 0.04	13.73 ± 0.09

3.2.1.2 [Ca²⁺] and [HCO₃⁻] additions: Alkalinity enhancement ~500 $\mu\text{mol kg}^{-1}$

We performed two [Ca²⁺] and [HCO₃⁻] addition experiments on 24 June 2024 and 15 July 2024 which each lasted four days. Alkalinity added in the form of HCO₃⁻ increases both alkalinity and DIC in a 1:1 ratio – hence, the change in pH relative to the Control is minimal for

Formatted: Font: Not Italic, Font color: Black

Formatted: Line spacing: 1.5 lines

this treatment (Fig. 2). Over the incubation period, this treatment resulted in no statistically significant difference in the production of PIC ($p > 0.05$, ANOVA and Linear Regression method, see Table 54 & 65), the $\Omega_{\text{aragonite}}$ only increased by 0.2 units in both experiments (Table 43). Chl a shows that changes in concentration over time were similar in both the control and the treatment (Fig. 2). The $[\text{Ca}^{2+}] + [\text{HCO}_3^-]$ addition experiments intentionally showed the largest increase within DIC pool compared to all other alkalinity enhancements. There was a concurrent increase in both organic carbon pools of POC and DOC, but it was not statistically dissimilar from the changes within the control condition ($p > 0.05$, ANOVA and Linear Regression method). The partitioning of carbon between the inorganic and organic pools is further discussed in Section 3.4.2 Carbon Partitioning.

Table 3. Initial Concentrations of Biogeochemical Parameters at Day 0 (Control) of 4 day Experiments


	<u>Start Date</u>	<u>PIC</u> <u>μgCL^{-1}</u>	<u>POC</u> <u>μgCL^{-1}</u>	<u>Chl</u> <u>$\mu\text{g L}^{-1}$</u>	<u>DOC</u> <u>$\mu\text{mol kg}^{-1}$</u>	<u>Control</u> <u>$\Omega_{\text{aragonite}}$</u>	<u>Treatment</u> <u>$\Omega_{\text{aragonite}}$</u>
<u>$\text{CaCl}_2 + \text{NaHCO}_3$</u> <u>(+500 $\mu\text{mol kg}^{-1}$)</u>	<u>06-24-24</u>	<u>3.90 ± 1.15</u>	<u>278 ± 20</u>	<u>2.37 ± 0.25</u>	<u>N/A</u>	<u>2.02 ± 0.06</u>	<u>2.28 ± 0.05</u>
	<u>07-15-24</u>	<u>2.51 ± 0.30</u>	<u>209 ± 18</u>	<u>0.60 ± 0.21</u>	<u>117 ± 15</u>	<u>1.88 ± 0.07</u>	<u>2.12 ± 0.05</u>
<u>NaOH</u> <u>(+500 $\mu\text{mol kg}^{-1}$)</u>	<u>06-17-24</u>	<u>0.95 ± 0.21</u>	<u>366 ± 33</u>	<u>1.82 ± 0.32</u>	<u>N/A</u>	<u>2.28 ± 0.02</u>	<u>8.21 ± 0.04</u>
	<u>07-22-24</u>	<u>1.16 ± 1.02</u>	<u>369 ± 26</u>	<u>1.91 ± 0.40</u>	<u>96.1 ± 12.8</u>	<u>1.92 ± 0.04</u>	<u>7.63 ± 0.07</u>
<u>CaO</u> <u>(+500 $\mu\text{mol kg}^{-1}$)</u>	<u>11-18-24</u>	<u>100.0 ± 49.3</u>	<u>171 ± 48</u>	<u>0.09 ± 0.02</u>	<u>29.2 ± 1.15</u>	<u>1.47 ± 0.02</u>	<u>7.97 ± 0.23</u>
	<u>01-27-25</u>	<u>28.4 ± 4.53</u>	<u>279 ± 83</u>	<u>0.56 ± 0.07</u>	<u>66.2 ± 2.16</u>	<u>1.23 ± 0.14</u>	<u>8.63 ± 0.16</u>
<u>CaO</u> <u>(+1000 $\mu\text{mol kg}^{-1}$)</u>	<u>11-04-24</u>	<u>4.83 ± 0.79</u>	<u>218 ± 25</u>	<u>0.76 ± 0.07</u>	<u>65.2 ± 1.59</u>	<u>1.81 ± 0.04</u>	<u>13.73 ± 0.09</u>

3.2.2 NaOH addition: Alkalinity Enhancement ~500 $\mu\text{mol kg}^{-1}$

NaOH additions occurred on 17 June 2024 and 22 July 2024. The addition of NaOH increases alkalinity but not DIC which results in a large change in $\Omega_{\text{aragonite}}$ from ~2 to 8.21 ± 0.04 (06-17-24; pH_T increases 0.63) and 7.63 ± 0.07 (07-22-24; pH_T increases 0.71). When compared to the control condition, the changes in PIC, POC, DOC, and Chl a in the treatment conditions are also not statistically significant (Fig. 3) via either of two statistical analyses ($p > 0.05$, ANOVA and Linear Regression methods). As with the $[\text{Ca}^{2+}] + [\text{HCO}_3^-]$ additions,

Formatted: Subscript

398 there are certain points in time where the controls diverge from the treatments, but there are no
399 consistent trends across multiple variables (i.e., no indication that one group (control/treatment)
400 is consistently higher/lower through time).



Formatted: Centered






Figure 2. Water Chemistry from (NaOH and $[\text{Ca}^{2+}] + [\text{HCO}_3^-]$ -addition) aAmendment eExperiments. Panels are grouped by the alkalinity treatment they received: Controls (green), NaOH (red) and $\text{CaCl}_2 + \text{NaHCO}_3$ (blue). The start date of the experiment is listed for reference to differentiate between duplicate experiments. The four variables displayed here are measured Alkalinity ($\mu\text{mol kg}^{-1}$), measured Dissolved Inorganic Carbon (DIC, $\mu\text{mol kg}^{-1}$), pH_T (Total Scale) calculated from CO2Sys.v3, and Dissolved Oxygen (O_2 , $\mu\text{mol kg}^{-1}$). Each plot shows the average of the treatment per timepoint (n =

Formatted: Subscript

3), and the error bars are the standard deviation between the triplicate bottles. Smaller symbols show individual bottles.





412
413 **Figure 3. Four dDay tTimeseries of bBiogeochemical rResponse to dDifferent aAlkalinity**
414 **eEnhancements.** Smaller symbols show individual bottles; larger symbols show the mean. Error bars are
415 the standard deviation between replicates ($n = 3$). The start date of the experiment and the amount of
416 alkalinity added are included as headers for reference. The variables displayed are Chlorophyll a (Chl

a, $\mu\text{g L}^{-1}$), Particulate Inorganic Carbon (PIC, $\mu\text{g L}^{-1}$), Particulate Organic Carbon (POC, $\mu\text{g L}^{-1}$), and Dissolved Organic Carbon (DOC, $\mu\text{mol kg}^{-1}$).

3.3 CaO ~~a~~Additions

3.3.1 CaO addition: ~~a~~Alkalinity ~~e~~Enhancement $\sim 1000 \mu\text{mol kg}^{-1}$ (4 November 2024)

We raised the $\Omega_{\text{aragonite}}$ value, with CaO addition, above the threshold for spontaneous precipitation of CaCO_3 in seawater ($\Omega_{\text{aragonite}} > 12$, $\Omega_{\text{calcite}} > 19$ (Marion et al., 2009; Morse and He, 1993)) according to the references cited. Increasing TA by $\sim 1000 \mu\text{mol kg}^{-1}$ (~~measured $\Delta\text{TA} = 923 \pm 12 \mu\text{mol kg}^{-1}$~~) raised the $\Omega_{\text{aragonite}}$ from 1.81 ± 0.04 to 13.73 ± 0.09 and pH_T was raised from 7.85 ± 0.01 to 8.91 ± 0.01 (error is standard deviation between replicate bottles). Yet in these incubations, we see no statistically significant increase in PIC either immediately after CaO addition or throughout the 4-day experiment ($p > 0.05$, nonparametric ANOVA and linear analysis, Table ~~54~~ & ~~65~~). We also see no statistically significant divergence between the Control bottles and CaO bottles in DOC or Chl a. However, the O_2 and POC trends are statistically different than the Control after CaO addition ($p < 0.05$ with linear analysis, Table ~~54~~). This is the only experiment where we see a statistically significant impact of CaO addition with a 95% confidence interval. There is production of $\text{O}_2 = 70 \pm 20 \mu\text{mol kg}^{-1}$ over 4 days in the Control, compared to no change in measured O_2 in the CaO treatment. In the Control, the POC production increases by $180 \pm 35 \mu\text{gC L}^{-1}$ ($\sim 15 \pm 3 \mu\text{mol kg}^{-1}$) compared to only a $70 \pm 40 \mu\text{gC L}^{-1}$ ($\sim 5.8 \pm 3.6 \mu\text{mol kg}^{-1}$) increase in the treated waters. Further implications of these parameters are discussed in Section 4.2.2.

3.3.2 CaO addition: ~~a~~Alkalinity ~~e~~Enhancement $\sim 500 \mu\text{mol kg}^{-1}$ with ~~h~~Higher ~~a~~Ambient PIC (18 November 2024)

We conducted two CaO addition experiments to enhance alkalinity by $\sim 500 \mu\text{mol kg}^{-1}$, similar to the amount of enhanced alkalinity in the $[\text{Ca}^{2+}] + [\text{HCO}_3^-]$ and NaOH treatments. In the 18 November 2024 experiment, PIC concentrations in the harbor water were ~ 20 times higher than the ambient PIC ($5 \pm 5 \mu\text{gC L}^{-1}$) of the experiments described above (Fig. 3 & Fig. 5). We used this opportunity as an analog to study an alkalinity addition with greater suspended solids – i.e., greater nucleation sites for secondary precipitation. The CaO addition raised pH_T from 7.75 ± 0.01 to 8.57 ± 0.01 , and $\Omega_{\text{aragonite}}$ from 1.47 ± 0.02 to 7.97 ± 0.23 . We saw no

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Superscript/ Subscript


Formatted: Font: 12 pt

Formatted: Font: 12 pt, Not Superscript/ Subscript

increase in PIC formation in the treatment compared to the control. Instead, we observed a loss of PIC occurring over time in both the control and treated bottles. Chl a, DOC, and dissolved O₂ show no statistically significant changes over time from the alkalinity addition; POC in the Control samples is higher than the treatment (Fig. 5), however, Day 0 variability negates any statistically significant impact.

3.3.3 CaO Addition: Alkalinity Enhancement ~500 $\mu\text{mol kg}^{-1}$ (27 January 2025)

We replicated the previous experiment with the goal to capture a lower ambient PIC signal, but in this experiment the average starting PIC was 6-10 times the normal background (30-50 $\mu\text{gC L}^{-1}$ compared to ~5 $\mu\text{gC L}^{-1}$). Upon CaO addition the system experienced a pH increase of 0.95 units ($\text{pH}_{\text{final}} = 8.63 \pm 0.01$), and $\Omega_{\text{aragonite}}$ increased from 1.23 ± 0.14 to 8.63 ± 0.16 . While there were samples in which the values of PIC concentration in the treatment samples averaged higher than the Controls, the overall 4-day trends are not statistically significant. The Control averages 1 $\mu\text{g L}^{-1}$ Chl a higher than the treatment and almost 100 $\mu\text{gC L}^{-1}$ POC higher than the treatment by Day 4. Yet, due to high variability between bottles, the differences in POC and DOC between the controls and treatment are not statistically different when tested with nonparametric ANOVA or linear analysis. Of note, the differences in Chl a overtime are statistically significant when analyzed with a linear regression analysis and a more lenient threshold for significance (Table 54, $p = 0.08$, $\alpha = 0.1$) but are not significant with a nonparametric ANOVA.



Formatted: Centered






Figure 4. Water Chemistry from (CaO) aAmendment cExperiments. Panels are grouped by the alkalinity treatment they received: Controls (green) and CaO (purple). The start date of the experiment is listed for reference to differentiate between duplicate experiments. The four variables displayed here are measured Alkalinity ($\mu\text{mol kg}^{-1}$), measured Dissolved Inorganic Carbon (DIC, $\mu\text{mol kg}^{-1}$), pH_T (Total Scale) calculated from CO2Sys.v3, and Dissolved Oxygen (O_2 , $\mu\text{mol kg}^{-1}$). Each plot shows the average of the treatment per timepoint ($n = 3$), and the error bars are the standard deviation between the triplicate bottles. Smaller symbols show individual bottles.

Formatted: Subscript



Formatted: Centered



476
477 **Figure 5. Four dDay tTimeseries of bBiogeochemical rResponse to dDifferent aAlkalinity**
478 **eEnhancements.** Smaller symbols show individual bottles, large symbol shows mean. Error bars are the
479 standard deviation between replicates (n = 3). The start date of the experiment and the amount of
480 alkalinity added are included as headers for reference. The variables displayed are Chlorophyll a (Chl a,
481 $\mu\text{g L}^{-1}$), Particulate Inorganic Carbon (PIC, $\mu\text{g L}^{-1}$), Particulate Organic Carbon (POC, $\mu\text{g L}^{-1}$), and
482 Dissolved Organic Carbon (DOC, $\mu\text{mol kg}^{-1}$).

483 Table 54. Summary of Linear Model Analysis by Experiment.

	Experiment Number	[ALK Added]	Variable	Control Slope $\pm$ se	Treatment Slope $\pm$ se	<i>P</i> -value Interaction Term	Significance
06-17-24	NaOH	500	PIC	0.23 $\pm$ 0.17	-0.02 $\pm$ 0.17	0.33	FALSE
	NaOH	500	POC	64.6 $\pm$ 14.5	69.3 $\pm$ 16.1	0.83	FALSE
	NaOH	500	CHL	0.87 $\pm$ 0.29	0.67 $\pm$ 0.33	0.67	FALSE
	NaOH	500	O ₂	4.14 $\pm$ 2.97	4.96 $\pm$ 3.65	0.86	FALSE
07-22-24	NaOH	500	PIC	0.69 $\pm$ 0.17	0.53 $\pm$ 0.17	0.50	FALSE
	NaOH	500	POC	21.4 $\pm$ 7.25	7.03 $\pm$ 7.25	0.18	FALSE
	NaOH	500	DOC	-0.16 $\pm$ 4.63	9.72 $\pm$ 4.63	0.15	FALSE
	NaOH	500	CHL	0.03 $\pm$ 0.12	-0.10 $\pm$ 0.12	0.46	FALSE
	NaOH	500	O ₂	9.61 $\pm$ 1.65	10.5 $\pm$ 1.65	0.69	FALSE
06-24-24	[Ca ²⁺] + [HCO ₃ ⁻]Ca ²⁺ + HCO ₃ ⁻	500					
	[Ca ²⁺] + [HCO ₃ ⁻]Ca ²⁺ + HCO ₃ ⁻	500	PIC	-0.16 $\pm$ 0.19	0.02 $\pm$ 0.19	0.53	FALSE
	[Ca ²⁺] + [HCO ₃ ⁻]Ca ²⁺ + HCO ₃ ⁻		POC	156 $\pm$ 25	170 $\pm$ 28.6	0.73	FALSE
	[Ca ²⁺] + [HCO ₃ ⁻]Ca ²⁺ + HCO ₃ ⁻	500	CHL	1.10 $\pm$ 0.54	0.26 $\pm$ 0.54	0.29	FALSE
	[Ca ²⁺] + [HCO ₃ ⁻]Ca ²⁺ + HCO ₃ ⁻	500	O ₂	13.5 $\pm$ 2.82	13.1 $\pm$ 3.12	0.93	FALSE
	[Ca ²⁺] + [HCO ₃ ⁻]Ca ²⁺ + HCO ₃ ⁻	500					
	[Ca ²⁺] + [HCO ₃ ⁻]Ca ²⁺ + HCO ₃ ⁻	500	PIC	0.72 $\pm$ 0.22	0.68 $\pm$ 0.22	0.91	FALSE
	[Ca ²⁺] + [HCO ₃ ⁻]Ca ²⁺ + HCO ₃ ⁻	500	POC	98.4 $\pm$ 16	98.9 $\pm$ 16	0.98	FALSE
07-15-24	[Ca ²⁺] + [HCO ₃ ⁻]Ca ²⁺ + HCO ₃ ⁻	500	DOC	2.83 $\pm$ 5.18	7.05 $\pm$ 5.18	0.57	FALSE
	[Ca ²⁺] + [HCO ₃ ⁻]Ca ²⁺ + HCO ₃ ⁻	500	CHL	0.78 $\pm$ 0.47	0.53 $\pm$ 0.47	0.71	FALSE
	[Ca ²⁺] + [HCO ₃ ⁻]Ca ²⁺ + HCO ₃ ⁻	500	O ₂	29.3 $\pm$ 2.81	22.8 $\pm$ 3.12	0.14	FALSE
	[Ca ²⁺] + [HCO ₃ ⁻]Ca ²⁺ + HCO ₃ ⁻	500					
	[Ca ²⁺] + [HCO ₃ ⁻]Ca ²⁺ + HCO ₃ ⁻	500					
	[Ca ²⁺] + [HCO ₃ ⁻]Ca ²⁺ + HCO ₃ ⁻	500					
11-18-24	CaO	500	PIC	-19.0 $\pm$ 6.39	-18.1 $\pm$ 6.39	0.92	FALSE
	CaO	500	POC	52.7 $\pm$ 14.4	27.8 $\pm$ 12.9	0.22	FALSE
	CaO	500	DOC	-0.41 $\pm$ 2.9	-1.12 $\pm$ 0.32	0.13	FALSE
	CaO	500	CHL	0.53 $\pm$ 0.12	0.61 $\pm$ 0.21	0.68	FALSE
	CaO	500	O ₂	1.92 $\pm$ 4.13	-0.64 $\pm$ 4.14	0.86	FALSE

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

01-27-25	CaO	500	PIC	1.58 ± 1.99	1.08 ± 1.99	0.86	FALSE
	CaO	500	POC	23.5 ± 11.1	8.03 ± 11.1	0.34	FALSE
	CaO	500	DOC	-0.49 ± 0.53	-0.74 ± 0.53	0.74	FALSE
	CaO	500	CHL	0.53 ± 0.08	0.32 ± 0.08	0.08*	FALSE
	CaO	500	O ₂	13.8 ± 0.97	10.9 ± 1.07	NA	$n < 3$
11-04-24	CaO	1000	PIC	-0.34 ± 0.15	-0.26 ± 0.15	0.68	FALSE
	CaO	1000	POC	45.7 ± 8.85	17.8 ± 8.85	0.04	TRUE
	CaO	1000	DOC	0.07 ± 0.36	-0.58 ± 0.32	0.20	FALSE
	CaO	1000	CHL	0.21 ± 0.13	0.24 ± 0.13	0.88	FALSE
	CaO	1000	O ₂	17.2 ± 4.86	-0.86 ± 4.37	0.01	TRUE

Table shows the slope $\pm$ the standard error around the slope of the Control and Treatment groups by the variable being analyzed. The p-value is shown, and the test result indicating statistical significance is shown when $p < 0.05$. The * marks a relaxed significance threshold ($\alpha = 0.1$). TRUE OR FALSE indicates whether the interaction is statistically significant or not. Bold for emphasis. NA written where there are less than three data points per time point for control and treatment.

Table 65. Summary of Two-Way Non-parametric Analysis of Variance (ANOVA) by Experiment.

	Experiment Number	[ALK Added]	Variable	P-value Interaction Term	Significance
06-17-24	NaOH	500	PIC	0.55	FALSE
	NaOH	500	POC	0.88	FALSE
	NaOH	500	CHL	0.88	FALSE
	NaOH	500	O ₂	0.93	FALSE
07-22-24	NaOH	500	PIC	0.52	FALSE
	NaOH	500	POC	0.22	FALSE
	NaOH	500	DOC	0.24	FALSE
	NaOH	500	CHL	0.44	FALSE
06-24-24	$[\text{Ca}^{2+}] + [\text{HCO}_3^-]\text{Ca}^{2+} + \text{HCO}_3^-$	500	PIC	0.39	FALSE
	$[\text{Ca}^{2+}] + [\text{HCO}_3^-]\text{Ca}^{2+} + \text{HCO}_3^-$	500	POC	0.94	FALSE
	$[\text{Ca}^{2+}] + [\text{HCO}_3^-]\text{Ca}^{2+} + \text{HCO}_3^-$	500	CHL	0.56	FALSE
	$[\text{Ca}^{2+}] + [\text{HCO}_3^-]\text{Ca}^{2+} + \text{HCO}_3^-$	500	O ₂	0.95	FALSE
	$[\text{Ca}^{2+}] + [\text{HCO}_3^-]\text{Ca}^{2+} + \text{HCO}_3^-$	500		0.89	FALSE
	$[\text{Ca}^{2+}] + [\text{HCO}_3^-]\text{Ca}^{2+} + \text{HCO}_3^-$	500		0.53	FALSE
07-15-24	$[\text{Ca}^{2+}] + [\text{HCO}_3^-]\text{Ca}^{2+} + \text{HCO}_3^-$	500	PIC		FALSE
	$[\text{Ca}^{2+}] + [\text{HCO}_3^-]\text{Ca}^{2+} + \text{HCO}_3^-$	500	POC		FALSE

	$[\text{Ca}^{2+}] + [\text{HCO}_3^-] \text{Ca}^{2+}$	500		0.18	FALSE
	$+ \text{HCO}_3^-$		DOC		
	$[\text{Ca}^{2+}] + [\text{HCO}_3^-] \text{Ca}^{2+}$	500		0.82	FALSE
	$+ \text{HCO}_3^-$		CHL		
	$[\text{Ca}^{2+}] + [\text{HCO}_3^-] \text{Ca}^{2+}$	500			FALSE
	$+ \text{HCO}_3^-$		O ₂	0.83	
11-18-24	CaO	500	PIC	0.74	FALSE
	CaO	500	POC	0.66	FALSE
	CaO	500	DOC	0.55	FALSE
	CaO	500	CHL	0.71	FALSE
	CaO	500	O ₂	0.98	FALSE
01-27-25	CaO	500	PIC	0.90	FALSE
	CaO	500	POC	0.86	FALSE
	CaO	500	DOC	0.78	FALSE
	CaO	500	CHL	0.97	FALSE
	CaO	500	O ₂	0.74	FALSE
11-04-24	CaO	1000	PIC	0.72	FALSE
	CaO	1000	POC	0.64	FALSE
	CaO	1000	DOC	0.15	FALSE
	CaO	1000	CHL	0.54	FALSE
	CaO	1000	O ₂	0.34	FALSE

The Scheirer Ray Hare test is used for this analysis. The interaction between treatment (Control vs. Treatment) condition and time on PIC, POC, Chl a, DOC, and dissolved O₂ is summarized by the p-value. TRUE indicates whether the interaction is statistically significant with an ($\alpha = 0.05$), FALSE shows an insignificant impact on the measured "Variable" column.

3.4 Carbon Budgets

3.4.1 Carbon Isotopes to Verify Estimates of Respiration

Organic carbon is subject to net increases due to production and decreases attributed to respiration. We added a ¹³C isotope spike to controls and treatment bottles to assess if an alkalinity addition would change organic matter remineralization; the remineralized organic matter would add unspiked DIC into a spiked DIC pool. The isotope addition ~~only~~ increased DIC by ~~2-5~~ $\mu\text{mol kg}^{-1}$, the starting isotopic $\delta^{13}\text{DIC}$ varied by bottle. Hence to compare across various days and treatments, we divided the measured $\delta^{13}\text{DIC}$ value from each bottle by the average $\delta^{13}\text{DIC}$ on Day 0 Control or Treatment value. An increase in organic matter remineralization would make the pool of $\delta^{13}\text{DIC}$ lighter. The effect of treatment on $\delta^{13}\text{DIC}$ was tested with a two-factor ANOVA where the variables 'Experiment Day' and 'Treatment' were taken as factors. The

508 interaction of these factors results in a $p > 0.05$ for all experiments. Hence, there were no
509 consistent and sustained changes in the $\delta^{13}\text{DIC}$ pool of the treatment bottles when compared to
510 the control bottles (Fig. 6). There was also no decrease in the $\delta^{13}\text{DIC}$ values due to invasion of
511 CO_2 even in the base addition experiments, because there was no headspace in the bottles, i.e.
512 this isotope tracer demonstrated that a closed system was maintained.

513

Formatted: Centered





Figure 6. Normalized Seawater (DIC) $\delta^{13}\text{C}$ ‰ over Time in all Experiments. Panel (07-15-24) + 500 $\mu\text{mol kg}^{-1}$ $\text{CaCl}_2 + \text{NaHCO}_3$, (07-22-24) + 500 $\mu\text{mol kg}^{-1}$ NaOH , (11-04-24) + 1000 $\mu\text{mol kg}^{-1}$ CaO , (11-18-24) + 500 $\mu\text{mol kg}^{-1}$ CaO with higher ambient PIC, and (01-27-25) + 500 $\mu\text{mol kg}^{-1}$ CaO . Each data point has an $n = 3$, and error bars are the standard deviation of the averaged values. Data was ‘normalized’ by dividing each measured value with the initial starting $\delta^{13}\text{C}$ value of the DIC on Day 0.

3.4.2 Carbon Partitioning

The purpose of conducting closed system experiments was to better understand and constrain the partitioning of carbon between various pools after increasing the amount of alkalinity. Here we present the distribution of carbon in various pools (DIC, DOC, PIC, POC) and track the movement of carbon over the span of four days (Fig. 7). These distributions are generated from the mean measured values of triplicate bottles at each time point. In our experiments, ~95 % of the carbon exists as DIC, ~3.5 % is stored as DOC, ~ 1.2 % is found in the POC pool, and < 0.1 % in the PIC pool. The non-DIC pool is highly variable across the “Day 0” starting conditions of all experiments (summarized in Table 43). In many experiments, the addition of alkalinity does not change carbon distribution relative to the controls (Fig. 7). The decrease in %DIC (Panel a & b, Fig. 7) is matched by an increase of either %POC or %DOC, which is expected from photosynthesis and organic matter cycling.




Figure 7. Total Carbon Budget for All Experiments. Panel (a) + 500 $\mu\text{mol kg}^{-1}$ $\text{CaCl}_2 + \text{NaHCO}_3$, (b) + 500 $\mu\text{mol kg}^{-1}$ NaOH , (c) + 500 $\mu\text{mol kg}^{-1}$ CaO , (d) + 1000 $\mu\text{mol kg}^{-1}$ CaO , and (e) + 500 $\mu\text{mol kg}^{-1}$ CaO with higher (ambient) particles of inorganic carbon. Bar plots are an average of 3 bottles per timepoint. Plots show changes in distribution of carbon over the duration of the experiment between treatments and controls. The total amount of C will always = 100%.

We can use a mass-balance approach to check if the carbon pools change over time. For example, one assumption may be that decline in DIC is compensated by increase in POC. In this case, one would expect a 1:1 relationship between ΔDIC and ΔPOC (Fig. 8a). However, the $[\text{Ca}^{2+}] + [\text{HCO}_3^-]$ treatments fall furthest from this line, which means these experiments show a deficit of POC production relative to DIC drawdown.

To account for DIC lost through other processes, we considered the formation of CaCO_3 as a DIC sink, assuming 1 mole of PIC growth is equivalent to 1 mole of DIC consumed. Our data indicate that the DIC lost to the precipitation of CaCO_3 is $< 1 \mu\text{mol kg}^{-1}$ over the span of 4 days; this is a trivial amount. We also considered CO_2 loss from the permeability of the PETG bottles. Assuming that CO_2 permeability for PETG is $125 \text{ cc-mil inch}^{-2} \text{ day}^{-1}$, wall thickness =

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

551 0.76 mm, surface area $\sim 300 \text{ inch}^2$ and the $\Delta p\text{CO}_2$ gradient between water inside the bottle vs.
 552 outside in the ocean = $300 \mu\text{atm}$, the maximum amount of CO_2 that could diffuse out of the
 553 bottle would be $3 \mu\text{mol kg}^{-1} \text{ day}^{-1}$. Therefore, only $\sim 13 \mu\text{mol kg}^{-1}$ DIC could potentially be lost to
 554 CaCO_3 precipitation or diffusion through the bottles' walls, and the $50\text{-}100 \mu\text{mol kg}^{-1}$ decrease in
 555 DIC in Figure 8(a) must result from production of POC and DOC.

556 As an additional test of carbon partitioning, we can check to see whether these
 557 experiments fall within Redfield stoichiometries by comparing the total drawdown of DIC with
 558 the change in dissolved oxygen (Fig. 8 (b)). A range of DIC: O_2 ratios are considered. The $[\text{Ca}^{2+}]$
 559 $+ [\text{HCO}_3^-]$ experiments and one of their controls fall furthest from the range of
 560 Redfield ratios. The same $[\text{Ca}^{2+}] + [\text{HCO}_3^-]$ experiments that show a deficit in POC
 561 also show a deficit in dissolved oxygen which would be expected if DIC consumption were
 562 producing DOC if the POC was being remineralized.

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt




Figure 8. Comparison of stoichiometric changes in DIC, Dissolved O₂, and POC in terms of the Redfield Ratio. (a) Cross plot of measured decrease of DIC versus the measured increase of POC; solid

line is the 1:1 line. Points that fall below the 1:1 line are thought to have increased DOC. (b) Cross plot of measured decrease of DIC versus measured production of dissolved oxygen. Plotted lines show the slopes of various Redfield Ratios: solid line ($-C:O_2 = -106:161$), dot-dash line ($-C:O_2 = -106:141$), and dashed line ($-C:O_2 = -106:-106$ i.e. a 1:1 ratio) (Anderson, 1995). All points in (a) and (b) have a $n = 3$, and the delta is calculated as the difference between the average of Day 0 and Day 4 bottle values. Error bars are propagated standard deviations of averaged values used to generate the delta values.

4. Discussion

If alkalinity is added to the ocean, we would need to know the effects of different minerals/alkalinity sources – i.e., to test if one produces more rapid changes to the ocean ecosystem than another. Thus, we conducted standardized experiments with different feedstocks that raised alkalinity by ~~the same~~similar thresholds (~ 500 and $\sim 1000 \mu\text{mol kg}^{-1}$). OAE functions on the principle of raising pH which will promote uptake of atmospheric CO_2 . One of the simulations we tested was the addition of $[\text{Ca}^{2+}] + [\text{HCO}_3^-]$ which represents the products of accelerated weathering of limestone (Dong et al., 2025) yet does not drastically raise pH like NaOH or CaO addition. CaO addition increases both pH and the $[\text{Ca}^{2+}]$, hence the resultant change in $\Omega_{\text{aragonite}}$ is slightly higher than the change in $\Omega_{\text{aragonite}}$ after NaOH addition. Here we discuss how these different regimes influenced abiotic effects such as precipitation of minerals and biotic effects such as organic matter cycling.

4.1 The ~~a~~absence of ~~s~~secondary CaCO_3 ~~p~~precipitation

The likelihood of observing mineral precipitation after OAE is controlled by several factors including supersaturation of the seawater with respect to a particular mineral, nucleation sites, temperature, time, and even interference from catalytic or inhibitor ions (Berner, 1975; Marion et al., 2009; Moras et al., 2024; Morse et al., 2007; Morse and He, 1993; Mucci and Morse, 1983; Naviaux et al., 2019; Nguyen Dang et al., 2017; Suitner et al., 2026). Here we summarize similar studies that used NaOH, NaHCO_3 , and CaO to find the minimum thresholds for precipitation of CaCO_3 after OAE (detailed experimental conditions are included in ~~Supplemental~~ Table B1 & B2 in Appendix B).

Several other studies that conducted similar manipulations of alkalinity additions to natural seawater (initial $\Omega_{\text{aragonite}} = 1-3$) have reported no observed precipitation of CaCO_3 (Hartmann et al., 2023; Moras et al., 2022; Ringham et al., 2024; Suitner et al., 2024). Of these, experiments conducted with $\text{NaHCO}_3 + \text{Na}_2\text{CO}_3$ or NaOH have shown stable seawater alkalinity up to $\Omega_{\text{aragonite}} \approx 15-19$ for at least four days and longer (~ 20 days) in closed systems (Hartmann et al., 2023; Moras et al., 2022; Suitner et al., 2024). In an open system that was allowed to equilibrate with atmospheric CO_2 , no precipitation was observed after NaOH addition up to $\Omega_{\text{aragonite}} \approx 27$ for at least 20 days (Ringham et al., 2024). At present, we are the only study to use CaO in an aqueous form. The one study that does not see CaCO_3 precipitation but uses solid CaO and $\text{Ca}(\text{OH})_2$ feedstocks to add alkalinity, only raised alkalinity by $250 \mu\text{mol kg}^{-1}$ (raising $\Omega_{\text{aragonite}}$ from 2 to 5), after which higher additions resulted in precipitation (Moras et al., 2022).

In these same experimental conditions discussed above, the threshold for CaCO_3 precipitation was surpassed after $\Omega_{\text{aragonite}}$ increased, temperature increased, or the mineral induction period where alkalinity is stable had ended. In some cases, precipitation was favorable due to the presence of nucleation sites (Hartmann et al., 2023; Hashim et al., 2025; Moras et al., 2022, 2024; Ringham et al., 2024; Suitner et al., 2024; Varliero et al., 2024). These are all parameters that vary in a given oceanic environment. With $\text{NaHCO}_3 + \text{Na}_2\text{CO}_3$ additions or NaOH additions where no solid feedstock was used, CaCO_3 precipitated as early as 1 day when $\Omega_{\text{aragonite}} > 30$ (Hartmann et al., 2023; Ringham et al., 2024; Suitner et al., 2024) or after 20 days when $\Omega_{\text{aragonite}}$ exceeded 15 (Hartmann et al., 2023; Suitner et al., 2024). At warmer surface temperatures (27°C) precipitation started at $\Omega_{\text{aragonite}} \approx 10$ (NaOH addition) within hours in a closed system (Hashim et al., 2025). In the experiments where nucleation sites were present, precipitation could be observed as low as $\Omega_{\text{aragonite}} \sim 7$ with CaO and $\text{Ca}(\text{OH})_2$ (Moras et al., 2022; Varliero et al., 2024). The influence of parameters such as surface area for nucleation and oversaturation with respect to $\Omega_{\text{aragonite}}$ on runaway CaCO_3 precipitation have been further constrained in observation driven modeling efforts (Suitner et al., 2026).

As our alkalinity additions only raised $\Omega_{\text{aragonite}}$ to 2 (for $[\text{Ca}^{2+}] + [\text{HCO}_3^-]$ addition) and $\Omega_{\text{aragonite}}$ to 7-8 (NaOH) for 4 days, which is within the range of no observed precipitation of other studies, we believe our results are consistent with existing literature. With our CaO addition we raised $\Omega_{\text{aragonite}}$ to 8-14, and we expected CaCO_3 as our saturation values were above that of

Formatted: Not Superscript/ Subscript

Formatted: Subscript

Moras et al. (2022). However, our experimental conditions did not have ~~mineral solid~~ feedstock like Moras et al. (2022), which could explain the lack of precipitation. Given our results, OAE practitioners in LA Harbour, perhaps analogous to other near-shore oceanic regions, could expect no CaCO_3 precipitation with ~~aqueous~~ NaOH, $[\text{Ca}^{2+}] + [\text{HCO}_3^-]$, or CaO addition at $\sim 500 \mu\text{mol kg}^{-1}$ for at least four days ~~if the alkalinity is added in an aqueous form~~. Thus far it is clear that the thresholds for heterogeneous precipitation (i.e., relevant if using mineral feedstock) versus pseudo-homogeneous precipitation (i.e., relevant if primarily using aqueous solutions for OAE in natural seawater) (Chen et al., 2005; Marion et al., 2009; Wolf et al., 2008) must be different, and may vary with different temperature, salinity, or even organic matter content (Moras et al., 2024; Suitner et al., 2026). Future studies could focus on comparing these precipitation thresholds by systematically changing parameters that influence the induction period before mineral precipitation.

4.2 Impact of ~~a~~Alkalinity on ~~o~~Organic ~~m~~Matter ~~c~~Cycling

4.2.1 Minimal ~~i~~Impacts of NaOH and $[\text{Ca}^{2+}]$ ~~±~~~~and~~ $[\text{HCO}_3^-]$ ~~a~~Additions

The Port of Los Angeles has a dynamic seasonal cycle that directly influences the carbon standing stock – driven by the seasonally dominant organisms within the water. The organic carbon biomass is highest in Spring at $300 \mu\text{gC L}^{-1}$, followed by $150 \mu\text{gC L}^{-1}$ in Summer, $100 \mu\text{gC L}^{-1}$ in Fall, and $80 \mu\text{gC L}^{-1}$ in the Winter (Connell et al., 2017). There is also seasonality in the types of organisms that are creating biomass and are thus responding to the alkalinity addition during our experiments.

The NaOH (starting $[\text{POC}] = 200 \mu\text{gC L}^{-1}$) and $[\text{Ca}^{2+}]$ ~~±~~~~and~~ $[\text{HCO}_3^-]$ (starting $[\text{POC}] = 400 \mu\text{gC L}^{-1}$) experiments were conducted in the Summer, when Diatoms make up more than half of the particulate biomass, followed by Picoeukaryotes in the LA Port region (Connell et al., 2017). The lack of impact on POC or Chl *a* production between the controls and treatments after NaOH and $[\text{Ca}^{2+}] + [\text{HCO}_3^-]$ addition ($\sim 500 \mu\text{mol kg}^{-1}$, pH_T 7.77 – 8.62) in this study is congruent with results seen from other Diatom specific studies (Gately et al., 2023; Oberlander et al., 2025) and natural community studies (Guo et al., 2025) where the resultant change in POC production was also negligible. Other studies have seen a decrease in Diatom concentration once the alkalinity addition has increased pH beyond 8.55 (Kousoulas et al., 2025; Oberlander et al., 2025) and note a decrease in Picoeukaryote concentration after NaOH addition (Ferderer et al.,

2022; Kousoulas et al., 2025; Oberlander et al., 2025). In this study, we do not have evidence to say if the species compositions shifted, however, these alkalinity additions did not statistically alter the POC concentrations when compared to the control (Fig. 3). Federer et al. (2022) saw POC decline after alkalinity addition, which they attributed to a possible alteration in the composition of the diatom community and increased heterotrophic bacterial activity in their alkalinity (NaOH and NaHCO₃) treatments.

Heterotrophic bacteria continuously recycle organic matter derived from primary production and thus will be affected by processes changing carbon biomass (Dittmar et al., 2021; Santinelli et al., 2024; Soong et al., 2020). Some of our results suggest that NaOH addition could change PIC and DOC production in waters incubated in the dark (See [Figure A2-Supplemental Information](#)), but this is not observed in the dark CaO or [Ca²⁺] + [HCO₃⁻] additions. In the non-shaded bottles of the [Ca²⁺] + [HCO₃⁻] experiments, there appear to be substantial amounts of DOC production, but statistically we cannot say if this is a result of the $\sim$ +500 $\mu\text{mol kg}^{-1}$ alkalinity enhancement. The observed heterogeneity may simply be caused by when the water was sampled. These results, when considered with existing literature suggest that more OAE impact studies should focus on changes in heterotrophic bacteria, metabolic reactions that utilize [HCO₃⁻], and recycling of DOC.

The pH in LA Harbor waters has a natural measured variability of 7.57-8.46 (Lyons and Birosik, 2007). Under the conditions of our incubations, the natural microbial populations may be acclimated to pH shifts of the order generated by the NaOH and [Ca²⁺] + [HCO₃⁻] additions ($\sim$ +500 $\mu\text{mol kg}^{-1}$, pH 7.77 – 8.62) in these experiments. It is not uncommon for coastal environments to experience pH oscillations of 1 or more pH units due to tidal fluctuations or metabolic processes such as photosynthesis and respiration. Hinga, (2002) reviewed 21 studies that assess the impact of pH on the growth rates of coastal marine phytoplankton. Some species were able to thrive at a wide pH range, but others were extremely sensitive to changes of <0.5 pH units. In LA Harbor there are also natural alkalinity changes experienced by the ecosystems, and therefore these water column ecosystems may be able to handle some extent of OAE. However, determining “safe” thresholds for OAE is a nuanced ecological and societal issue that requires further scientific consensus.

4.2.2 Potential Impact of CaO Addition

The impacts to Chl a and POC from the CaO addition when compared to the unamended conditions show different outcomes. Of the two experiments conducted around $\sim 500 \mu\text{mol kg}^{-1}$, one (01-27-25; $\text{measured } \Delta\text{TA} = 612 \pm 13 \mu\text{mol kg}^{-1}$) shows that Chl a was statistically lower. This trend was not seen in Chl a in the experiment conducted at $\sim 1000 \mu\text{mol kg}^{-1}$ (11-04-24; $\text{measured } \Delta\text{TA} = 923 \pm 12 \mu\text{mol kg}^{-1}$). POC production was reduced by $100 \mu\text{gC L}^{-1}$ after 4-days from the $\sim 500 \mu\text{mol kg}^{-1}$ CaO additions (but the trend is not statistically significant due to high initial variability). POC production is reduced by $200 \mu\text{gC L}^{-1}$ after the $\sim 1000 \mu\text{mol kg}^{-1}$ addition, *which is statistically significant*. This coincides with statistically different (lower) dissolved oxygen after a $\sim 1000 \mu\text{mol kg}^{-1}$ CaO addition as compared to the control. The impacts of CaO addition at $\sim 500 \mu\text{mol kg}^{-1}$ may not be statistically significant on LA Harbor organic matter cycling, but at $\sim 1000 \mu\text{mol kg}^{-1}$ the impacts do become statistically significant.

Other studies that test the impact of ocean liming – either simulated by CaO, Ca(OH)_2 , or even NaOH + CaCl_2 addition – have seen minimal effects on growth rates, grazing rate, and community composition when the resultant change has been below pH 9 (de Castro et al., 2025; Traboni et al., 2025). In our study, the starting pH_T of the seawater is around 7.68-7.85 and the CaO additions bring this pH_T to 8.57-8.91 (~ 500 – $1000 \mu\text{mol kg}^{-1}$ experiments). Several studies, that raised pH > 9 with CaO, have shown a decrease in organism quality (i.e., nutrient composition) and growth rate (Bhaumik et al., 2025) or decreased motility and even increased mortality (Camatti et al., 2024; Pedersen and Hansen, 2003). CaO addition, that reduces pCO_2 below $\sim 100 \mu\text{atm}$, could cause issues in primary production from the CO_2 limitation. This has been shown in coccolithophore growth (Bach et al., 2011; Faucher et al., 2025; Riebesell et al., 1993; Sett et al., 2014) and for other marine phytoplankton (Hansen, 2002; Pedersen and Hansen, 2003). These results draw attention to how reorganization of community composition (with smaller phytoplankton that are better adapted to lower pCO_2 availability (Chrachri et al., 2018; Flynn et al., 2012; Wolf-Gladrow and Riebesell, 1997)) could result from alkalinity additions. This in ~~turn~~ could reduce the transfer efficiency of POM to the deeper ocean (Boyd and Newton, 1995). In this study we do not incorporate a characterization of size classes, but future alkalinity enhancement studies should also try to incorporate an analysis of changes in particle size class, among a suite of other ecological impacts (Marx et al., 2025 Preprint).

Formatted: Font: 12 pt, Font color: Auto

5. Conclusions

Regional studies that systematically explore the impacts on biomass transfer but also characterize the changing biomass will be highly informative when considering the impacts of OAE on carbon partitioning. In our LA Harbor waters, POC sometimes ~~POC~~ increased or decreased over 4-day incubations. However, there was a consistent pattern of less POC production in CaO ~~-added~~-treatment bottles –lower by 100-200 $\mu\text{gC L}^{-1}$ compared to control samples; as was statistically lower at $\Omega_{\text{aragonite}} = 14$. Future studies should resolve changes in POC particle class size and leverage ‘omics tools to understand what parts of the organic matter cycle may be impacted by CaO addition. Studies should also draw attention to the impacts of increased alkalinity in regions where the amended water mass sinks below the euphotic zone and heterotrophic reactions may dominate carbon cycling.

Our experiments show that even factors that should increase the likelihood of CaCO_3 precipitation – supersaturation & greater nucleation sites – did not lead to CaCO_3 precipitation over a four-day incubation period at $\Omega_{\text{aragonite}} \leq 14$ in closed but photosynthetically active systems. Additionally, Chlorophyll as a proxy for the community of phytoplankton, did not show sensitivity to any of the three alkalinity types at an addition threshold of $\approx +500 \mu\text{mol kg}^{-1}$.

There may be other environmental parameters that influence PIC and POC production in concert with alkalinity enhancement. We recommend conducting longer duration experiments across a wide range of waters and ecosystems, especially regions that may experience smaller shifts in alkalinity. However, natural ecosystems that do experience more variability in alkalinity may be resilient to some OAE intervention.

Appendix A. Summary of data from “Dark-Bottles” during alkalinity enhancement experiments

Formatted: Font: Bold

Formatted: Font: Bold, Font color: Black

Formatted: Normal, Justified, Indent: Left: 0"




Figure A1: Summary of carbonate chemistry measured in “Dark Bottles”. Data in figure panels (First Row) Total Alkalinity $\mu\text{mol kg}^{-1}$, (Second Row) Dissolved Inorganic Carbon $\mu\text{mol kg}^{-1}$, and (Third Row) pH_T (as calculated on the Total Scale from the Total Alkalinity and DIC pair in CO2sys.v3 – detailed in Methods). Bottles that were devoid of light (deployed during the in-situ experiments) are shown in these figures; Day 0 Bottles from the normal light exposure experiments shown for reference (n=3). Day 2 and Day 4 bottles have an n=1 (Statistical analysis was not conducted on this data – only being presented for qualitative assessment). Treatments include Control (green), NaOH (red), $\text{CaCl}_2 + \text{NaHCO}_3$ (blue), and CaO (purple).




Figure A2: Summary of biogeochemical variables measured in “Dark Bottles”. Data in figure panels (First Row) Chl-a $\mu\text{g L}^{-1}$, (Second Row) Particulate Inorganic Carbon $\mu\text{g C L}^{-1}$, (Third Row) Particulate Organic Carbon $\mu\text{g C L}^{-1}$, and (Fourth Row) Dissolved Organic Carbon $\mu\text{mol kg}^{-1}$. Bottles that were devoid of light (deployed during the in-situ experiments) are shown in these figures; Day 0 Bottles from the normal light exposure experiments shown for reference ($n=3$). Day 2 and Day 4 bottles have an $n=1$ (Statistical analysis was not conducted on this data – only being presented for qualitative assessment). Treatments include Control (green), NaOH (red), $\text{CaCl}_2 + \text{NaHCO}_3$ (blue), and CaO (purple).

In the 4-day experiments we also included 4 bottles that received no light (2 controls and 2 treatments), while the remaining 18 bottles were subjected to the regular day-night cycle. A summary of the carbonate system manipulations can be seen in Figure A1. The biogeochemical changes to alkalinity additions are summarized in Figure A2. There were no replicate bottles on Day 2 and 4, hence, we did not conduct any statistical analysis on this dataset. We have included this dataset for qualitative results that may encourage further study of the impacts of alkalinity on dark metabolic reactions and organic matter cycling. There are no consistent trends in Chl a or particulate organic carbon concentrations across treatments. We see an increase of particulate

inorganic carbon and dissolved organic carbon in the NaOH treatment bottles relative to the control conditions, but we caution against making any conclusive judgements about the effective alkalinity on dark reactions from this dataset.

Appendix B. CaCO₃ precipitation after OAE

Table B1: Experimental conditions where CaCO₃ precipitation is NOT observed.

<u>Study</u>	<u>Feedstock</u>	<u>Media</u>	<u>Ω_{aragonite}</u>	<u>Temperature</u> (°C)	<u>Time</u> (days)
<u>Hartman et al. 2023</u>	<u>NaHCO₃</u>	<u>0.2 μm filtered SW</u>	<u>< 15</u>	<u>24</u>	<u>4</u>
	<u>+ Na₂CO₃</u>	<u>55 μm filtered SW</u>	<u>~ 6</u>	<u>24</u>	<u>4</u>
<u>Suitner et al. 2024</u>	<u>NaHCO₃</u>	<u>0.2 μm filtered SW</u>	<u>< 15</u>	<u>12-16</u>	<u>20</u>
	<u>+ Na₂CO₃</u>	<u>55 μm filtered SW</u>	<u>< 15</u>	<u>12-15</u>	<u>20</u>
<u>Moras et al. 2022</u>	<u>Na₂CO₃</u>	<u>0.2 μm filtered SW</u>	<u>~ 9</u>	<u>21</u>	<u>40</u>
		<u>0.2 μm filtered SW</u>	<u>~ 9</u>	<u>21</u>	<u>5</u>
		<u>+ Quartz crystals</u>			
<u>Hartman et al. 2023</u>	<u>NaOH</u>	<u>0.2 μm filtered SW</u>	<u><20</u>	<u>24</u>	<u>4</u>
		<u>55 μm filtered SW</u>	<u><20</u>	<u>24</u>	<u>4</u>
<u>Suitner et al. 2024</u>	<u>NaOH</u>	<u>0.2 μm filtered SW</u>	<u>< 15</u>	<u>11-13</u>	<u>25</u>
		<u>55 μm filtered SW</u>	<u>~19</u>	<u>10-11</u>	<u>25</u>
<u>Ringham et al. 2024</u>	<u>NaOH</u>	<u>Unfiltered SW</u>	<u>< 30</u>	<u>19-21</u>	<u>20-60</u>
<u>Moras et al. 2022</u>	<u>CaO</u>	<u>0.2 μm filtered SW;</u>	<u>5</u>	<u>21</u>	<u>40</u>
	<u>Ca(OH)₂</u>	<u>solid feedstock</u>	<u>4</u>	<u>21</u>	<u>30</u>

Formatted: Font: Not Bold, Italic
Formatted: Normal, Don't keep with next

Table B2: Experimental conditions where CaCO₃ precipitation is observed.

<u>Study</u>	<u>Feedstock</u>	<u>Media</u>	<u>$\Omega_{\text{aragonite}}$</u>	<u>Temperature (°C)</u>	<u>Onset of Precipitation (days)</u>
<u>Suitner et al. 2024</u>	<u>NaHCO₃ + Na₂CO₃</u>	<u>0.2 μm filtered SW</u>	<u>15-45</u>	<u>12-16</u>	<u>1-20</u>
<u>Moras et al. 2022</u>	<u>Na₂CO₃</u>	<u>0.2 μm filtered SW + Quartz crystals</u>	<u>~ 9</u>	<u>21</u>	<u>>5</u>
<u>Hartman et al. 2023</u>	<u>NaOH</u>	<u>0.2 μm filtered SW</u>	<u>$\sim 16-19$</u>	<u>24</u>	<u>0-1</u>
		<u>55 μm filtered SW</u>	<u>$\sim 15-17$</u>	<u>24</u>	<u>0-1</u>
<u>Suitner et al. 2024</u>	<u>NaOH</u>	<u>0.2 μm filtered SW</u>	<u>$\geq 15-30$</u>	<u>11-13</u>	<u>0-25</u>
		<u>55 μm filtered SW</u>	<u>$\geq 19-30$</u>	<u>10-11</u>	<u>1-15</u>
<u>Ringham et al. 2024</u>	<u>NaOH</u>	<u>Unfiltered SW</u>	<u>> 30</u>	<u>19-21</u>	<u>1</u>
<u>Hashim et al. 2025</u>	<u>NaOH</u>	<u>Unfiltered SW</u>	<u>10-30</u>	<u>27</u>	<u><1</u>
<u>Moras et al. 2022</u>	<u>CaO</u>	<u>0.2 μm filtered SW; solid feedstock</u>	<u>~ 7</u>	<u>21</u>	<u>0.2</u>
	<u>Ca(OH)₂</u>		<u>~ 7</u>	<u>21</u>	<u>0.2</u>
<u>Moras et al. 2024</u>	<u>Mg(OH)₂</u>	<u>0.2 μm filtered SW; $\leq 63 \mu\text{m}$ feedstock</u>	<u>8-10</u>	<u>24</u>	<u>4-8</u>
		<u>0.2 μm filtered SW; $>180 \mu\text{m}$ feedstock</u>	<u>8-10</u>		<u>2-4</u>
		<u>0.2 μm filtered SW; 63-180 μm feedstock; S =20</u>	<u>8-10</u>		<u>2-4</u>
		<u>0.2 μm filtered SW; 63-180 μm feedstock; S =28</u>			<u>6-8</u>
		<u>0.2 μm filtered SW; 63-180 μm feedstock; S = 36</u>			<u>10-14</u>

*Seawater abbreviated to SW; Salinity abbreviated to S.

These tables summarize the conditions under which CaCO₃ precipitation was and was not observed, as discussed in Section 4.1 Absence of secondary CaCO₃ precipitation.

Formatted: Justified, Indent: First line: 0"

Data Availability

Data are publicly available at <https://doi.org/10.5281/zenodo.20059856>.

Author Contributions

RW and WB designed the experiments with consultation from NR. RW conducted these experiments with the help of DR, RA, and EJ. RW was responsible for sample analysis, data curation, and formal statistical analysis with consultation from NR and WB. Funding was acquired by WB. The manuscript was written and edited by RW with contributions from WB.

Competing Interests

WB is the co-founder of Calcareia, Inc. The authors declare that they have no other competing interests.

Acknowledgements

We would like to thank the USC Wrigley Institute of Environment and Sustainability, AltaSea, and Calcareia (Pierre Forin), for access to their laboratory and dock resources. We acknowledge our presence on the ancestral and unceded territory of the Kizh (Gabrieleño) people, and we pay our respect to the members and elders of these communities. We also thank Doug Capone, Myrna Jacobson, Sarah Feakins, and the Department of Environmental Engineering at the University of Southern California for access to their instruments to make Chl-a and dissolved organic carbon measurements. Additionally, we thank the two anonymous reviewers and the associate editor, Perran Cook, for their efforts in reviewing this manuscript.

Financial Support

This research was conducted while RW was a recipient of the Wrigley Institute Graduate Fellowship in 2023. The research was funded by the Lott Innovation Award and the Faculty Innovation Award (to WB) through the USC Wrigley Institute for Environmental Studies (WIES). We acknowledge an NSF-REU award to USC WIES for support of RA (2024) and a Zinsmeyer Summer Research Award to EJM (2023) through grant no. OCE-2244583.

References

- Anderson, L. A.: On the hydrogen and oxygen content of marine phytoplankton, Deep-Sea Research Part I, 42, 1675–1680, [https://doi.org/10.1016/0967-0637\(95\)00072-E](https://doi.org/10.1016/0967-0637(95)00072-E), 1995.
- Archer, D. E.: An atlas of the distribution of calcium carbonate in sediments of the deep sea, Global Biogeochem. Cycles, 10, 159–174, <https://doi.org/10.1029/95GB03016>, 1996.
- Bach, L. T.: The additionality problem of ocean alkalinity enhancement, Biogeosciences, 21, 261–277, <https://doi.org/10.5194/bg-21-261-2024>, 2024.
- Bach, L. T., Riebesell, U., and Schulz, K. G.: Distinguishing between the effects of ocean acidification and ocean carbonation in the coccolithophore *Emiliania huxleyi*, Limnol. Oceanogr., 56, 2040–2050, <https://doi.org/10.4319/lo.2011.56.6.2040>, 2011.
- Bach, L. T., Gill, S. J., Rickaby, R. E. M., Gore, S., and Renforth, P.: CO₂ Removal With Enhanced Weathering and Ocean Alkalinity Enhancement: Potential Risks and Co-benefits for Marine Pelagic Ecosystems, Frontiers in Climate, 1, <https://doi.org/10.3389/fclim.2019.00007>, 2019.
- Bach, L. T., Ferderer, A. J., LaRoche, J., and Schulz, K. G.: Technical note: Ocean Alkalinity Enhancement Pelagic Impact Intercomparison Project (OAEPiIP), Biogeosciences, 21, 3665–3676, <https://doi.org/10.5194/bg-21-3665-2024>, 2024.
- Barker, S., Archer, D., Booth, L., Elderfield, H., Henderiks, J., and Rickaby, R. E. M.: Globally increased pelagic carbonate production during the Mid-Brunhes dissolution interval and the CO₂ paradox of MIS 11, Quat. Sci. Rev., 25, 3278–3293, <https://doi.org/10.1016/j.quascirev.2006.07.018>, 2006.
- Berelson, W. M., Balch, W. M., Najjar, R., Feely, R. A., Sabine, C., and Lee, K.: Relating estimates of CaCO₃ production, export, and dissolution in the water column to measurements of CaCO₃ rain into sediment traps and dissolution on the sea floor: A revised global carbonate budget, Global Biogeochem. Cycles, 21, 1–15, <https://doi.org/10.1029/2006GB002803>, 2007.
- Berner, R. A.: The role of magnesium in the crystal growth of calcite and aragonite from sea water, Geochim. Cosmochim. Acta, 39, 489–504, [https://doi.org/10.1016/0016-7037\(75\)90102-7](https://doi.org/10.1016/0016-7037(75)90102-7), 1975.
- Berner, R. A. and Berner, E. K.: Silicate Weathering and Climate, in: Tectonic Uplift and Climate Change, Springer US, Boston, MA, 353–365, https://doi.org/10.1007/978-1-4615-5935-1_15, 1997.
- Berner, R. A., Lasaga, A. C., and Garrels, R. M.: The carbonate-silicate geochemical cycle and its effect on atmospheric carbon dioxide over the past 100 million years, Am. J. Sci., 283, 641–683, <https://doi.org/10.2475/ajs.283.7.641>, 1983.
- Bhaumik, A., Faucher, G., Henning, M., Meunier, C. L., and Boersma, M.: Prey dynamics as a buffer: enhancing copepod resilience to ocean alkalinity enhancement, Environmental Research Letters, 20, <https://doi.org/10.1088/1748-9326/adaa8c>, 2025.

- 860 Bialonski, S., Caron, D. A., Schloen, J., Feudel, U., Kantz, H., and Moorthi, S. D.: Phytoplankton
861 dynamics in the Southern California Bight indicate a complex mixture of transport and biology,
862 *J. Plankton Res.*, 38, 1077–1091, <https://doi.org/10.1093/plankt/fbv122>, 2016.
- 863 Boyd, P. and Newton, P.: Evidence of the potential influence of planktonic community structure
864 on the interannual variability of particulate organic carbon flux, *Deep Sea Research Part I:
865 Oceanographic Research Papers*, 42, 619–639, [https://doi.org/10.1016/0967-0637\(95\)00017-Z](https://doi.org/10.1016/0967-0637(95)00017-Z),
866 1995.
- 867 Cai, W.-J. and Jiao, N.: Wastewater alkalinity addition as a novel approach for ocean negative
868 carbon emissions, *The Innovation*, 3, 100272, <https://doi.org/10.1016/j.xinn.2022.100272>, 2022.
- 869 Camatti, E., Valsecchi, S., Caserini, S., Barbaccia, E., Santinelli, C., Basso, D., and Azzellino, A.:
870 Short-term impact assessment of ocean liming: A copepod exposure test, *Mar. Pollut. Bull.*, 198,
871 115833, <https://doi.org/10.1016/j.marpolbul.2023.115833>, 2024.
- 872 Caron, D. A., Connell, P. E., Schaffner, R. A., Schnetzer, A., Fuhrman, J. A., Countway, P. D.,
873 and Kim, D. Y.: Planktonic food web structure at a coastal time-series site: I. Partitioning of
874 microbial abundances and carbon biomass, *Deep Sea Research Part I: Oceanographic Research
875 Papers*, 121, 14–29, <https://doi.org/10.1016/j.dsr.2016.12.013>, 2017.
- 876 Carter, C. M., Ross, A. H., Schiel, D. R., Howard-Williams, C., and Hayden, B.: In situ
877 microcosm experiments on the influence of nitrate and light on phytoplankton community
878 composition, *J. Exp. Mar. Biol. Ecol.*, 326, 1–13, <https://doi.org/10.1016/j.jembe.2005.05.006>,
879 2005.
- 880 Caserini, S., Pagano, D., Campo, F., Abbà, A., De Marco, S., Righi, D., Renforth, P., and Grosso,
881 M.: Potential of Maritime Transport for Ocean Liming and Atmospheric CO₂ Removal, *Frontiers
882 in Climate*, 3, 1–18, <https://doi.org/10.3389/fclim.2021.575900>, 2021.
- 883 de Castro, I., Ribeiro, S. C., Louvado, A., Gomes, N. C. M., Cachão, M., Brito de Azevedo, E.,
884 and Barcelos e Ramos, J.: Ocean liming effect on a North Atlantic microbial community:
885 changes in composition and rates, *Front. Mar. Sci.*, 12,
886 <https://doi.org/10.3389/fmars.2025.1602158>, 2025.
- 887 Chen, T., Neville, A., and Yuan, M.: Calcium carbonate scale formation—assessing the initial
888 stages of precipitation and deposition, *J. Pet. Sci. Eng.*, 46, 185–194,
889 <https://doi.org/10.1016/j.petrol.2004.12.004>, 2005.
- 890 Chou, H.-T.: On the dilution of liquid waste in ship's wake, *J. Mar. Sci. Technol.*, 149–154,
891 <https://doi.org/10.1126/science.155.3770.1621>, 1996.
- 892 Chrachri, A., Hopkinson, B. M., Flynn, K., Brownlee, C., and Wheeler, G. L.: Dynamic changes
893 in carbonate chemistry in the microenvironment around single marine phytoplankton cells, *Nat.
894 Commun.*, 9, 74, <https://doi.org/10.1038/s41467-017-02426-y>, 2018.
- 895 de Coninck, H., Revi, A., Babiker, M., Bertoldi, P., Buckeridge, M., Cartwright, A., Dong, W.,
896 Ford, J., Fuss, S., Hourcade, J.-C., Ley, D., Mechler, R., Newman, P., Revokatova, A., Schultz,

- 897 K., Steg, L., and Sugiyama, T.: Strengthening and Implementing the Global Response In: Global
898 Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above
899 pre-industrial levels and related global greenhouse gas emission pathways, in the context of
900 strengthening the global response to the threat of climate change, sustainable development, and
901 efforts to eradicate poverty, in: Global Warming of 1.5°C, edited by: Masson-Delmotte, V., Zhai,
902 P., Portner, H.-O., Roberts, D., Skea, J., Shukla, P. R., Pirani, A., Moufouma-Okia, W., Pean, C.,
903 Pidcock, R., Connors, S., Matthews, J. B. R., Chen, Y., Zhou, X., Gomis, M. I., Lonny, E.,
904 Maycock, T., Tignor, M., and Waterfield, T., Cambridge University Press, Cambridge, UK and
905 New York, NY, USA, 313–444, <https://doi.org/10.1017/9781009157940.006>, 2018.
- 906 Connell, P. E., Campbell, V., Gellene, A. G., Hu, S. K., and Caron, D. A.: Planktonic food web
907 structure at a coastal time-series site: II. Spatiotemporal variability of microbial trophic activities,
908 Deep Sea Research Part I: Oceanographic Research Papers, 121, 210–223,
909 <https://doi.org/10.1016/j.dsr.2017.01.007>, 2017.
- 910 Daehler, C. C. and Strong, D. R.: Can You Bottle Nature? The Roles of Microcosms in
911 Ecological Research, Ecology, 77, 663–664, <https://doi.org/10.2307/2265487>, 1996.
- 912 Dickson, A. G.: Thermodynamics of the dissociation of boric acid in synthetic seawater from
913 273.15 to 318.15 K, Deep Sea Research Part A. Oceanographic Research Papers, 37, 755–766,
914 [https://doi.org/10.1016/0198-0149\(90\)90004-F](https://doi.org/10.1016/0198-0149(90)90004-F), 1990.
- 915 Dickson, A. G. and Millero, F. J.: A comparison of the equilibrium constants for the dissociation
916 of carbonic acid in seawater media, Deep Sea Research Part A. Oceanographic Research Papers,
917 34, 1733–1743, [https://doi.org/10.1016/0198-0149\(87\)90021-5](https://doi.org/10.1016/0198-0149(87)90021-5), 1987.
- 918 Dickson, A. G. , Sabine, C. L. , and Christian, J. Robert.: Guide to best practices for ocean CO₂
919 measurements, North Pacific Marine Science Organization, 2007.
- 920 Dittmar, T., Lennartz, S. T., Buck-Wiese, H., Hansell, D. A., Santinelli, C., Vanni, C., Blasius, B.,
921 and Hehemann, J.-H.: Enigmatic persistence of dissolved organic matter in the ocean, Nat. Rev.
922 Earth Environ., 2, 570–583, <https://doi.org/10.1038/s43017-021-00183-7>, 2021.
- 923 Dong, S., Berelson, W. M., Forin, P., Gutierrez, M., Carroll, D., Menemenlis, D., Kyi, A. Y., and
924 Adkins, J. F.: Potential of CO₂ sequestration through accelerated weathering of limestone on
925 ships, Sci. Adv., 11, <https://doi.org/10.1126/sciadv.adr7250>, 2025.
- 926 Dunne, J. P., Sarmiento, J. L., and Gnanadesikan, A.: A synthesis of global particle export from
927 the surface ocean and cycling through the ocean interior and on the seafloor, Global
928 Biogeochem. Cycles, 21, <https://doi.org/10.1029/2006GB002907>, 2007.
- 929 Eisaman, M. D., Geilert, S., Renforth, P., Bastianini, L., Campbell, J., Dale, A. W., Foteinis, S.,
930 Grasse, P., Hawrot, O., Löscher, C. R., Rau, G. H., and Rønning, J.: Assessing the technical
931 aspects of ocean-alkalinity-enhancement approaches, State of the Planet, 2-oae2023, 1–29,
932 <https://doi.org/10.5194/sp-2-oae2023-3-2023>, 2023.
- 933 Falkowski, P., Scholes, R. J., Boyle, E., Canadell, J., Canfield, D., Elser, J., Gruber, N., Hibbard,
934 K., Höglberg, P., Linder, S., Mackenzie, F. T., Moore III, B., Pedersen, T., Rosenthal, Y.,

- Seitzinger, S., Smetacek, V., and Steffen, W.: The Global Carbon Cycle: A Test of Our Knowledge of Earth as a System, *Science* (1979)., 290, 291–296, <https://doi.org/10.1126/science.290.5490.291>, 2000.
- Faucher, G., Haunost, M., Paul, A. J., Tietz, A. U. C., and Riebesell, U.: Growth response of *Emiliania huxleyi* to ocean alkalinity enhancement, *Biogeosciences*, 22, 405–415, <https://doi.org/10.5194/bg-22-405-2025>, 2025.
- Ferderer, A., Chase, Z., Kennedy, F., Schulz, K. G., and Bach, L. T.: Assessing the influence of ocean alkalinity enhancement on a coastal phytoplankton community, *Biogeosciences*, 19, 5375–5399, <https://doi.org/10.5194/bg-19-5375-2022>, 2022.
- Flynn, K. J., Blackford, J. C., Baird, M. E., Raven, J. A., Clark, D. R., Beardall, J., Brownlee, C., Fabian, H., and Wheeler, G. L.: Changes in pH at the exterior surface of plankton with ocean acidification, *Nat. Clim. Chang.*, 2, 510–513, <https://doi.org/10.1038/nclimate1489>, 2012.
- Fraser, L. H. and Keddy, P.: The role of experimental microcosms in ecological research, *Trends Ecol. Evol.*, 12, 478–481, [https://doi.org/10.1016/S0169-5347\(97\)01220-2](https://doi.org/10.1016/S0169-5347(97)01220-2), 1997.
- Gately, J. A., Kim, S. M., Jin, B., Brzezinski, M. A., and Iglesias-Rodriguez, M. D.: Coccolithophores and diatoms resilient to ocean alkalinity enhancement: A glimpse of hope?, *Sci. Adv.*, 9, <https://doi.org/10.1126/sciadv.adg6066>, 2023.
- Giesy, J. P. : Microcosms in ecological research : selected papers from a symposium held at Augusta, Georgia, November 8-10, 1978, Technical Information Center, U.S. Dept. of Energy ; Available from National Technical Information Service, U.S. Dept. of Commerce, 1110 pp., 1980.
- Guo, J. A., Strzepek, R. F., Yuan, Z., Swadling, K. M., Townsend, A. T., Achterberg, E. P., Browning, T. J., and Bach, L. T.: Effects of ocean alkalinity enhancement on plankton in the Equatorial Pacific, *Commun. Earth Environ.*, 6, <https://doi.org/10.1038/s43247-025-02248-7>, 2025.
- Hansen, P.: Effect of high pH on the growth and survival of marine phytoplankton: implications for species succession, *Aquatic Microbial Ecology*, 28, 279–288, <https://doi.org/10.3354/ame028279>, 2002.
- Hartmann, J., Suitner, N., Lim, C., Schneider, J., Marin-Samper, L., Arístegui, J., Renforth, P., Taucher, J., and Riebesell, U.: Stability of alkalinity in ocean alkalinity enhancement (OAE) approaches - consequences for durability of CO₂ storage, *Biogeosciences*, 20, 781–802, <https://doi.org/10.5194/bg-20-781-2023>, 2023.
- Hashim, M. S., Marx, L., Klein, F., Dean, C. L., Burdige, E., Hayden, M., McCorkle, D. C., and Subhas, A. V.: Mineral formation during shipboard ocean alkalinity enhancement experiments in the North Atlantic, *Biogeosciences*, 22, 7149–7165, <https://doi.org/10.5194/bg-22-7149-2025>, 2025.

- 971 Hinga, K.: Effects of pH on coastal marine phytoplankton, *Mar. Ecol. Prog. Ser.*, 238, 281–300,
972 <https://doi.org/10.3354/meps238281>, 2002.
- 973 Holland, H. D. : The chemistry of the atmosphere and oceans, Wiley, 351 pp., 1978.
- 974 Iglesias-Rodríguez, M. D., Rickaby, R. E. M., Singh, A., and Gately, J. A.: Laboratory
975 experiments in ocean alkalinity enhancement research, *State of the Planet*, 2-oae2023, 1–18,
976 <https://doi.org/10.5194/sp-2-oae2023-5-2023>, 2023.
- 977 Ives, A. R., Foufopoulos, J., Klopfer, E. D., Klug, J. L., and Palmer, T. M.: Bottle or Big-Scale
978 Studies: How do we do Ecology?, *Ecology*, 77, 681–685, <https://doi.org/10.2307/2265491>, 1996.
- 979 Kheshgi, H. S.: Sequestering atmospheric carbon dioxide by increasing ocean alkalinity, *Energy*,
980 20, 915–922, [https://doi.org/10.1016/0360-5442\(95\)00035-F](https://doi.org/10.1016/0360-5442(95)00035-F), 1995.
- 981 Kousoulas, K., Ferderer, A., Eriksen, R., and Bach, L. T.: Winners and losers under hydroxide-
982 based ocean alkalinity enhancement in a Tasmanian plankton community, *J. Phycol.*,
983 <https://doi.org/10.1111/jpy.70052>, 2025.
- 984 de la Broise, D. and Palenik, B.: Immersed in situ microcosms: A tool for the assessment of
985 pollution impact on phytoplankton, *J. Exp. Mar. Biol. Ecol.*, 341, 274–281,
986 <https://doi.org/10.1016/J.JEMBE.2006.10.045>, 2007.
- 987 Lawton, J. H., Naeem, S., Woodfin, R. M., Brown, V. K., Gange, A., Godfray, H. J. C., Heads, P.
988 A., Lawler, S., Magda, D., Thomas, C. D., Thompson, L. J., and Young, S.: The Ecotron: a
989 controlled environmental facility for the investigation of population and ecosystem processes,
990 *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 341, 181–194, <https://doi.org/10.1098/rstb.1993.0102>,
991 1993.
- 992 Lee, K., Kim, T.-W., Byrne, R. H., Millero, F. J., Feely, R. A., and Liu, Y.-M.: The universal ratio
993 of boron to chlorinity for the North Pacific and North Atlantic oceans, *Geochim. Cosmochim.*
994 *Acta*, 74, 1801–1811, <https://doi.org/10.1016/j.gca.2009.12.027>, 2010.
- 995 Lunstrum, A. and Berelson, W.: CaCO₃ dissolution in carbonate-poor shelf sands increases with
996 ocean acidification and porewater residence time, *Geochim. Cosmochim. Acta*, 329, 168–184,
997 <https://doi.org/10.1016/j.gca.2022.04.031>, 2022.
- 998 Lunstrum, A., Hartmann, J., Moras, C., and Keller, D.: A framework for understanding efficiency
999 losses of Ocean Alkalinity Enhancement, <https://doi.org/10.70212/cdrxiv.2026517.v1>, 2026.
- 1000 Lyons, M. J. and Birosik, S.: WATER QUALITY IN THE DOMINGUEZ CHANNEL AND LOS
1001 ANGELES/LONG BEACH HARBOR WATERSHED MANAGEMENT AREA , 2007.
- 1002 Marion, G. M., Millero, F. J., and Feistel, R.: Precipitation of solid phase calcium carbonates and
1003 their effect on application of seawater SA-T-P models, *Ocean Science*, 5, 285–291,
1004 <https://doi.org/10.5194/os-5-285-2009>, 2009.
- 1005 Marx, L., Rheuban, J., McCorkle, D., Murray, C., Guo, Y., Wang, Z., Michel, A., Chen, K., Kim,
1006 H., and Subhas, A.: Development of the ecological activity index as an integrative ecosystem

- assessment and monitoring asset for ocean alkalinity enhancement,
<https://doi.org/10.21203/rs.3.rs-6371725/v1>, 28 May 2025.
- Moras, C. A., Bach, L. T., Cyronak, T., Joannes-Boyau, R., and Schulz, K. G.: Ocean alkalinity enhancement - avoiding runaway CaCO_3 precipitation during quick and hydrated lime dissolution, *Biogeosciences*, 19, 3537–3557, <https://doi.org/10.5194/bg-19-3537-2022>, 2022.
- Moras, C. A., Cyronak, T., Bach, L. T., Joannes-Boyau, R., and Schulz, K. G.: Effects of grain size and seawater salinity on magnesium hydroxide dissolution and secondary calcium carbonate precipitation kinetics: implications for ocean alkalinity enhancement, *Biogeosciences*, 21, 3463–3475, <https://doi.org/10.5194/bg-21-3463-2024>, 2024.
- Morse, J. W. and He, S.: Influences of T, S and PCO_2 on the pseudo-homogeneous precipitation of CaCO_3 from seawater: implications for whiting formation, *Mar. Chem.*, 41, 291–297, [https://doi.org/10.1016/0304-4203\(93\)90261-L](https://doi.org/10.1016/0304-4203(93)90261-L), 1993.
- Morse, J. W., Arvidson, R. S., and Lüttge, A.: Calcium Carbonate Formation and Dissolution, *Chem. Rev.*, 107, 342–381, <https://doi.org/10.1021/cr050358j>, 2007.
- Mucci, A. and Morse, J. W.: The incorporation of Mg^{2+} and Sr^{2+} into calcite overgrowths: influences of growth rate and solution composition, *Geochim. Cosmochim. Acta*, 47, 217–233, [https://doi.org/10.1016/0016-7037\(83\)90135-7](https://doi.org/10.1016/0016-7037(83)90135-7), 1983.
- National Academies of Sciences, E. and M.: A Research Strategy for Ocean-based Carbon Dioxide Removal and Sequestration, National Academies Press, Washington, DC, 1–308 pp., <https://doi.org/10.17226/26278>, 2022.
- Naviaux, J. D., Subhas, A. V., Dong, S., Rollins, N. E., Liu, X., Byrne, R. H., Berelson, W. M., and Adkins, J. F.: Calcite dissolution rates in seawater: Lab vs. in-situ measurements and inhibition by organic matter, *Mar. Chem.*, 215, 103684, <https://doi.org/10.1016/j.marchem.2019.103684>, 2019.
- Nguyen Dang, D., Gascoin, S., Zanibellato, A., G. Da Silva, C., Lemoine, M., Riffault, B., Sabot, R., Jeannin, M., Chateigner, D., and Gil, O.: Role of Brucite Dissolution in Calcium Carbonate Precipitation from Artificial and Natural Seawaters, *Cryst. Growth Des.*, 17, 1502–1513, <https://doi.org/10.1021/acs.cgd.6b01305>, 2017.
- Oberlander, J. L., Burke, M. E., London, C. A., and Macintyre, H. L.: Assessing the impacts of simulated ocean alkalinity enhancement on viability and growth of nearshore species of phytoplankton, *Biogeosciences*, 22, 499–512, <https://doi.org/10.5194/bg-22-499-2025>, 2025.
- Parsons, T. R.: The Future of Controlled Ecosystem Enclosure Experiments, in: *Marine mesocosms: Biological and Chemical Research in Experimental Ecosystems*, edited by: Grice, G. D. and Reeve, M. R., Springer-Verlag, New York, 411–418, 1982.
- Paul, A. J., Haunost, M., Goldenberg, S. U., Hartmann, J., Sánchez, N., Schneider, J., Suitner, N., and Riebesell, U.: Ocean alkalinity enhancement in an open ocean ecosystem: Biogeochemical

- 1043 responses and carbon storage durability, *EGUsphere*, 2024, 1–31,
1044 <https://doi.org/10.5194/egusphere-2024-417>, 2024.
- 1045 Pedersen, F. and Hansen, P.: Effects of high pH on a natural marine planktonic community, *Mar.*
1046 *Ecol. Prog. Ser.*, 260, 19–31, <https://doi.org/10.3354/meps260019>, 2003.
- 1047 Penman, D. E., Caves Rugenstein, J. K., Ibarra, D. E., and Winnick, M. J.: Silicate weathering as
1048 a feedback and forcing in Earth’s climate and carbon cycle, *Earth. Sci. Rev.*, 209, 103298,
1049 <https://doi.org/10.1016/j.earscirev.2020.103298>, 2020.
- 1050 Perez, F. F. and Fraga, F.: Association constant of fluoride and hydrogen ions in seawater, *Mar.*
1051 *Chem.*, 21, 161–168, [https://doi.org/10.1016/0304-4203\(87\)90036-3](https://doi.org/10.1016/0304-4203(87)90036-3), 1987.
- 1052 Pierrot, D., Epitalon, J. M., Orr, J. C., Lewis, E., and Wallace, D. W. R.: MS Excel program
1053 developed for CO₂ system calculations – version 3.0, 2021.
- 1054 Pukelsheim, F.: The Three Sigma Rule, *Am. Stat.*, 48, 88–91,
1055 <https://doi.org/10.1080/00031305.1994.10476030>, 1994.
- 1056 Rau, G. H. and Caldeira, K.: Enhanced carbonate dissolution: A means of sequestering waste
1057 CO₂ as ocean bicarbonate, *Energy Convers. Manag.*, 40, 1803–1813,
1058 [https://doi.org/10.1016/S0196-8904\(99\)00071-0](https://doi.org/10.1016/S0196-8904(99)00071-0), 1999.
- 1059 Renforth, P.: Perspective: The European Union is not prepared for geochemical carbon dioxide
1060 removal*, *Nº*, 37–72 pp., 2025.
- 1061 Renforth, P. and Henderson, G.: Assessing ocean alkalinity for carbon sequestration, *Reviews of*
1062 *Geophysics*, 55, 636–674, <https://doi.org/10.1002/2016RG000533>, 2017.
- 1063 Ridgwell, A. and Zeebe, R.: The role of the global carbonate cycle in the regulation and
1064 evolution of the Earth system, *Earth Planet. Sci. Lett.*, 234, 299–315,
1065 <https://doi.org/10.1016/j.epsl.2005.03.006>, 2005.
- 1066 Riebesell, U., Wolf-Gladrow, D. A., and Smetacek, V.: Carbon dioxide limitation of marine
1067 phytoplankton growth rates, *Nature*, 361, 249–251, <https://doi.org/10.1038/361249a0>, 1993.
- 1068 Riebesell, U., Zondervan, I., Rost, B., Tortell, P. D., Zeebe, R. E., and Morel, F. M. M.: Reduced
1069 calcification of marine plankton in response to increased atmospheric CO₂, *Nature*, 407, 364–
1070 367, <https://doi.org/10.1038/35030078>, 2000.
- 1071 Riebesell, U., Körtzinger, A., and Oschlies, A.: Sensitivities of marine carbon fluxes to ocean
1072 change, *Proceedings of the National Academy of Sciences*, 106, 20602–20609,
1073 <https://doi.org/10.1073/pnas.0813291106>, 2009.
- 1074 Riebesell, U., Lee, K., Nejstgaard, J. C., and Fisheries, I.: Pelagic mesocosms Part 2 :
1075 Experimental design of perturbation experiments Pelagic mesocosms, 2010.
- 1076 Ringham, M. C., Hirtle, N., Shaw, C., Lu, X., Herndon, J., Carter, B. R., and Eisaman, M. D.: An
1077 assessment of ocean alkalinity enhancement using aqueous hydroxides: kinetics, efficiency, and

- precipitation thresholds, *Biogeosciences*, 21, 3551–3570, <https://doi.org/10.5194/bg-21-3551-2024>, 2024.
- Santinelli, C., Valsecchi, S., Retelletti Brogi, S., Bachi, G., Checcucci, G., Guerrazzi, M., Camatti, E., Caserini, S., Azzellino, A., and Basso, D.: Ocean liming effects on dissolved organic matter dynamics, *Biogeosciences*, 21, 5131–5141, <https://doi.org/10.5194/bg-21-5131-2024>, 2024.
- Savoie, A. M., Ringham, M., Torres Sanchez, C., Carter, B. R., Dougherty, S., Feely, R. A., Hegeman, D., Herndon, J., Khangaonkar, T., Loretz, J., Minck, T., Pelman, T., Premathilake, L., Subban, C., Vance, J., and Ward, N. D.: Novel field trial for ocean alkalinity enhancement using electrochemically derived aqueous alkalinity, *Frontiers in Environmental Engineering*, 4, <https://doi.org/10.3389/fenv.2025.1641277>, 2025.
- Sett, S., Bach, L. T., Schulz, K. G., Koch-Klavsén, S., Lebrato, M., and Riebesell, U.: Temperature Modulates Coccolithophorid Sensitivity of Growth, Photosynthesis and Calcification to Increasing Seawater pCO₂, *PLoS One*, 9, e88308, <https://doi.org/10.1371/journal.pone.0088308>, 2014.
- Soong, J. L., Fuchslueger, L., Marañón-Jimenez, S., Torn, M. S., Janssens, I. A., Penuelas, J., and Richter, A.: Microbial carbon limitation: The need for integrating microorganisms into our understanding of ecosystem carbon cycling, *Glob. Chang. Biol.*, 26, 1953–1961, <https://doi.org/10.1111/gcb.14962>, 2020.
- Stachowski-Haberkorn, S., Becker, B., Marie, D., Haberkorn, H., Coroller, L., and de la Broise, D.: Impact of Roundup on the marine microbial community, as shown by an in situ microcosm experiment, *Aquatic Toxicology*, 89, 232–241, <https://doi.org/10.1016/j.aquatox.2008.07.004>, 2008.
- Subhas, A. V., Rollins, N. E., Berelson, W. M., Dong, S., Erez, J., and Adkins, J. F.: A novel determination of calcite dissolution kinetics in seawater, *Geochim. Cosmochim. Acta*, 170, 51–68, <https://doi.org/10.1016/j.gca.2015.08.011>, 2015.
- Subhas, A. V., Marx, L., Reynolds, S., Flohr, A., Mawji, E. W., Brown, P. J., and Cael, B. B.: Microbial ecosystem responses to alkalinity enhancement in the North Atlantic Subtropical Gyre, *Frontiers in Climate*, 4, <https://doi.org/10.3389/fclim.2022.784997>, 2022.
- Suitner, N., Faucher, G., Lim, C., Schneider, J., Moras, C. A., Riebesell, U., and Hartmann, J.: Ocean alkalinity enhancement approaches and the predictability of runaway precipitation processes: results of an experimental study to determine critical alkalinity ranges for safe and sustainable application scenarios, *Biogeosciences*, 21, 4587–4604, <https://doi.org/10.5194/bg-21-4587-2024>, 2024.
- Suitner, N., Hartmann, J., Varliero, S., Faucher, G., Suessle, P., and Moras, C. A.: Surface area and Ω -aragonite oversaturation as controls of the runaway precipitation process in ocean alkalinity enhancement, *Biogeosciences*, 23, 3965–3980, <https://doi.org/10.5194/bg-23-3965-2026>, 2026.

- 1116 Sulpis, O., Boudreau, B. P., Mucci, A., Jenkins, C., Trossman, D. S., Arbic, B. K., and Key, R.
 1117 M.: Current CaCO_3 dissolution at the seafloor caused by anthropogenic CO_2 , *Proc. Natl. Acad.*
 1118 *Sci. U. S. A.*, 115, 11700–11705, <https://doi.org/10.1073/pnas.1804250115>, 2018.
- 1119 Traboni, C., Nocera, A. C., Romano, F., Courboulès, J., Chantzaras, C., Magiopoulos, I.,
 1120 Varliero, S., Basso, D., and Pitta, P.: Plankton do not care: Minimal effects of ocean liming on
 1121 plankton growth and grazing in the Eastern Mediterranean, *Limnol. Oceanogr.*,
 1122 <https://doi.org/10.1002/lno.70136>, 2025.
- 1123 (UNFCCC) United Nations Framework Convention on Climate Change: NDC Synthesis Report,
 1124 2022.
- 1125 U.S. EPA: Standard Operating Procedure for In Vitro Determination of Chlorophyll a in
 1126 Freshwater Phytoplankton by Fluorescence LG405, 2021.
- 1127 Varliero, S., Buono, A., Caserini, S., Raos, G., and Macchi, P.: Chemical Aspect of Ocean
 1128 Liming for CO_2 Removal: Dissolution Kinetics of Calcium Hydroxide in Seawater, *ACS*
 1129 *Engineering Au*, 4, 422–431, <https://doi.org/10.1021/acsengineeringau.4c00008>, 2024.
- 1130 Walker, J. C. G., Hays, P. B., and Kasting, J. F.: A negative feedback mechanism for the long-
 1131 term stabilization of Earth’s surface temperature, *J. Geophys. Res. Oceans*, 86, 9776–9782,
 1132 <https://doi.org/10.1029/JC086iC10p09776>, 1981.
- 1133 Wani, R. P., Roper, D. L., Agrawal, R., Lim, E. J., Rollins, N. E., and Berelson, W. M.: Dataset
 1134 for “The Impact of NaOH , CaO , and $\text{Ca}^{2+} + \text{HCO}_3^-$ Addition on PIC and POC Formation in
 1135 Los Angeles Harbor Waters,” <https://doi.org/https://doi.org/10.5281/zenodo.20059856>, 19
 1136 February 2026.
- 1137 Wolf, S. E., Leiterer, J., Kappl, M., Emmerling, F., and Tremel, W.: Early Homogenous
 1138 Amorphous Precursor Stages of Calcium Carbonate and Subsequent Crystal Growth in Levitated
 1139 Droplets, *J. Am. Chem. Soc.*, 130, 12342–12347, <https://doi.org/10.1021/ja800984y>, 2008.
- 1140 Wolf-Gladrow, D. and Riebesell, U.: Diffusion and reactions in the vicinity of plankton: A
 1141 refined model for inorganic carbon transport, *Mar. Chem.*, 59, 17–34,
 1142 [https://doi.org/10.1016/S0304-4203\(97\)00069-8](https://doi.org/10.1016/S0304-4203(97)00069-8), 1997.
- 1143 Zachos, J. C., Röhl, U., Schellenberg, S. A., Sluijs, A., Hodell, D. A., Kelly, D. C., Thomas, E.,
 1144 Nicolo, M., Raffi, I., Lourens, L. J., McCarren, H., and Kroon, D.: Rapid Acidification of the
 1145 Ocean During the Paleocene-Eocene Thermal Maximum, *Science* (1979)., 308, 1611–1615,
 1146 <https://doi.org/10.1126/science.1109004>, 2005.
- 1147 Zondervan, I., Zeebe, R. E., Rost, B., and Riebesell, U.: Decreasing marine biogenic
 1148 calcification: A negative feedback on rising atmospheric $p\text{CO}_2$, *Global Biogeochem. Cycles*, 15,
 1149 507–516, <https://doi.org/10.1029/2000GB001321>, 2001.
- 1150
- 1151