

Instantaneous radiative forcings due to the first indirect effect linked to ~~warm~~-low-level liquid clouds in the Amazon

Andre Cezar Pugliesi¹, Alexandre Lima Correia², and Micael Amore Cecchini¹

¹Institute of Astronomy, Geophysics and Atmospheric Sciences, USP, São Paulo, Brazil

²Institute of Physics, USP, São Paulo, Brazil

Correspondence: Andre Cezar Pugliesi (andre.cezar.silva@gmail.com)

Abstract. Much of the present uncertainty in effective radiative forcing due to aerosol–cloud interactions (ERFaci) arises from aerosol–cloud adjustments. Nevertheless, estimating instantaneous radiative forcing due to aerosol–cloud interactions (IRFaci) remains important because it provides observation-based benchmarks for model evaluation. Such estimates are particularly scarce for ~~warm~~-low-level liquid Amazonian clouds, particularly from approaches combining surface-based remote sensing with in situ aircraft observations. Here, we estimate IRFaci for low-level ~~warm~~-liquid clouds over Amazon using the GoAmazon2014/5 datasets. Cloud microphysical properties were constrained with ground-based remote sensing and in situ measurements, used to configure cloud representations and coupled to libRadtran simulations of daily top-of-atmosphere upward irradiance. To reduce uncertainty in baseline atmosphere, two clean reference states are defined, including one designed to represent the seasonal variability of natural background aerosol conditions. Campaign-mean IRFaci values were -11.8 W m^{-2} (interquartile range: -23.0 to -2.4 W m^{-2}) and -1.3 W m^{-2} (-5.8 to 0.3 W m^{-2}) for the two reference-state definitions. The first estimate matches the maximum literature IRFaci per AOD unit in Amazon; the second aligns is in order of magnitude with IPCC’s global mean IRFaci, $-0.7 \pm 0.5 \text{ W m}^{-2}$. Sensitivity tests showed-a-strong-for 2014 showed a dependence of IRFaci on aerosol load under clean conditions, decreasing with higher loads. Although it does not quantify aerosol–cloud adjustments or ERFaci, this research provides an observationally constrained estimate of IRFaci in the Amazon, serving as a benchmark for future Amazon-focused studies of ERFaci.

1 Introduction

Anthropogenic factors generate disturbances in the global climate system and cause climatic radiative forcings (RF), defined as differences in global net irradiance between a reference atmospheric state and a state perturbed by an external climate agent (IPCC, 2013). The radiative imbalance caused by RF can prompt the atmosphere to seek a new equilibrium through variations in its internal temperature (IPCC, 2013). This re-equilibration can be modulated by the temperature of the oceans and/or continents (Levitus et al., 2012; Trenberth et al., 2014), whether or not adjustments are made to the tropospheric and/or stratospheric temperature profiles (Solomon et al., 2010; Santer et al., 2013). Effective radiative forcings (ERF) are calculated after these profiles have adjusted and reached a new radiative balance. In contrast, instantaneous radiative forcings (IRF) are obtained when no changes in temperature profiles are considered (IPCC, 2013).

25 Atmospheric aerosols act as cloud condensation nuclei (CCN) and ice nucleation particles (INP), playing a crucial role in cloud formation (Petters and Kreidenweis, 2007). Aerosol–cloud interactions (ACI) influence cloud microphysics (Wang et al., 2024), altering the droplet size distribution (Cecchini et al., 2016), optical properties (Zhao and Garrett, 2015), lifetime (Albrecht, 1989) and precipitation formation (Machado et al., 2018). Consequently, ACIs involving anthropogenic aerosols significantly affect global radiative balance and represent one of the main sources of uncertainty in climate models (Forster et al., 2021). The radiative forcing resulting from ACI (RFaci) can also be categorized as effective (ERFaci) or instantaneous (IRFaci). ERFaci accounts for adjustments in the initial changes in cloud droplet and ice crystal formation rates. These adjustments stem from variations in cloud fraction (f_c), cloud liquid water content (LWC), convective intensity, precipitation efficiency, and cloud lifetime, in addition to cloud formation or suppression in remote areas due to altered atmospheric circulation (Forster et al., 2021). IRFaci corresponds to the increase in cloud optical depth and albedo for a constant LWC. Environments with 30 high aerosol concentrations increase the number of small droplets of clouds, increasing the albedo of the cloud that reflects more radiation to space — a mechanism known as the "cloud albedo effect," the "Twomey effect," or the "first indirect aerosol effect" (Twomey, 1974, 1977).

The estimation of ERFaci and IRFaci on global and regional scales has been the focus of many studies (e.g. Dong and Mace, 2003; Lebsock et al., 2008; Hasekamp et al., 2019; Diamond et al., 2020). The complexity in quantifying RFaci lies mainly 40 in representing the reference atmosphere (Haywood and Boucher, 2000; Hamilton, Lee, Pringle, Reddington, Spracklen, and Carslaw, 2014), especially when the pre-industrial (PI) period is considered (Lee et al., 2016). This representation depends on a set of hypotheses and assumptions, including a degree of arbitrariness (Gryspeerdt et al., 2023) and constitutes an important source of uncertainty (Carslaw et al., 2013). Therefore, a detailed evaluation of the aerosol conditions in the reference state can reduce the uncertainties associated with the RFaci estimates (Gryspeerdt et al., 2023).

45 Warm clouds are regarded as the main mediators of aerosol radiative forcing through ACI (Heyn et al., 2017; Mülmenstädt and Feingold, 2018). Several studies have used modeling or satellite remote sensing data to quantify and analyze RFaci values associated with warm clouds (e.g. Quaas et al., 2009; Chen et al., 2014; Christensen et al., 2017; Toll et al., 2019), often disregarding continental areas. Among the more regionalized efforts (e.g. Keil and Haywood, 2003; Mace et al., 2006; Marsing et al., 2023; Wang et al., 2023; Herbert and Stier, 2023), only Herbert and Stier (2023) aimed to quantify IRFaci in the Amazon 50 region, also using satellite sensor data. This shows that research on instantaneous forcings due to the Twomey effect in warm clouds over the Amazon rainforest remains limited, particularly when the estimates are derived from in situ or ground-based observations.

This study proposes a new method to estimate IRFaci caused by aerosol pollution in low-level ~~warm-liquid~~ clouds in the central Amazon. The IRFaci calculations are based on ground-based and in situ measurements and are performed according to 55 two distinct "clean" reference atmospheric states. We hypothesize that a reference state based on irradiance values that follow the natural seasonal variability of aerosols could reduce the uncertainties in IRFaci estimates. We also assess the sensitivity of ~~warm-cloud~~low-level liquid-cloud IRFaci to different aerosol concentrations in the Amazonian atmosphere.

2 Methodology

2.1 Spatio-temporal domain

60 The annual cycle of aerosol number concentration in the Amazon is characterized by extreme variations, from 250 particles cm^{-3} during the wet season to values as high as 30,000 particles cm^{-3} in the dry season (Pöschl et al., 2010). This contrast makes the Amazon region a natural laboratory for studying aerosol–cloud interactions. In this study, datasets obtained during the GoAmazon2014/5 campaign (Martin et al., 2016), carried out in 2014 and 2015 in the Brazilian Amazon, were used. The experimental sites of GoAmazon2014/5 were selected based on seasonal timing and the prevailing wind direction. The
65 sites designated as “T0” were located north of the city of Manaus, a region undisturbed by the urban pollution plume due to the prevailing northeasterly winds. The Sites “T1” and “T2” were positioned south of Manaus. The “T3” and “T3u” sites were located southwest of the city, alternately influenced by clean and polluted air masses (Martin et al., 2016) due to variations around the mean wind direction.

In this work, we used data collected from the T3 site (3.2° S, 60.6° W), which accommodated the largest number of instru-
70 ments during both GoAmazon2014/5 campaign years. At T3, instruments operated by the U.S. Department of Energy (DOE) under the Atmospheric Radiation Measurement Program (ARM, <https://www.arm.gov/>) were deployed, including the ARM Mobile Facility 1 (AMF-1) and the ARM Mobile Aerosol Observing System (MAOS; (Uin et al., 2019)). Furthermore, during the two intensive operational periods (IOP1 and IOP2, from 1 February to 31, March 2014, and from 15 August to 15, October 2014, respectively) of GoAmazon2014/5, other instrumental platforms were also employed, such as the Gulfstream-159
75 aircraft (“G1”; (~~Schmid et al., 2014~~)[Schmid et al. \(2014\)](#)) and the HALO aircraft (High Altitude and Long Range Research Aircraft; (~~Wendisch et al., 2016~~)[Wendisch et al. \(2016\)](#)). The HALO aircraft flew as part of a collaboration effort with the Brazilian-German project ACRIDICON-CHUVA (Wendisch et al., 2016).

2.2 Ground-based datasets

The atmospheric and cloud properties obtained by ground-based remote sensing included the f_c , the liquid water path (LWP),
80 the cloud base height, and the cloud top height (CBH and CTH, respectively). All datasets were maintained and provided continuously by the ARM program. The data set f_c originated from a Value-Added Product (~~VAP; (Riihimäki et al., 2019)~~
[RADiative FLUX ANALysis VAP, Riihimäki et al. \(2019\)](#)) that combined measurements from different radiometers to derive total, direct and diffuse shortwave and longwave downward and upward irradiances, as well as f_c for shortwave (Long et al., 2006) and longwave (Dürr and Philipona, 2004) radiation. The f_c values were retrieved at one-minute intervals and did not
85 distinguish between different types of cloud.

The LWP values ~~were also (sourced from the MicroWave Radiometer RETrievals VAP and with 1-sigma uncertainty,~~
[Gaustad et al. \(2011\)\)](#) ~~were~~ derived through an algorithm (Turner et al., 2007) that combines atmospheric thermodynamic state information — vertical profiles of air temperature, pressure, and relative humidity, [used in independent steps of the](#)
[Turner et al. \(2007\) method](#) — with brightness temperature measurements at 23.8 GHz and 31.4 GHz obtained by a surface-
90 based microwave radiometer (MWR). ~~As in f_c , the~~ [The](#) base and top heights of the observed cloud layers were derived from ~~a~~

~~VAP (Kollias et al., 2007) that~~ the Active Remote Sensing of Clouds VAP (ARSCL VAP, Kollias et al. (2007)) which combined measurements from a W-band ARM Cloud Radar (WACR, operating at 95.04 GHz) and a micropulse lidar system (MPL), with CBH and CTH values sampled every 5 seconds. An important observation is that the LWP data measured using the MWR could, in principle, be validated from in-situ LWC measurements made in clouds during flights over T3 (more on these LWC data in Sect. 2.3), when co-located spatially and temporally with cloud radar and lidar (WACR + MPL) measurements, used to estimate CBH and CTH. However, there were no cases of simultaneous measurements from the three instruments at both IOPs and, for this reason, we were unable to cross-validate the LWP data.

The total aerosol scattering coefficient ($\beta_{sct.}$), the total concentration of organic mass (“Total Organics” or “Org.”), the number concentration of cloud condensation nuclei (CCN) and the spectral aerosol optical depth (AOD) were among the aerosol properties analyzed. The values of $\beta_{sct.}$ were derived (Uin, 2024) using a pair of nephelometers measuring $\beta_{sct.}$ as a function of relative humidity at intervals of one-min, at three wavelengths λ (700, 550 and 450 nm), and for two cuts in particle size (1 μm and 10 μm). The values of Total Organics were obtained using an Aerosol Chemical Speciation Monitor (ACSM), a mass spectrometer based on thermal vaporization and subsequent ionization of aerosol particles smaller than 1 μm (Watson, 2017).

The CCN number concentrations at the T3 site were measured using a cloud condensation nuclei particle counter (CCNC). This instrument measures aerosol concentrations by sampling air through a column with thermodynamically unstable supersaturated water vapor, which can condense on aerosol particles. The resulting grown particles are counted and sized by an Optical Particle Counter (OPC). Thus, CCNC quantifies the number concentration of particles activated as a function of the instrument’s internal supersaturation (Uin and Enekwizu, 2024).

Spectral AOD values constituted the final aerosol-related variable used in this study. Due to the limited availability of samples, two complementary datasets were employed. The first was obtained from a sun photometer belonging to AERONET (AErosol RObotic NETwork; <https://aeronet.gsfc.nasa.gov/>), an instrument equipped with a 25 cm collimator system mounted on a robotic base that can be programmed to point either toward the sky or directly at the Sun. This photometer measures spectral solar and sky radiances at eight wavelengths (340, 380, 440, 500, 675, 870, 1020, and 1640 nm) and uses them to calculate AOD values and other quantities (Giles et al., 2019), after correcting for molecular scattering and absorption caused by ozone and other atmospheric gases. The AERONET-derived AOD values used here correspond to the 500 nm wavelength, being measured at variable sampling rates (ranging from 1.9 to 551.3 minutes).

~~This~~ The AERONET’s data set was complemented with AOD values retrieved from solar irradiance measurements acquired by a MultiFilter Rotating Shadowband Radiometer (MFRSR) operated by ARM. These irradiances serve as input for a VAP that applies the algorithms described by Harrison and Michalsky (1994), Michalsky et al. (2001), and Alexandrov et al. (2004) to derive spectral AOD values at six wavelengths (415, 500, 615, 673, 870, and 940 nm), with a sampling interval of 20 seconds.

2.3 In situ datasets

In situ measurements of the liquid water content in clouds, total water content (TWC), total cloud droplet concentration (N_d), cloud droplet size distribution ($N(D)$), and effective cloud droplet diameter (D_{eff}) were performed using instruments onboard the G-1 and HALO aircraft.

Of the 35 flights conducted by the G-1 during IOP 1 and IOP 2, 32-29 flew over the retangular area (60.75° W, 3.3° S, 60.45° W, 3.1° S) of 0.2° x 0.3° centered at the T3 site and were used in this study. ~~Measurements of~~ LWC and TWC measurements were carried out using the WCM-2000 instrument, a sensor composed of a scoop-shaped device equipped with several heated wires. Some of these wires are directly exposed to the airflow and cloud water droplets, while a separate reference wire is exposed only to the airflow. A power control system maintains constant voltage in one wire and constant temperature in the others. In the case of constant temperature operation, the “King-type” probe (King et al., 1978) uses the system’s power values, relative airspeed, ambient temperature, individual wire dimensions, and the latent heat of water vaporization to calculate LWC and TWC each second (for calculation details, see the user manual available at <https://www.scieng.com/pdf/WCM2000User.pdf>).

The N_d and $N(D)$ values measured during the G-1 flights were obtained using the Fast Cloud Droplet Probe (FCDP; manual available at <http://www.specinc.com/node/123>), an instrument that employs a focused laser beam to size hydrometeors based on the amount of scattered light during particle–beam interaction. The FCDP has a resolution of 3 μm , measures droplet diameters between 1 and 50 μm , and divides them into 20 equally spaced size bins.

The HALO aircraft conducted 14 research flights between September and October 2014, during the GoAmazon2014/5 IOP 2. The flights were designed to cover five distinct mission types, focusing on different processes involving clouds, aerosols, chemistry, and radiation (Wendisch et al., 2016). The flights were numbered chronologically as ACXX, with X ranging from 07 to 20. Among these, only two flew over clean densely forested areas to the north and northwest of Manaus (AC09 and AC18, conducted on 11 September and 28 September, 2014, respectively) ~~and are used here~~. Those flights were ~~selected to serve as the background reference because~~ conducted over two regions practically unaffected by biomass burning plumes from the deforestation arc in the Amazon, characterizing the cleanest possible atmosphere among all flights performed, as indicated in Cecchini et al. (2016). Despite the low number of samples, these flights are representative of the cleanest possible atmospheric regime within the polluted season in the region, which is why they constitute the best available samples to characterize the second clean scenario used in this study. Given that the clouds measured during it were not directly affected by biomass-burning aerosols, those flights were selected to serve as a background reference.

LWC and D_{eff} HALO measurements were obtained using the Cloud and Aerosol Spectrometer with Detector for Polarization (CAS-DPOL), a spectrometer that also measures the cloud droplet size distribution from 0.5 to 50 μm (Wendisch et al., 2016). Filtering of LWC and D_{eff} for warm-low-level liquid clouds was performed using flight altitude data (above sea level, ASL) from the Basic HALO Measurement and Sensor System (BAHAMAS), considering a maximum flight height of 3 km.

Table 1 compiles the atmospheric, cloud, and aerosol variables used in this study. The table also shows the corresponding instrument, measurement unit, experimental platform, and sampling interval for each variable. For CBH, CTH, and LWP, 1-minute averages were calculated to facilitate the derivation of microphysical parameters for subsequent warm-low-level

liquid-cloud modeling. For AOD, $\beta_{sct.}$, Total Organics, CCN, and f_c , daily and/or diurnal averages were calculated. The mathematical formulations of the variables presented in Table 1 are provided in Sect. 2.4.

Table 1. Cloud and aerosol properties used in this work.

Variable	Unit	Instrument	Platform	Sampling interval	Observation
LWC	g m^{-3}	WCM-2000	G-1	1 second	IOP 1 and 2
LWC	g m^{-3}	CAS-DPOL	HALO	1 second	IOP 2
TWC	g m^{-3}	WCM-2000	G-1	1 second	IOP 1 and 2
N_d	drops L^{-1}	FCDP	G-1	1 second	IOP 1 and 2
$N(D)$	$\text{drops L}^{-1} \mu\text{m}^{-1}$	FCDP	G-1	1 second	IOP 1 and 2
D_{eff}	μm	CAS-DPOL	HALO	1 second	IOP 2
Altitude	m ASL	BAHAMAS	HALO	1 second	IOP 2
CBH	m AGL	WACR + MPL	Ground	5 seconds	2014–2015
CTH	m AGL	WACR + MPL	Ground	5 seconds	2014–2015
LWP	g m^{-2}	MWR	Ground	20 seconds	2014–2015
f_c	%	Radiometers	Ground	1 minute	Shortwave; 2014–2015
AOD	unitless	AERONET	Ground	Variable	550 nm; 2014–2015
AOD	unitless	MFRSR	Ground	20 seconds	500 nm; 2014–2015
$\beta_{sct.}$	Mm^{-1}	Nephelometer (NEPH3W)	Ground	1 minute	500 nm; 1 μm cut; 2014–2015
Total organics (Org.)	$\mu\text{g m}^{-3}$	ACSM	Ground	1 minute	~50–700 nm; 2014–2015
CCN	particles cm^{-3}	CCNC	Ground	1 minute	2014–2015

2.4 Clean and polluted scenarios

Periods with lower and higher average aerosol number concentration during the GoAmazon2014/5 campaign will be referred to as “clean” and “polluted” scenarios, respectively. The term “clean” does not imply a complete absence of anthropogenic aerosols in the atmosphere, but rather a distinctly lower concentration compared to the “polluted” scenario. The latter corresponds to episodes in 2014 and 2015 with a higher frequency of biomass-burning events and advection of the Manaus pollution plume over the T3 site.

The identification of clean and polluted scenarios during GoAmazon2014/5 was based on a manual analysis of daily means of $\beta_{sct.}$, Org., CCN, and AOD. Among these quantities, $\beta_{sct.}$ and AOD are directly proportional to aerosol loading and result from the interaction between aerosols and incident electromagnetic radiation, particularly at wavelengths between approximately 0.3 μm and 4 μm — the solar spectrum range. This interaction is described by Mie scattering (Mie, 1908), which applies Maxwell’s electromagnetic theory to describe radiation scattered by spherical particles with radius r comparable to or greater than the incident wavelength. Macroscopically, $\beta_{sct.}$ at position s' along an optical path s for aerosols with refractive index $n(\lambda)$ is given by Eq. (1), where $N_p(r, s')$ is the number of aerosol particles per unit air volume per size interval at position s' , and $Q_{sct.}$

is the scattering efficiency factor. Spectral AOD is related to the amount of optically active aerosols in the atmospheric column. To derive it, one must consider the solar zenith angle (θ_0 or SZA, formed between the vertical direction and the propagation path of the incident radiation beam) and the linear absorption coefficient $\beta_{abs.}$ (Eq. (2)), resulting in the formulation shown in Eq. (3).

$$\beta_{sct.}(\lambda, s') = \pi r^2 Q_{sct.} \left[\frac{2\pi r}{\lambda}, n(\lambda) \right] \int_0^\infty N_p(r, s') dr \quad (1)$$

$$\beta_{abs.}(\lambda, s') = \pi r^2 Q_{abs.} \left[\frac{2\pi r}{\lambda}, n(\lambda) \right] \int_0^\infty N_p(r, s') dr \quad (2)$$

$$AOD(\lambda) = \int_0^s [\beta_{sct.}(\lambda, s) + \beta_{abs.}(\lambda, s)] \cos(\theta_0) ds \quad (3)$$

Daily means of $\beta_{sct.}$, Org., and CCN for 2014 are displayed in Fig. 1. The period between days 10 and 90 of 2014 was selected as representative of the clean scenario, whereas the intervals between days 210–277 and 300–345 were chosen to represent polluted periods. For 2015, the same procedure was applied to the daily time series of $\beta_{sct.}$, Org., and AOD. Figure 2, which presents these series, shows that aerosol influence peaked during the second half of the year. During the polluted period of 2015, the $\beta_{sct.}$ and Org. peaks were greater than those observed in 2014, which exhibited a shorter clean period compared to 2015. From the time series shown in Figs. 1 and 2, the start and end days of the clean and polluted scenarios for 2014 and 2015 were determined. Table 2 summarizes these intervals and the weighted means of $\beta_{sct.}$, Org., CCN, and AOD calculated for them. This reference will be extensively used in the calculation and analysis of IRFaci.

Table 2. Days that define the clean and polluted scenarios for 2014 and 2015. The values in parentheses are the weighted means of $\beta_{sct.}$, Org. and CCN (for 2014, respectively) and of $\beta_{sct.}$, Org. and AOD from AERONET (for 2015, respectively) calculated for the corresponding interval.

Year	Clean	Polluted 1	Polluted 2
2014	Jan. 11th to Apr. 1st (7.7, 2.3, 400.2)	July 30th to Oct. 5th (30.8, 9.1, 1748.5)	Oct. 28th to Dec. 12th (22.4, 6.8, 1411.8)
2015	Jan. 1st to July 30th (6.7, 2.9, 0.1)	Sep. 23th to Nov. 30th (104.9, 28.8, 0.6)	—

The clean scenarios of 2014 and 2015 define the periods within which the cleanest days were used as the basis for the determination of the first atmospheric reference state. The selection of these cleanest days proceeded as follows: first, all days sharing common data for the variables $\beta_{sct.}$, Org., CCN, and AOD (for AOD, since the AERONET dataset had many empty data points, only data derived from the MFRSR were considered) were identified. For these days, daytime medians (06:00–18:00 LT) were calculated for each of the four variables. After normalizing these medians, the K-Means clustering algorithm (Lloyd, 1982) was applied using 2, 3, and 4 clusters. For each cluster and each variable, a characteristic median was assigned. Comparing the




Figure 1. Daily averages of $\beta_{sct.}$, Org. and CCN for 2014. The legend indicates the instruments responsible for data acquisition and the days that demilit the clean and polluted scenarios. The identification of cleanest days within the clean scenario was carried out using a clustering technique described in Sect. 2.4.




Figure 2. Daily averages of $\beta_{sct.}$, AOD and Org. for 2015. CCN values were not considered because, for most of 2015, measurements made by CCNPC were classified as incorrect by ARM quality flags. The sparse nature of the MFRSR AOD data outside the day 150-270 window was complemented by the AERONET AOD data.

medians across different cluster counts helped determine the optimal number of clusters, corresponding to cases in which no overlap occurred between the characteristic medians of distinct groups.

From the identification of optimal cluster counts for 2014 and 2015, the minimum characteristic median values were obtained, allowing the identification of the cleanest days within each clean scenario. For 2014, six days in February (11, 17, 23, 24, 26, and 28) and ten in March (2, 6, 7, 8, 12, 20, 21, 22, 25, and 28) were selected. For 2015, the cleanest days were 25 February and 3–4 May. The mean upward irradiance values at the top of the atmosphere (TOA) simulated for these days were later used as a fixed reference for the IRFaci calculations.

2.5 Extension of $D_{eff.}$ and LWC measurements for the GoAmazon2014/5 period

Upward irradiance values at the TOA were obtained using the libRadtran radiative transfer code (Emde et al., 2016), which simulated ~~warm-low-level liquid~~ clouds modeled from LWC values (Eq. (4); (~~Morrison et al., 2005~~)Morrison et al. (2005)), where ρ is the liquid water density, r is the cloud droplet radius, $N(D)$ is given by Eq. (5), and $r_{eff.}$ by Eq. (6a) (Stephens, 1978). However, experimental in situ values of LWC, TWC, $N(D)$, N_d , and $D_{eff.} = 2 r_{eff.}$ were obtained only during G-1 flights, limiting the temporal coverage of these datasets. The procedure used to ~~extend their availability~~ derive new $r_{eff.}$ and LWC values based on those data consisted of:

1. Deriving empirical relationships between the in situ measured LWC and $r_{eff.}$ values;
2. Using surface-based datasets of CBH, CTH, and LWP to calculate $LWC_{ground} = LWP/(CTH-CBH)$;
3. Inserting LWC_{ground} values into the empirical relationships obtained in the first step to extrapolate $r_{eff.}$ for the entire GoAmazon2014/5 campaign period.

The first step in obtaining empirical relationships between in situ LWC and $r_{eff.}$ was to consider only positive simultaneous measurements of N_d and LWC, ensuring that the aircraft was within a cloud. Subsequently, only cases with zero ice water content (IWC, equals the mathematical difference between TWC and LWC) were retained, thereby filtering ~~warm-liquid~~ clouds. During data screening, cases where $N_d < 0.3$ particles cm^{-3} and $LWC < 0.02$ g m^{-3} were excluded, removing experimental points associated with fog or extremely tenuous clouds. These filters were applied for both IOP's, with the aim of ensuring consistency of data selection for two intensive periods. Finally, $r_{eff.}$ was computed according to Eq. (6b), where $\langle r_j \rangle$ represents the j -th mean of one of the 20 size intervals (bins) in which $N(D)$ values were classified.

$$LWC = \frac{4\pi}{3} \rho \int_0^{\infty} r^3 N(D) dr \quad (4)$$

$$N(D) = N_0 r^m e^{-\Lambda r} \quad (5)$$

$$220 \quad r_{\text{eff.}} = \frac{\int_0^\infty r^3 N(D) dr}{\int_0^\infty r^2 N(D) dr} \quad (6a)$$

$$r_{\text{eff.}} = \frac{\sum_{j=1}^{20} \langle r_j \rangle^3 N(D)_j dr}{\sum_{j=1}^{20} \langle r_j \rangle^2 N(D)_j dr} \quad (6b)$$

The scatterplot of $r_{\text{eff.}}$ versus LWC for IOP 1 did not exhibit patterns that could be easily fitted by simple functions (e.g., linear or quadratic). Consequently, the data were divided into 20 equally spaced LWC bins (with a width of 0.05 g m^{-3} , between 0.02 and 1.02 g m^{-3}), within which LWC and $r_{\text{eff.}}$ medians were calculated. Similarly to this study, Reid et al. (1999) explored
 225 the relationship between LWC and $r_{\text{eff.}}$ of ~~warm-liquid~~ clouds over the Brazilian Amazon. Those authors used in situ aircraft measurements focusing on ~~warm-liquid~~ stratiform and cumuliform clouds with bases located within the planetary boundary layer. Their datasets, obtained between August and September 1995 — when the Amazonian atmosphere is typically affected by smoke and biomass-burning aerosols — produced $r_{\text{eff.}}$ -LWC distributions for cumulus and stratocumulus clouds that closely resembled those observed here for IOP 1.

230 For IOP 2, the LWC/N_d ratios were considered instead of LWC alone. These ratios were then applied to Eqs. (7) and (8), corresponding to cumulus and stratocumulus clouds, respectively (Reid et al., 1999). In their analysis, Reid et al. (1999) concluded that, after testing several fitting functions, the hyperbolic arccosine function provided the best fit — motivating its adoption here to adjust the median pairs for IOPs 1 and 2 (Fig. 3).

$$r_{\text{eff.}} = 6.67 \left(\frac{\text{LWC}}{N_d} \right)^{0.31} \quad (7)$$

$$235 \quad r_{\text{eff.}} = 7.50 \left(\frac{\text{LWC}}{N_d} \right)^{0.33} \quad (8)$$

Following the procedure to ~~extend the dataset coverage, derive new $r_{\text{eff.}}$ and LWC values, the second step was to use~~ surface-derived values of CBH, CTH, and LWP ~~were used~~ to compute $\text{LWC}_{\text{ground}}$. These $\text{LWC}_{\text{ground}}$ values were then inserted into the empirical relationships shown in Fig. 3, generating new $r_{\text{eff.}}$ estimates for ~~warm-liquid~~ clouds during IOPs 1 and 2. This procedure enabled the construction of an expanded database, covering all days within the clean and polluted periods defined
 240 by the intervals listed in Table 2, thereby increasing data availability across 2014 and 2015.

For transition periods, hybrid adjustments were made using weighted averages that combined both empirical functions. For instance, during the first ~~transitional-transition~~ days of 2014, the ~~computed-calculated~~ $r_{\text{eff.}}$ values had a greater weight from the clean-period equation, while the last days were progressively influenced by the polluted-period function. This approach enabled the generation of $r_{\text{eff.}}$ values as a function of $\text{LWC}_{\text{ground}}$ for nearly every day of the GoAmazon2014/5 campaign,
 245 consequently allowing the modeling of ~~warm-liquid~~ clouds for almost the entire observation period.




Figure 3. Dispersion of medians of r_{eff} and LWC calculated in the 20 bins for IOP 1 and 2. The medians included experimental points from all altitudes where clouds with liquid water were identified and for which $N_d \geq 0.3 \text{ particles cm}^{-3}$ and $\text{LWC} \geq 0.02 \text{ g m}^{-3}$. Although no filter was applied to distinguish between cumuliform and stratiform clouds, these clouds were selected using these same filters in both IOP. The adjustments (hyperbolic arc cosine function) originated the equations shown in the graph's legend.

2.6 Configuration of Warm-Low-level liquid Cloud Models

Selection of warm-low-level liquid clouds was also performed through filtering of the LWC_{ground} and CTH datasets. The interval $[0.2-0.4] \text{ g m}^{-3}$ for LWC_{ground} encompasses typical values of cumulus and stratiform clouds in both clean and polluted continental environments (Hess et al., 1998), and was therefore applied as a filter to the LWC_{ground} data. Additionally, only clouds with CTH less than or equal to 3 km were considered to ensure that the final results represented warm-low-level clouds exclusively. Added to the filters previously applied to the in situ r_{eff} and LWC values, the filters applied to the LWC_{ground} and the corresponding calculated r_{eff} data justify the term "low-level liquid", which includes both shallow cumulus and low stratiform cloud types in this work.

The effective radius of liquid clouds increases with altitude in the atmosphere (Martins et al., 2011; Rosenfeld and Woodley, 2003), and this behavior was adopted as the justifying the selection of this behavior as conceptual model to simulate characteristic warm-low-level liquid clouds for each day of the GoAmazon2014/5. To represent the vertical increase of r_{eff} with

altitude, the clouds were configured with two horizontal layers, ensuring that the r_{eff} and LWC values in the models were higher in the upper layer. The base height of the modeled clouds was set according to the 12.5th percentile of CBH values, while the top height corresponded to the 87.5th percentile of the CTH data distribution. The central height was defined as the arithmetic mean between the base and top heights. The 12.5th and 87.5th percentiles of the daily r_{eff} and LWC_{ground} distributions were similarly selected to describe the r_{eff} and LWC values at the base and top, respectively, while the middle layer was represented by the median of the same distributions. Although these statistics are insufficient to explicitly capture intraday variability or transient processes, constituting a low-dimensional description, they are adequate to characterize the central tendency and dispersion of the variables mentioned. Table 3 summarizes the statistical parameters used to simulate representative ~~warm~~-low-level liquid clouds for most days of the GoAmazon2014/5 campaign.

Table 3. Configuration of “daily” modeled clouds in LibRadtran, which does not require the complete droplet size distribution as input. The statistics shown were calculated from daily distributions of LWC_{ground} (equals the ratio between LWP and CTH-CBH difference) and of the corresponding r_{eff} , which was calculated from adjusted equations shown in Fig. 3.

Cloud region	Height (km)	LWC (g m^{-3})	r_{eff} (μm)
Top	CTH 87.5th percentile	87.5 percentile	87.5 percentile
Center	Mean between base and top	50.0 percentile	50.0 percentile
Base	CBH 12.5th percentile	12.5 percentile	12.5 percentile

2.7 Radiative transfer code configuration to obtain upward irradiance at TOA

The integrated upward irradiance values (hereafter referred to as F) at the TOA were obtained using the libRadtran radiative transfer model, which applies the DISORT method (DIScrete Ordinate Radiative Transfer solver; Stamnes et al. (1988)) to solve the radiative transfer equation after accounting for adjustments to atmospheric boundary conditions. In addition, libRadtran uses the “medium” spectral resolution mode (bandwidth = 5 cm^{-1}) of the REPTRAN parameterization (Gasteiger et al., 2014) as the standard spectral configuration. The wavelength interval selected for the simulations ranged from 300 to 1000 nm, encompassing the entire visible spectrum and a small portion of the near-infrared region. The solar spectrum used extended from 250 to 10,000 nm, with a resolution of 0.1 nm.

REPTRAN is also the default method employed by libRadtran to represent atmospheric molecular scattering, applicable to both the solar and terrestrial spectra. For this, vertical profiles of the main atmospheric gases (O_2 , O_3 , NO_2 , CO_2 , and water vapor) must be provided. The “standard tropical atmosphere” file available in libRadtran already includes vertical profiles for these gases, along with profiles of air density, temperature, and pressure. The profiles of less abundant trace gases such as N_2O , CO , and CH_4 were taken from the “standard U.S. atmosphere” dataset. Considering this, the tropical atmosphere data provided by the Earth Observation Data Group (EODG) of the University of Oxford (<http://eodg.atm.ox.ac.uk/RFM/atm/minor.atm>) were used here to generate new vertical profiles for these gases. All the profiles mentioned were used equally in all simulations.

seeking to follow the definition of "instantaneous" forcing, namely, forcings obtained with the tropospheric and stratospheric profiles of constant T.

The files used to simulate ~~the representative daily warm~~ representative daily low-level liquid clouds considered the exact geographic location and elevation of the T3 site. Since T3 was situated in a forested region, a surface albedo of 14% was adopted. The mean CO₂ mixing ratios estimated for 2014 and 2015 according to the Keeling curve (Keeling et al., 2017) were also ~~incorporated~~ taken into account. All simulations were configured to represent upward irradiance at TOA for 16:00 UTC (12:00 LT), ensuring that the Sun was near zenith at the simulation time. Moreover, the f_c was set to 100% for all simulations, assuming that the simulated ~~stratiform warm~~ clouds completely covered the sky at 12:00 LT. Typical examples of input files (extension .inp) used in LibRadtran can be found at the link <https://doi.org/10.5281/zenodo.20610494>.

2.8 IRFaci estimates for the two clean atmospheric reference states

The upward integrated irradiance values at the TOA obtained from libRadtran (hereafter $F_{libRadtran}$) do not account for variations in the Earth–Sun distance (d). The necessary correction was made using Eq. (9), which assumes a plane-parallel atmosphere. The values of d and the solar zenith angle (θ_0) used in Eq. (9) were calculated using spreadsheets provided by NOAA's Earth System Research Laboratories (<https://gml.noaa.gov/grad/solcalc/calcdetails.html>). These spreadsheets compute various astronomical parameters for any location on Earth; therefore, the d and θ_0 data were obtained specifically for the T3 site. With these parameters and the $F_{libRadtran}$ outputs, the corrected irradiance values ($F_{corr.}$) were computed, accounting for temporal variations in d throughout 2014 and 2015 at T3.

$$F_{corr.} = \frac{d^2 F_{libRadtran}}{\cos(\theta_0)} \quad (9)$$

Instantaneous radiative forcings (IRF) were then calculated as the difference between $F_{corr.}$ in the reference state and $F_{corr.}$ in the anthropogenically perturbed state (Eq. (10)). Moreover, since the IRF calculations were based on integrated irradiances simulated for 12:00 LT at T3, the estimation of daily radiative forcing (IRF_{daily}, corresponding to 00:00 to 23:59) required consideration of the variation of the θ_0 throughout the day. This correction was implemented using Eq. 11, which incorporated 120 values of θ_0 sampled every 6 minutes during daylight hours. Because T3 is located near the Equator, the durations of daytime and nighttime were considered nearly identical, and the morning and afternoon periods were approximately equal in length. For this reason, IRF_{daily} corresponds to either the morning or afternoon interval, justifying the division by 2 in Eq. (11).

$$IRF = F_{corr}^{reference} - F_{corr}^{perturbed} \quad (10)$$

$$IRF_{daily} = \frac{1}{2} \frac{IRF \sum_{n=06:00}^{18:00} \cos(\theta_0)_n}{120} \quad (11)$$

As previously discussed, two clean atmospheric reference values were used in Eq. (10). The first, a fixed value of 689.9 W m⁻², corresponds to the mean of the corrected irradiances obtained from the cleanest days of the GoAmazon2014/5 campaign. This

reference represents natural background atmospheric conditions during the 19 cleanest days of 2014 and 2015, which may differ from those characterizing the polluted periods of the same years. To account for seasonal variability in cloud microphysical properties, a second reference with variable irradiance was implemented. This approach ~~involved generating warm-, which consists of a first approximation to account for the natural variation in the microphysical characteristics of clouds, involved~~
 315 ~~generating low-level liquid~~ cloud models (example shown in Table 4) based on LWC data (within the $[0.2\text{--}0.4]$ g m^{-3} interval) and $r_{eff.}$ values corresponding to the cleanest conditions identified during IOP 2. The measurements used ~~originated-came~~
 from HALO aircraft flights AC09 and AC18~~-, which represents the cleanest cases available among HALO flights made during the polluted period of 2014.~~

For these models, libRadtran simulated upward integrated irradiances at TOA for $\theta_0 = 0$ and an Earth–Sun distance of 1
 320 AU. The resulting irradiances, 766.7 W m^{-2} and 769.1 W m^{-2} , yielded a mean value of 767.9 W m^{-2} , which was considered representative of the cleanest possible atmospheric conditions for September 2014. This mean was used as the reference for IRF calculations only during the polluted periods of 2014 and 2015. For the transitional days, reference irradiances were calculated to vary linearly (Fig. 4) between the clean and polluted reference values (689.9 W m^{-2} and 767.9 W m^{-2} , respectively). ~~This~~
 325 ~~The assumption of this linear variation between the cleanest days of the clean period and the cleanest days within the polluted period of 2014 came after an analysis of the evolution of reff between these two scenarios, which occurs in an approximately linear manner throughout 2014, justifying the approximation used.~~

~~Using the seasonal reference based on aerosol availability aimed to reduce the seasonal contamination of thermodynamic variables (temperature, pressure, and water vapor profiles were the same for all simulations) and isolate the effect of aerosols on forcings. Since the effects of seasonal contamination due to dynamics on macrophysical characteristics of clouds were already~~
 330 ~~embedded in the statistics of these properties (such as LWP, CBH, and CTH), this~~ second approach allowed the computation of new daily forcing values less affected by the seasonal variability of natural atmospheric conditions ~~over-in~~ the T3 region.

Table 4. Cloud configuration representing 11 September 2014 (AC09), ~~considered~~ one of the two “cleanest”-days ~~with measurements available~~ during ~~the cleanest days of~~ IOP 2. The corresponding intervals of LWC ($r_{eff.}$) for 28 September 2014 were 0.00 to 0.24 g m^{-3} (0.00 to $5.48 \text{ }\mu\text{m}$), 0.24 to 0.27 g m^{-3} (5.48 to $8.03 \text{ }\mu\text{m}$), and 0.27 to 0.00 g m^{-3} (8.03 to $0.00 \text{ }\mu\text{m}$) for base, center, and top portions, respectively. The average LWC and $r_{eff.}$ values for these two days were 0.26 g m^{-3} and $6.89 \text{ }\mu\text{m}$, respectively. The zero extremes in cloud configuration are necessary for libRadtran to correctly simulate the boundaries of the modeled clouds.

Cloud region	Height (km)	LWC (g m^{-3})	$r_{eff.}$ (μm)
Top	2 to 3	0.28 to 0.00	8.30 to 0.00
Center	1 to 2	0.26 to 0.28	6.09 to 8.30
Base	0 to 1	0.00 to 0.26	0.00 to 6.09




Figure 4. Integrated and corrected upward irradiances ($F_{corr.}$) at TOA for 2014 and 2015 (bluish line varying over the day of the year). The values were calculated for each day and correspond to local solar noon at the T3 site. They are also unaffected by Earth’s orbital variation, so the variation over the years is due to changes in the cloud properties. The areas highlighted in blue (brown) represent the clear (polluted) periods for the two years. Days that do not show F values contain incomplete datasets (CBH, CTH, or LWP), making ~~warm~~-cloud modeling impossible. The solid (dashed) red (black) line corresponds to the F value used as a fixed (variable) reference to obtain the IRFaci estimates.

3 Results

3.1 First clean state - “Fixed” irradiance reference

The IRF_{daily} values were obtained assuming a cloud fraction of 100%. However, to regionalize the results according to the actual cloudiness observed at T3, each daily value was also multiplied by the corresponding daytime mean f_c for that specific day. Figure 5, which presents the IRF_{daily} values for the first clean-atmosphere reference, shows that the results incorporating the f_c measured at T3 generally display smaller absolute magnitudes compared to those associated with $f_c = 100\%$. The observed daily oscillation arises from the way the ~~warm~~low-level liquid-cloud models were configured — that is, based on the percentiles and medians of $r_{eff.}$ and LWC. Figure 5 also clearly reveals the mean reduction in IRF_{daily} magnitude as the more polluted periods approach, with values becoming increasingly negative during those episodes.




Figure 5. IRF_{daily} values are calculated based on the fixed clean atmosphere reference for 2014 and 2015. Blue line ($f_c = 100\%$, the same of " $f_c = 1$ ") indicates the results for an f_c of 100%, while the gold line shows the values according to the f_c measured at T3.

To evaluate the relative behavior of IRF_{daily} values across different aerosol-loading scenarios, boxplots were generated for each case. Figure 6 displays the results for 2014, along with the means and medians for each scenario. Overall, IRF_{daily} values that do not account for the measured f_c at T3 exhibit distributions shifted toward more negative values compared to those incorporating f_c variability, indicating that a constant $f_c = 100\%$ produces markedly stronger negative instantaneous radiative forcings. The statistics show progressively more negative values as the clean days give way to polluted ones, with results for the second 2014 transition period showing an increasing trend consistent with the decreasing aerosol parameters displayed in Fig. 1. It is also noteworthy that, although positive values occur only under the clean scenario (at the upper limit of the seventy-fifth percentile), both the means and medians for all scenarios remain negative.

The IRF_{daily} values computed for the polluted period of 2015 (Fig. 7) exhibit means, medians, and percentiles that are more negative than those of the clean and transitional periods of the same year. While mean and median values range from 0 to -8 W m^{-2} during the clean period, the values associated with the polluted scenario lie between approximately -15 and -30 W m^{-2} . An important observation is that the mean IRF_{daily} values for $f_c = 100\%$ and for the measured cloud fraction at T3 are nearly identical under cleaner atmospheric conditions, which is not the case for the polluted scenario. This is noteworthy




Figure 6. Boxplots, means and medians of IRF_{daily} calculated according to the first clean atmosphere reference for 2014 and separated according to the different aerosol availability at T3. The polluted period was divided into two subintervals due to the significant drop in concentrations of $\beta_{sct.}$, Org. and CCN shown in Fig. 1.

thy because—even with elevated aerosol concentrations and reduced cloud cover during polluted periods—the instantaneous
 355 radiative forcing associated with the few existing clouds is more negative than that produced by the more numerous clouds in the clean season.




Figure 7. Boxplots, means and medians of IRF_{daily} calculated according to the first clean atmosphere reference for 2015 and separated according to the aerosol availability at T3.

The boxplots discussed above were complemented by histograms of the annual IRF_{daily} distributions for 2014 and 2015, shown in Figs. 8 and 9. The results in Fig. 8 indicate that the overall statistics for 2014 are more negative than those for 2015: the 2015 median corresponds to about 47% of that for 2014, whereas the mean represents 66% of the 2014 mean. The results scaled by the daytime mean f_c (Fig. 9) display more homogeneous distributions approaching normality, with the lower and upper quartiles of the corresponding boxplots confined to narrower intervals than those for the distributions that ignore f_c variability at T3. This outcome suggests that multiplying by the daytime mean f_c effectively selects and constrains the final dataset, removing extreme values and producing probability density functions with reduced kurtosis magnitude.




Figure 8. Boxplots and histograms of IRF_{daily} for 2014 (left) and 2015 (right) for $f_c = \pm 100\%$.




Figure 9. Boxplots and histograms of IRF_{daily} for 2014 and 2015 for f_c observed in the T3 site.

3.2 Second clean state - “~~Sazonal~~Seasonal” irradiance reference

365 The IRF_{daily} values obtained using a variable irradiance value for a clean-atmosphere reference are presented here. It is important to recall that this seasonal irradiance reference approach aimed to reduce the likelihood that the results would be

influenced by the natural atmospheric variability over T3, ~~thereby thus~~ better isolating the aerosol effect on ~~warm~~ low-level liquid clouds. As shown in Fig. 10, the IRF_{daily} values in this case fluctuate around zero, unlike the forcings computed using the fixed irradiance reference value (Fig. 5). Nevertheless, the results are, on average, nearly constant throughout 2014 and 2015, with most data points exhibiting negative values. Subtle variations can also be observed that may or may not be associated, on average, with aerosol properties.




Figure 10. IRF_{daily} calculated according to the variable reference for clean atmosphere for 2014 and 2015. The blue line shows the results for an f_c of 100%, while the gold line shows the values according to the f_c measured in T3. Vertical scale as in Fig. 5 for comparison.

The IRF_{daily} boxplots for the different aerosol-loading scenarios in 2014 and 2015 (Figs. 11 and 12, respectively) indicate that the distributions multiplied by the measured f_c at T3 are more homogeneous and less skewed than those that do not account for this parameter. The distributions for all scenarios in both years show central values (between the twenty-fifth and seventy-fifth percentiles) between -10 W m^{-2} and approximately $+8.0 \text{ W m}^{-2}$. The means and medians for the clean and polluted scenarios of the two years are negative, with greater magnitudes for the clean periods. This behavior, associated with a larger dispersion of IRF_{daily} values during cleaner conditions, can be explained—at least for the distributions that do not include the T3’s cloud fraction—by greater variability in the r_{eff} values of the modeled clouds in clean periods. For the results multiplied

by f_c , this effect may also occur, as higher f_c values during clean conditions make the means and medians even more negative
 380 than those observed during polluted periods.




Figure 11. Boxplots, means and medians of IRF_{daily} calculated for the second clean atmosphere reference for 2014 and separated according to the different aerosol availability at T3.

The results derived from the variable irradiance reference allow the derivation of annual statistics in which the first indirect aerosol effect is more clearly isolated. The resulting statistics for 2014 and 2015 (Figs. 13 and 14) show that, compared to the values obtained using the fixed irradiance reference, the IRF_{daily} magnitudes are smaller. The effect of seasonal variations in atmospheric conditions on the warm-low-level liquid cloud droplet size is illustrated by the percentage variations in the
 385 mean r_{eff} . during the clean periods of 2014 and 2015 relative to the mean r_{eff} . of the clean reference state. These differences were 21.6% and 20.7%, respectively, while the polluted periods exhibited reductions of 18.7% for both years. The percentage variations in TOA flux were 1.62% and 0.50% for the clean intervals, while for the polluted intervals, there were increases of 0.82% and 0.34%, even though LWC was assumed to be the same for both clean and polluted periods.




Figure 12. Statistics of IRF_{daily} calculated for the second clean atmosphere reference for 2015 and separated according to the different aerosol availability at T3.




Figure 13. Boxplots and histograms of IRF_{daily} for 2014 and 2015 for $f_c = \pm 100\%$ according to the second reference for a clean atmosphere.




Figure 14. Boxplots and histograms of IRF_{daily} for 2014 and 2015 for $f_c = \pm 100\%$ measured at T3, according to the second reference for a clean atmosphere.

3.3 IRFaci sensibilities for clean and polluted scenarios

390 According to Wei et al. (2020), aerosol optical depth (AOD) is widely recognized as a critical indicator to understand atmospheric physics and regional air quality due to its ability to quantify aerosol loading in the atmosphere. Considering this, one way to assess the sensitivity of daily instantaneous forcings to aerosol availability ~~during the clean period is via AOD.~~

In this study, the sensitivity of IRF_{daily} to AOD was assessed as follows: first, we calculated the daily medians of AOD for the clean and polluted scenarios of each year, and for each of these scenarios, we selected the 10 days with the lowest medians. 395 Then, all days of a given ~~year is to compute the ratio between the mean IRF_{daily} scenario~~ were used to calculate the average AOD (for the clean scenario of 2014, for example, we called this average " $T_{clean14}$ "), while another average was calculated only for the 10 cleanest days selected previously (" $t_{clean14}$ " for the clean ~~season and~~ scenario of 2014, for example). Finally, we used Eq. 12 to obtain the percentage variation of AOD ~~for the same clean period. The same procedure can then be applied to the polluted season of that year~~ within each scenario, taking " t " as the clean reference and seeking to determine by what 400 percentage the average " T " is greater than " t ". To test the hypothesis that $\Delta AOD\%_{clean} < \Delta AOD\%_{polluted}$ for a given year, we applied a bootstrap test and obtained the statistical significance (5% level) of the percentage variations for each year. The results ($IC = [-67.8, -23.4]$, $p\text{-value} = 0.00$ and $[-16.1, 21.9]$, $p\text{-value} = 0.92$ for 2014 and ~~the sensitivities obtained for the clean and polluted periods can be compared.~~ 2015, respectively) show that the hypothesis was confirmed only for 2014. As with AOD, the average IRF_{daily} values for each scenario and for the 10 cleanest days within each scenario were calculated 405 ($T_{IRF_{polluted14}}$ and $T_{IRF_{clean14}}$ for example). The difference between these averages ($\Delta IRF_{daily} = T_{IRF} - t_{IRF}$) was used to calculate sensitivity S (Eq. 13, with units of $W m^{-2} \%^{-1}$), for each scenario of each year. The final step consisted of performing

a new statistical significance test, considering the hypothesis that S in the clean scenario is greater than in the polluted scenario. The results (p-value = 0.00 and 0.39 for 2014 and 2015, respectively) for the confidence level of 5% showed that this occurred only in 2014.

410 Table 5 presents the calculated sensitivities for the clean and polluted periods of 2014 and 2015, derived from the results based on the seasonal irradiance reference while assuming $f_c = 1$. The 2014 results indicate that the sensitivity of low-warm-low-level liquid clouds to variations in aerosol load may be greater under cleaner atmospheric conditions than in already highly polluted environments.

$$\Delta AOD\% = 100 \cdot \frac{T - t}{t} \tag{12}$$

415
$$S = \frac{|\Delta IRF_{daily}|}{\Delta AOD\%} \tag{13}$$

Table 5. Percentage variations of AOD (ΔAOD) and differences of daily IRFaci, and sensitivities according to the clean and the polluted periods of 2014 and 2015. The average AOD values for the constituent days of the clean and the polluted references were equal to 0.07 and 0.20, respectively. The average IRF_{daily} for the polluted period of 2014, equal to -1.53, was calculated from the values referring to the two polluted sub-periods of 2014 (-0.25 and -2.81, respectively).

Year - Scenario	$ \text{Mean-IRF}_{daily} (W m^{-2})$	Mean-AOD	$\Delta AOD\%$ (%)	Sensitivity	t_{IRF} ($W m^{-2}$)	$ \Delta IRF $
2014 - Clean	3.79-0.09	31.26-0.06	12.12-45.5	-3.79	-10.91	7.12
2014 - Polluted	1.53-0.28	37.54-0.15	4.08-81.18	-1.64	-4.87	3.23
2015 - Clean	1.09-0.13	81.10-0.07	1.34-81.02	-1.09	-0.32	0.77
2015 - Polluted	0.79-0.59	191.04-0.34	0.41-70.96	-0.79	0.97	1.76

4 Summary and Discussion

420 The Twomey effect related to low-level liquid clouds corresponds to the increase in cloud optical depth and albedo for a constant LWC. Those increases are caused by high aerosol concentrations that, in turn, increase the number of small droplets of clouds (the total cloud droplet concentration N_d is augmented too), becoming the dispute for water inside the cloud higher and diminishing the r_{eff} of cloud droplets. The occurrence of this mechanism can be seen in Fig. 15, which shows that, for a given fixed bin of LWC, the N_d and the r_{eff} values vary differently for each IOP. For the [0.2–0.4] $g m^{-3}$ LWC interval, in particular, higher N_d values and lower r_{eff} values occur in IOP 2, as expected. The fact that this mechanism occurs in conjunction with the IRF_{daily} distribution characterized by negative medians (see Figs. 8, 9, 13, and 14) demonstrates the existence of the Twomey effect linked to the modeled low-level liquid clouds.




Figure 15. Medians of N_d , r_{eff} and LWC according to LWC bins for both IOP.

425 The instantaneous radiative forcings due to the first indirect aerosol effect obtained in this study were separated into those calculated using a constant irradiance reference and those derived from a seasonal irradiance reference. The annual means for $f_c = \pm 100\%$ in 2014 and 2015, corresponding to the fixed reference, were -14.20 $[-24.07; -3.79]$ W m^{-2} and -9.37 $[-21.92; -1.09]$ W m^{-2} , respectively. For the seasonal reference, the results were -1.78 $[-7.48; 0.89]$ W m^{-2} and -0.82 $[-4.12; -0.29]$ W m^{-2} , while the interquartile ranges (IQR) were 8.37 W m^{-2} and 3.83 W m^{-2} , significantly smaller than those obtained for the fixed-irradiance reference (20.28 and 20.83 W m^{-2}). These results demonstrate greater IRFaci variability in 2014 than in 2015 (the interannual variability of the means was -4.83 and -0.96 W m^{-2} for the first and second references, respectively) and indicate a reduced degree of uncertainty when ~~accounting for seasonal variations in the microphysical structure of the analyzed clouds~~ seasonal aerosol variations are taken into account when calculating the forcings. Given that CCN and AOD concentrations were notably high during the second half of both years, these results are consistent in magnitude with previous findings (e.g. Wall et al., 2023; Wang et al., 2023).

435 An analysis of the probability density distributions of CBH, CTH, r_{eff} , LWP_{ground} and β_{sct} for the two references showed that, despite the presence of more scattering aerosols in the second reference (medians equal to 16.8 and 4.8 M m^{-1} for the two references, respectively) and relative discrepancies in cloud height (higher for the second reference than for the first), when compared to the most polluted days of 2014 (August and September), the second reference can still be considered relatively clean.

For both references, positive $\text{IRF}_{\text{daily}}$ values were obtained. These correspond to instances in which the reference TOA upward irradiance exceeded that of the perturbed state. This does not imply that ~~low-warm-low-level liquid~~ clouds have a net warming effect on the surface. As ~~the-warm~~-cloud models were constructed from daily statistical distributions of CBH, CTH, LWC, and r_{eff} , the occurrence of positive instantaneous forcings arises from the adopted methodology itself, which
 445 sought to reflect the natural variability of daytime cloud properties. Furthermore, it should be noted that the days selected to represent the two reference states were chosen ~~based-on-on the basis of~~ aerosol availability rather than ~~warm-low-level liquid~~ cloud characteristics. Therefore, daily fluctuations in irradiances and $\text{IRF}_{\text{daily}}$ values are expected, representing the natural day-to-day variability of ~~low-warm-low-level liquid~~ cloud properties.

~~For both references, interannual differences were observed, with more intense IRF values occurring in 2014.~~ Since the first
 450 reference, obtained using a simpler and more direct procedure, assumes a fixed irradiance value for the entire GoAmazon2014/5 period, larger IRF magnitudes are expected—particularly during polluted periods. This first reference involves fewer intermediate assumptions than the second and does not require non-contaminated data from polluted seasons. However, it neglects the seasonal variability of the atmosphere’s natural conditions over the study period, which may influence the microphysics of the analyzed clouds and the resulting IRF. The variable irradiance reference, in turn, follows the seasonal atmospheric variations,
 455 reducing the likelihood that the results are biased by natural variability. Thus, although it represents a first-order approximation, the results obtained using the seasonal irradiance reference are more appropriate for analyzing instantaneous radiative forcings associated with the first indirect aerosol effect.

The differences between the IRF_{aci} values computed using the two reference states align with Gryspeerdt et al. (2023), who emphasized the importance of clean-atmosphere states in determining resultant instantaneous forcings. The slightly higher
 460 percentage variations in integrated TOA flux during clean periods compared to polluted ones highlight the influence of cleaner atmospheric conditions on IRF_{aci} calculations. The results in Table 5 further indicate that ~~low-warm-low-level liquid~~ clouds may exhibit greater sensitivity to anthropogenic aerosols under cleaner conditions than in atmospheres already affected by high ~~pollution levels~~ levels of pollution.

The daily IRF_{aci} values obtained when considering f_c variations at T3 and using the seasonal irradiance reference were
 465 $-1.63 [-4.87; 0.65] \text{ W m}^{-2}$ and $-1.01 [-2.52; -0.23] \text{ W m}^{-2}$ for 2014 and 2015, respectively. The IRF_{aci} values associated with the first indirect effect during the GoAmazon2014/5 campaign (averaged from the $\text{IRF}_{\text{daily}}$ means of 2014 and 2015, for $f_c = +100\%$) corresponding to the two reference approaches were $-11.79 [-23.0; -2.44] \text{ W m}^{-2}$ and $-1.30 [-5.8; 0.59] \text{ W m}^{-2}$, respectively. Despite differing observational conditions (experimental platform and reference irradiance), both sets of results fall within the range reported by Herbert and Stier (2023), namely total instantaneous forcings—largely due to biomass-burning
 470 aerosols—of up to -50 W m^{-2} for $\text{AOD} \approx 1$.

Although direct comparison must be interpreted with caution, the second clean-atmosphere reference yielded IRF_{aci} values ~~closer in order of magnitude~~ to the estimate ($-0.7 \pm 0.5 \text{ W m}^{-2}$, specifically related to the Twomey effect) reported in the latest IPCC Assessment (Forster et al., 2021). This underscores the importance of defining an appropriate atmospheric reference state in IRF_{aci} calculations, which are significantly influenced by the chosen irradiance baseline.

Quantification of IRFaci associated with ~~warm-low-level liquid~~ clouds in the Amazon was achieved by evaluating the first indirect aerosol effect in the region, using an unique approach that combines in situ measurements, ground-based observations, and cloud modeling based on these datasets.

480 The IRFaci values calculated using the fixed irradiance reference became increasingly negative as polluted scenarios approached. The averages computed for $f_c = 100\%$ and for f_c measured at T3 are virtually identical under clean atmospheric conditions, which is not the case for polluted scenarios. This indicates that, although cloud cover is greater during cleaner periods, the influence of cloud cover on the amount of radiation reflected to space becomes more significant under polluted conditions.

485 Using ~~of~~ the seasonal irradiance reference reduced the variance of the resulting IRFaci, which oscillated around zero—unlike the forcings derived from the fixed irradiance reference. In addition, the IRFaci values obtained with the seasonal reference exhibited distributions with higher positive skewness, predominated by values smaller than the negative mean. The resulting interquartile range was also smaller, indicating reduced uncertainties.

490 The results demonstrate greater IRFaci variability in 2014 than in 2015. This interannual variability depends on the aerosol load in the atmosphere, the occurrence of which varies significantly each year, depending on meteorological factors such as the occurrence of droughts or social and economic pressures, and also on public policy decisions. For the generalization of the approach used in this work to other periods, it is therefore fundamental to adequately record the daily and seasonal variations in the properties of aerosols and clouds.

495 The result obtained for GoAmazon2014/5 (-11.8 W m^{-2} when $f_c = \del{+100\%} for the first clean-atmosphere reference) ~~presents is~~ an order of magnitude consistent with previous studies. The value obtained using the second reference (-1.3 W m^{-2}) is ~~closer to~~ of an order of magnitude consistent with the latest estimate reported by the IPCC. The percentage variations of AOD and associated IRFaci sensitivities for ~~the~~ clean and polluted periods of 2014 (~~31.3% and 37.5%; 12.1 and 4.1 per %~~) and 2015 (~~81.1% and 191.0%; 1.3 and 0.4~~45.5% and 81.2%; 15.6 and 4.0 W m^{-2} per %) suggest that the sensitivity of ~~low-warm~~ low-level liquid clouds to anthropogenic aerosols may be greater under clean conditions than in atmospheres already affected by high ~~pollution-levels~~ levels of pollution.$

500 The results obtained stem from an effort to estimate instantaneous radiative forcings entirely from in situ and ground-based remote sensing data collected over ~~the~~ a specific region of the Brazilian Amazon. This represents a unique approach to calculating instantaneous radiative forcings associated with the first indirect effect of anthropogenic aerosols on ~~warm-low-level liquid~~ clouds in the ~~region~~ Amazon. The findings contribute to a better understanding of the first indirect effect related to the influence that anthropogenic aerosols may exert on low ~~warm-liquid~~ low-level liquid clouds in the Amazon and can also serve as a reference
505 for model evaluation and future studies aiming to quantify IRFaci or ~~ERFaci in the even~~ ERFaci in that region.

Data availability. Publicly available datasets were analyzed in this study. These data can be found at ARM: <https://adc.arm.gov/discovery/>, accessed on 20 February 2026 19:50 UTC, and HALO database: <https://halo-db.pa.op.dlr.de/mission/5>, accessed on 20 February 2026 19:50 UTC. Furthermore, a folder named "00-Data.zip" disponible at <https://doi.org/10.5281/zenodo.20610494> compiles all datasets used in this study.

510 *Author contributions.* Pugliesi, A. C. and Correia, A. L. conceptualized the study. Pugliesi, A. C. and Cecchini, M. curated the data. Pugliesi, A. C. initiated the study, developed the methodology, performed the simulations and did the investigation and formal analysis. Pugliesi, A. C. prepared the manuscript. Correia, A. L. and Cecchini, M. review the manuscript. All authors have read and agreed to the published version of the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

515 *Acknowledgements.* We thank ARM, G-1 and HALO teams for the datasets used in this study. We also thank everyone involved in organizing and carrying out the GoAmazon2014/5 campaign. Andre Cezar Pugliesi thanks the Coordination for the Improvement of Higher Education Personnel (*Coordenação de Aperfeiçoamento de Pessoal de Nível Superior*, CAPES, Brazil) and the *São Paulo* Research Foundation (FAPESP, Brazil) for research grants 88882.332852/2010-01 and 2024/17448-9, respectively. Micael A. Cecchini was funded by FAPESP grant 2022/13257-9, *Serrapilheira* grant 2211-41823. Andre Cezar Pugliesi acknowledges the use of Grammarly (<https://www.grammarly.com/>)
520 to improve the grammar of this manuscript. After using this tool, the authors reviewed and revised the text as needed and assume full responsibility for the content of the manuscript.

References

- Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, <https://doi.org/10.1126/science.245.4923.1227>, 1989.
- 525 Alexandrov, M. D., Marshak, A., Cairns, B., Lacis, A. A., and Carlson, B. E.: Automated Cloud Screening Algorithm for MFRSR Data, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2003GL019105>, 2004.
- Carslaw, K. S., Lee, L. A., Reddington, C. L., Pringle, K. J., Rap, A., Forster, P. M., Mann, G. W., Spracklen, D. V., Woodhouse, M. T., Regayre, L. A., and Pierce, J. R.: Large Contribution of Natural Aerosols to Uncertainty in Indirect Forcing, *Nature*, 503, 67–71, <https://doi.org/10.1038/nature12674>, 2013.
- 530 Cecchini, M. A., Machado, L. A. T., Comstock, J. M., Mei, F., Wang, J., Fan, J., Tomlinson, J. M., Schmid, B., Albrecht, R., Martin, S. T., and Artaxo, P.: Impacts of the Manaus Pollution Plume on the Microphysical Properties of Amazonian Warm-Phase Clouds in the Wet Season, *Atmospheric Chemistry and Physics*, 16, 7029–7041, <https://doi.org/10.5194/acp-16-7029-2016>, 2016.
- Chen, Y.-C., Christensen, M. W., Stephens, G. L., and Seinfeld, J. H.: Satellite-Based Estimate of Global Aerosol–Cloud Radiative Forcing by Marine Warm Clouds, *Nature Geoscience*, 7, 643–646, <https://doi.org/10.1038/ngeo2214>, 2014.
- 535 Christensen, M. W., Neubauer, D., Poulsen, C. A., Thomas, G. E., McGarragh, G. R., Povey, A. C., Proud, S. R., and Grainger, R. G.: Unveiling Aerosol–Cloud Interactions – Part 1: Cloud Contamination in Satellite Products Enhances the Aerosol Indirect Forcing Estimate, *Atmospheric Chemistry and Physics*, 17, 13 151–13 164, <https://doi.org/10.5194/acp-17-13151-2017>, 2017.
- Diamond, M. S., Director, H. M., Eastman, R., Possner, A., and Wood, R.: Substantial Cloud Brightening From Shipping in Subtropical Low Clouds, *AGU Advances*, 1, e2019AV000 111, <https://doi.org/10.1029/2019AV000111>, 2020.
- 540 Dong, X. and Mace, G. G.: Arctic Stratus Cloud Properties and Radiative Forcing Derived from Ground-Based Data Collected at Barrow, Alaska, *Journal of Climate*, 16, 445–461, [https://doi.org/10.1175/1520-0442\(2003\)016<0445:ASCPAR>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<0445:ASCPAR>2.0.CO;2), 2003.
- Dürr, B. and Philipona, R.: Automatic Cloud Amount Detection by Surface Longwave Downward Radiation Measurements, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2003JD004182>, 2004.
- Emde, C., Buras-Schnell, R., Kylling, A., Mayer, B., Gasteiger, J., Hamann, U., Kylling, J., Richter, B., Pause, C., Dowling, T., and Bugliaro, L.: The libRadtran Software Package for Radiative Transfer Calculations (Version 2.0.1), *Geoscientific Model Development*, 9, 1647–1672, <https://doi.org/10.5194/gmd-9-1647-2016>, 2016.
- 545 Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J. L., Frame, D., Lunt, D. J., Mauritsen, T., Palmer, M. D., Watanabe, M., Wild, M., and Zhang, H.: The Earth’s Energy Budget, Climate Feedbacks and Climate Sensitivity, Tech. rep., Cambridge University Press, <https://doi.org/10.1017/9781009157896.009>, 2021.
- 550 Gasteiger, J., Emde, C., Mayer, B., Buras, R., Buehler, S. A., and Lemke, O.: Representative Wavelengths Absorption Parameterization Applied to Satellite Channels and Spectral Bands, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 148, 99–115, <https://doi.org/10.1016/j.jqsrt.2014.06.024>, 2014.
- Gaustad, K. L., Turner, D. D., and McFarlane, S. A.: MWRRET Value-Added Product: The Retrieval of Liquid Water Path and Precipitable Water Vapor from Microwave Radiometer (MWR) Data Sets, Tech. Rep. DOE/SC-ARM/TR-081.2, U.S. Department of Energy, Atmospheric Radiation Measurement Climate Research Facility, https://www.arm.gov/publications/tech_reports/doe-sc-arm-tr-081.2.pdf, 2011.
- 555 Giles, D. M., Sinyuk, A., Sorokin, M. G., Schafer, J. S., Smirnov, A., Slutsker, I., Eck, T. F., Holben, B. N., Lewis, J. R., Campbell, J. R., Welton, E. J., Korkin, S. V., and Lyapustin, A. I.: Advancements in the Aerosol Robotic Network (AERONET) Version 3 Database

- Automated near-Real-Time Quality Control Algorithm with Improved Cloud Screening for Sun Photometer Aerosol Optical Depth (AOD) Measurements, *Atmospheric Measurement Techniques*, 12, 169–209, <https://doi.org/10.5194/amt-12-169-2019>, 2019.
- Gryspeerdt, E., Povey, A. C., Grainger, R. G., Hasekamp, O., Hsu, N. C., Mulcahy, J. P., Sayer, A. M., and Sorooshian, A.: Uncertainty in Aerosol–Cloud Radiative Forcing Is Driven by Clean Conditions, *Atmospheric Chemistry and Physics*, 23, 4115–4122, <https://doi.org/10.5194/acp-23-4115-2023>, 2023.
- Hamilton, D. S., Lee, L. A., Pringle, K. J., Reddington, C. L., Spracklen, D. V., and Carslaw, K. S.: Occurrence of Pristine Aerosol Environments on a Polluted Planet, *Proceedings of the National Academy of Sciences*, 111, 18466–18471, <https://doi.org/10.1073/pnas.1415440111>, 2014.
- Harrison, L. and Michalsky, J.: Objective Algorithms for the Retrieval of Optical Depths from Ground-Based Measurements, *Applied Optics*, 33, 5126–5132, <https://doi.org/10.1364/AO.33.005126>, 1994.
- Hasekamp, O. P., Gryspeerdt, E., and Quaas, J.: Analysis of Polarimetric Satellite Measurements Suggests Stronger Cooling Due to Aerosol–Cloud Interactions, *Nature Communications*, 10, 5405, <https://doi.org/10.1038/s41467-019-13372-2>, 2019.
- Haywood, J. and Boucher, O.: Estimates of the Direct and Indirect Radiative Forcing Due to Tropospheric Aerosols: A Review, *Reviews of Geophysics*, 38, 513–543, <https://doi.org/10.1029/1999RG000078>, 2000.
- Herbert, R. and Stier, P.: Satellite Observations of Smoke–Cloud–Radiation Interactions over the Amazon Rainforest, *Atmospheric Chemistry and Physics*, 23, 4595–4616, <https://doi.org/10.5194/acp-23-4595-2023>, 2023.
- Hess, M., Koepke, P., and Schult, I.: Optical Properties of Aerosols and Clouds: The Software Package OPAC, *Bulletin of the American Meteorological Society*, https://journals.ametsoc.org/view/journals/bams/79/5/1520-0477_1998_079_0831_opoaac_2_0_co_2.xml, 1998.
- Heyn, I., Block, K., Mülmenstädt, J., Gryspeerdt, E., Kühne, P., Salzmann, M., and Quaas, J.: Assessment of Simulated Aerosol Effective Radiative Forcings in the Terrestrial Spectrum, *Geophysical Research Letters*, 44, 1001–1007, <https://doi.org/10.1002/2016GL071975>, 2017.
- IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (Eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 Pp., Cambridge University Press, 2013.
- Keeling, R. F., Morgan, E. J., and Keeling, C. D.: Atmospheric Monthly In Situ CO₂ Data - Mauna Loa Observatory, Hawaii (Archive 2024-11-13). In Scripps CO₂ Program Data. UC San Diego Library Digital Collections., <https://doi.org/10.6075/J08W3BHW>, 2017.
- Keil, A. and Haywood, J. M.: Solar Radiative Forcing by Biomass Burning Aerosol Particles during SAFARI 2000: A Case Study Based on Measured Aerosol and Cloud Properties, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD002315>, 2003.
- King, W. D., Parkin, D. A., and Handsworth, R. J.: A Hot-Wire Liquid Water Device Having Fully Calculable Response Characteristics, *Journal of Applied Meteorology and Climatology*, 17, 1809–1813, [https://doi.org/10.1175/1520-0450\(1978\)017<1809:AHWLWD>2.0.CO;2](https://doi.org/10.1175/1520-0450(1978)017<1809:AHWLWD>2.0.CO;2), 1978.
- Kollias, P., Miller, M. A., Luke, E. P., Johnson, K. L., Clothiaux, E. E., Moran, K. P., Widener, K. B., and Albrecht, B. A.: The Atmospheric Radiation Measurement Program Cloud Profiling Radars: Second-Generation Sampling Strategies, Processing, and Cloud Data Products, *Journal of Atmospheric and Oceanic Technology*, 24, 1199–1214, <https://doi.org/10.1175/JTECH2033.1>, 2007.
- Lebsock, M. D., Stephens, G. L., and Kummerow, C.: Multisensor Satellite Observations of Aerosol Effects on Warm Clouds, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2008JD009876>, 2008.

- Lee, L. A., Reddington, C. L., and Carslaw, K. S.: On the Relationship between Aerosol Model Uncertainty and Radiative Forcing Uncertainty, *Proceedings of the National Academy of Sciences*, 113, 5820–5827, <https://doi.org/10.1073/pnas.1507050113>, 2016.
- Levitus, S., Antonov, J. I., Boyer, T. P., Baranova, O. K., Garcia, H. E., Locarnini, R. A., Mishonov, A. V., Reagan, J. R., Seidov, D., Yarosh, E. S., and Zweng, M. M.: World Ocean Heat Content and Thermosteric Sea Level Change (0–2000 m), 1955–2010, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL051106>, 2012.
- Lloyd, S.: Least Squares Quantization in PCM, *IEEE Transactions on Information Theory*, 28, 129–137, <https://doi.org/10.1109/TIT.1982.1056489>, 1982.
- Long, C. N., Ackerman, T. P., Gaustad, K. L., and Cole, J. N. S.: Estimation of Fractional Sky Cover from Broadband Shortwave Radiometer Measurements, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006475>, 2006.
- Mace, G. G., Benson, S., and Kato, S.: Cloud Radiative Forcing at the Atmospheric Radiation Measurement Program Climate Research Facility: 2. Vertical Redistribution of Radiant Energy by Clouds, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD005922>, 2006.
- Machado, L. A. T., Calheiros, A. J. P., Biscaro, T., Giangrande, S., Silva Dias, M. A. F., Cecchini, M. A., Albrecht, R., Andreae, M. O., Araujo, W. F., Artaxo, P., Borrmann, S., Braga, R., Burleyson, C., Eichholz, C. W., Fan, J., Feng, Z., Fisch, G. F., Jensen, M. P., Martin, S. T., Pöschl, U., Pöhlker, C., Pöhlker, M. L., Ribaud, J.-F., Rosenfeld, D., Saraiva, J. M. B., Schumacher, C., Thalman, R., Walter, D., and Wendisch, M.: Overview: Precipitation Characteristics and Sensitivities to Environmental Conditions during GoAmazon2014/5 and ACRIDICON-CHUVA, *Atmospheric Chemistry and Physics*, 18, 6461–6482, <https://doi.org/10.5194/acp-18-6461-2018>, 2018.
- Marsing, A., Meerkötter, R., Heller, R., Kaufmann, S., Jurkat-Witschas, T., Krämer, M., Rolf, C., and Voigt, C.: Investigating the Radiative Effect of Arctic Cirrus Measured in Situ during the Winter 2015–2016, *Atmospheric Chemistry and Physics*, 23, 587–609, <https://doi.org/10.5194/acp-23-587-2023>, 2023.
- Martin, S. T., Artaxo, P., Machado, L. a. T., Manzi, A. O., Souza, R. a. F., Schumacher, C., Wang, J., Andreae, M. O., Barbosa, H. M. J., Fan, J., Fisch, G., Goldstein, A. H., Guenther, A., Jimenez, J. L., Pöschl, U., Silva Dias, M. A., Smith, J. N., and Wendisch, M.: Introduction: Observations and Modeling of the Green Ocean Amazon (GoAmazon2014/5), *Atmospheric Chemistry and Physics*, 16, 4785–4797, <https://doi.org/10.5194/acp-16-4785-2016>, 2016.
- Martins, J. V., Marshak, A., Remer, L. A., Rosenfeld, D., Kaufman, Y. J., Fernandez-Borda, R., Koren, I., Correia, A. L., Zubko, V., and Artaxo, P.: Remote Sensing the Vertical Profile of Cloud Droplet Effective Radius, Thermodynamic Phase, and Temperature, *Atmospheric Chemistry and Physics*, 11, 9485–9501, <https://doi.org/10.5194/acp-11-9485-2011>, 2011.
- Michalsky, J. J., Schlemmer, J. A., Berkheiser, W. E., Berndt, J. L., Harrison, L. C., Laulainen, N. S., Larson, N. R., and Barnard, J. C.: Multiyear Measurements of Aerosol Optical Depth in the Atmospheric Radiation Measurement and Quantitative Links Programs, *Journal of Geophysical Research: Atmospheres*, 106, 12 099–12 107, <https://doi.org/10.1029/2001JD900096>, 2001.
- Mie, G.: Beiträge Zur Optik Trüber Medien, Speziell Kolloidaler Metallösungen, *Annalen der Physik*, 330, 377–445, <https://doi.org/10.1002/andp.19083300302>, 1908.
- Morrison, H., Curry, J. A., and Khvorostyanov, V. I.: A New Double-Moment Microphysics Parameterization for Application in Cloud and Climate Models. Part I: Description, *Journal of the Atmospheric Sciences*, 62, 1665–1677, <https://doi.org/10.1175/JAS3446.1>, 2005.
- Mülmenstädt, J. and Feingold, G.: The Radiative Forcing of Aerosol–Cloud Interactions in Liquid Clouds: Wrestling and Embracing Uncertainty, *Current Climate Change Reports*, 4, 23–40, <https://doi.org/10.1007/s40641-018-0089-y>, 2018.
- Petters, M. D. and Kreidenweis, S. M.: A Single Parameter Representation of Hygroscopic Growth and Cloud Condensation Nucleus Activity, *Atmospheric Chemistry and Physics*, 7, 1961–1971, <https://doi.org/10.5194/acp-7-1961-2007>, 2007.

- 635 Pöschl, U., Martin, S. T., Sinha, B., Chen, Q., Gunthe, S. S., Huffman, J. A., Borrmann, S., Farmer, D. K., Garland, R. M., Helas, G., Jimenez, J. L., King, S. M., Manzi, A., Mikhailov, E., Pauliquevis, T., Petters, M. D., Prenni, A. J., Roldin, P., Rose, D., Schneider, J., Su, H., Zorn, S. R., Artaxo, P., and Andreae, M. O.: Rainforest Aerosols as Biogenic Nuclei of Clouds and Precipitation in the Amazon, *Science*, 329, 1513–1516, <https://doi.org/10.1126/science.1191056>, 2010.
- Quaas, J., Ming, Y., Menon, S., Takemura, T., Wang, M., Penner, J. E., Gettelman, A., Lohmann, U., Bellouin, N., Boucher, O., Sayer, A. M., Thomas, G. E., McComiskey, A., Feingold, G., Hoose, C., Kristjánsson, J. E., Liu, X., Balkanski, Y., Donner, L. J., Ginoux, P. A., Stier, P., Grandey, B., Feichter, J., Sednev, I., Bauer, S. E., Koch, D., Grainger, R. G., Kirkevå, G. A., Iversen, T., Seland, Ø., Easter, R., Ghan, S. J., Rasch, P. J., Morrison, H., Lamarque, J.-F., Iacono, M. J., Kinne, S., and Schulz, M.: Aerosol Indirect Effects – General Circulation Model Intercomparison and Evaluation with Satellite Data, *Atmospheric Chemistry and Physics*, 9, 8697–8717, <https://doi.org/10.5194/acp-9-8697-2009>, 2009.
- 640 Reid, J. S., Hobbs, P. V., Rangno, A. L., and Hegg, D. A.: Relationships between Cloud Droplet Effective Radius, Liquid Water Content, and Droplet Concentration for Warm Clouds in Brazil Embedded in Biomass Smoke, *Journal of Geophysical Research: Atmospheres*, 104, 6145–6153, <https://doi.org/10.1029/1998JD200119>, 1999.
- Riihimäki, L., Gaustad, K., and Long, C.: Radiative Flux Analysis (RADFLUXANAL) Value-Added Product: Retrieval of Clear-Sky Broadband Radiative Fluxes and Other Derived Values, Tech. Rep. DOE/SC-ARM-TR-228, 1569477, DOE-ARM, <https://doi.org/10.2172/1569477>, 2019.
- 650 Rosenfeld, D. and Woodley, W. L.: Closing the 50-Year Circle: From Cloud Seeding to Space and Back to Climate Change through Precipitation Physics, in: *Cloud Systems, Hurricanes, and the Tropical Rainfall Measuring Mission (TRMM)*, pp. 59–80, Meteorological Monographs, American Meteorological Society, 2003.
- Santer, B. D., Painter, J. F., Bonfils, C., Mears, C. A., Solomon, S., Wigley, T. M. L., Gleckler, P. J., Schmidt, G. A., Doutriaux, C., Gillett, N. P., Taylor, K. E., Thorne, P. W., and Wentz, F. J.: Human and Natural Influences on the Changing Thermal Structure of the Atmosphere, *Proceedings of the National Academy of Sciences*, 110, 17 235–17 240, <https://doi.org/10.1073/pnas.1305332110>, 2013.
- Schmid, B., Tomlinson, J. M., Hubbe, J. M., Comstock, J. M., Mei, F., Chand, D., Pekour, M. S., Kluzek, C. D., Andrews, E., Biraud, S. C., and McFarquhar, G. M.: The DOE ARM Aerial Facility, *Bulletin of the American Meteorological Society*, 95, 723–742, <https://doi.org/10.1175/BAMS-D-13-00040.1>, 2014.
- 660 Solomon, S., Rosenlof, K. H., Portmann, R. W., Daniel, J. S., Davis, S. M., Sanford, T. J., and Plattner, G.-K.: Contributions of Stratospheric Water Vapor to Decadal Changes in the Rate of Global Warming, *Science*, 327, 1219–1223, <https://doi.org/10.1126/science.1182488>, 2010.
- Stamnes, K., Tsay, S.-C., Wiscombe, W., and Jayaweera, K.: Numerically Stable Algorithm for Discrete-Ordinate-Method Radiative Transfer in Multiple Scattering and Emitting Layered Media, *Applied Optics*, 27, 2502–2509, <https://doi.org/10.1364/AO.27.002502>, 1988.
- 665 Stephens, G. L.: Radiation Profiles in Extended Water Clouds. II: Parameterization Schemes, *Journal of the Atmospheric Sciences*, [https://doi.org/https://doi.org/10.1175/1520-0469\(1978\)035<2123:RPIEWC>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0469(1978)035<2123:RPIEWC>2.0.CO;2), 1978.
- Toll, V., Christensen, M., Quaas, J., and Bellouin, N.: Weak Average Liquid-Cloud-Water Response to Anthropogenic Aerosols, *Nature*, 572, 51–55, <https://doi.org/10.1038/s41586-019-1423-9>, 2019.
- Trenberth, K. E., Fasullo, J. T., and Balmaseda, M. A.: Earth’s Energy Imbalance, *Journal of Climate*, <https://doi.org/10.1175/JCLI-D-13-00294.1>, 2014.
- 670

- Turner, D. D., Clough, S. A., Liljegren, J. C., Clothiaux, E. E., Cady-Pereira, K. E., and Gaustad, K. L.: Retrieving Liquid Water Path and Precipitable Water Vapor From the Atmospheric Radiation Measurement (ARM) Microwave Radiometers, *IEEE Transactions on Geoscience and Remote Sensing*, 45, 3680–3690, <https://doi.org/10.1109/TGRS.2007.903703>, 2007.
- Twomey, S.: Pollution and the Planetary Albedo, *Atmospheric Environment* (1967), 8, 1251–1256, [https://doi.org/10.1016/0004-6981\(74\)90004-3](https://doi.org/10.1016/0004-6981(74)90004-3), 1974.
- Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *Journal of the Atmospheric Sciences*, 1977.
- Uin, J.: Nephelometer Instrument Handbook, Tech. Rep. DOE/SC-ARM-TR-165, U.S. Department of Energy, Atmospheric Measurement user facility, Richland, Washington, 2024.
- Uin, J. and Enekwizu, O.: Cloud Condensation Nuclei Particle Counter Instrument Handbook, Tech. Rep. DOE/SCARM-TR-168, U.S. Department of Energy, Atmospheric Measurement user facility, Richland, Washington, 2024.
- Uin, J., Aiken, A. C., Dubey, M. K., Kuang, C., Pekour, M., Salwen, C., Sedlacek, A. J., Senum, G., Smith, S., Wang, J., Watson, T. B., and Springston, S. R.: Atmospheric Radiation Measurement (ARM) Aerosol Observing Systems (AOS) for Surface-Based In Situ Atmospheric Aerosol and Trace Gas Measurements, *Journal of Atmospheric and Oceanic Technology*, 36, 2429–2447, <https://doi.org/10.1175/JTECH-D-19-0077.1>, 2019.
- Wall, C. J., Storelvmo, T., and Possner, A.: Global Observations of Aerosol Indirect Effects from Marine Liquid Clouds, *Atmospheric Chemistry and Physics*, 23, 13 125–13 141, <https://doi.org/10.5194/acp-23-13125-2023>, 2023.
- Wang, M., Su, J., Xu, Y., Han, X., Peng, N., and Ge, J.: Radiative Contributions of Different Cloud Types to Regional Energy Budget over the SACOL Site, *Climate Dynamics*, 61, 1697–1715, <https://doi.org/10.1007/s00382-022-06651-0>, 2023.
- Wang, Y., Li, J., Fang, F., Zhang, P., He, J., Pöhlker, M. L., Henning, S., Tang, C., Jia, H., Wang, Y., Jian, B., Shi, J., and Huang, J.: In-Situ Observations Reveal Weak Hygroscopicity in the Southern Tibetan Plateau: Implications for Aerosol Activation and Indirect Effects, *npj Climate and Atmospheric Science*, 7, 1–11, <https://doi.org/10.1038/s41612-024-00629-x>, 2024.
- Watson, T. B.: Aerosol Chemical Speciation Monitor (ACSM) Instrument Handbook, Tech. Rep. DOE/SC-ARM-TR-196, 1375336, DOE-ARM, <https://doi.org/10.2172/1375336>, 2017.
- Wei, X., Chang, N.-B., Bai, K., and Gao, W.: Satellite Remote Sensing of Aerosol Optical Depth: Advances, Challenges, and Perspectives, *Critical Reviews in Environmental Science and Technology*, 50, 1640–1725, <https://doi.org/10.1080/10643389.2019.1665944>, 2020.
- Wendisch, M., Pöschl, U., Andreae, M. O., Machado, L. A. T., Albrecht, R., Schlager, H., Rosenfeld, D., Martin, S. T., Abdelmonem, A., Afchine, A., Araújo, A. C., Artaxo, P., Aufmhoff, H., Barbosa, H. M. J., Borrmann, S., Braga, R., Buchholz, B., Cecchini, M. A., Costa, A., Curtius, J., Dollner, M., Dorf, M., Dreiling, V., Ebert, V., Ehrlich, A., Ewald, F., Fisch, G., Fix, A., Frank, F., Fütterer, D., Heckl, C., Heidelberg, F., Hüneke, T., Jäkel, E., Järvinen, E., Jurkat, T., Kanter, S., Kästner, U., Kenntner, M., Kesselmeier, J., Klimach, T., Knecht, M., Kohl, R., Kölling, T., Krämer, M., Krüger, M., Krisna, T. C., Lavric, J. V., Longo, K., Mahnke, C., Manzi, A. O., Mayer, B., Mertes, S., Minikin, A., Molleker, S., Münch, S., Nillius, B., Pfeilsticker, K., Pöhlker, C., Roiger, A., Rose, D., Rosenow, D., Sauer, D., Schnaiter, M., Schneider, J., Schulz, C., de Souza, R. A. F., Spanu, A., Stock, P., Vila, D., Voigt, C., Walser, A., Walter, D., Weigel, R., Weinzierl, B., Werner, F., Yamasoe, M. A., Ziereis, H., Zinner, T., and Zöger, M.: ACRIDICON-CHUVA Campaign: Studying Tropical Deep Convective Clouds and Precipitation over Amazonia Using the New German Research Aircraft HALO, *Bulletin of the American Meteorological Society*, 97, 1885–1908, <https://doi.org/10.1175/BAMS-D-14-00255.1>, 2016.
- Zhao, C. and Garrett, T. J.: Effects of Arctic Haze on Surface Cloud Radiative Forcing, *Geophysical Research Letters*, 42, 557–564, <https://doi.org/10.1002/2014GL062015>, 2015.